Sensitivity to the control-sample selection

Variant	N_{FRB}	$N_{control}$	Quartile counts (face-on → edge-on)	A-Dr p_m	MW two-sided p_m	MW one-sided p_m
Sérsic-index range						
$0.4 < n < 1.5$ (adopted)	56	24,450	14,19,10,13	0.54	0.46	0.23
$0.4 < n < 2.0$	56	31,212	14,18,10,14	0.64	0.55	0.27
$0.4 < n < 2.5$	56	36,700	14,18,10,14	0.68	0.66	0.32
$n < 2.5$	56	37,670	14,18,10,14	0.69	0.63	0.31
all n (no morphology cut)	56	52,682	12,16,14,14	0.74	0.76	0.62
Magnitude limit						
control $r \leq 22$, all 56 hosts	56	24,450	14,19,10,13	0.54	0.46	0.23
control $r \leq 21$, all 56 hosts	56	7,423	12,20,10,14	0.74	0.69	0.35
both samples $r \leq 21$	45	7,423	11,17,7,10	0.44	0.34	0.17
control $r \leq 20$, all 56 hosts	56	1,426	12,20,10,14	0.84	0.88	0.44
both samples $r \leq 20$	32	1,426	9,10,6,7	0.75	0.49	0.25
Colour and star formation						
star-forming only (quiescent_flag = 0)	56	23,331	14,18,10,14	0.57	0.49	0.25
blue, $g - i < 1.2$	56	11,720	14,18,10,14	0.63	0.52	0.26
red, $g - i \geq 1.2$	56	12,730	14,19,10,13	0.45	0.39	0.20

p_m: fraction of 10,000 random control samples of the same size that differ from the control at least as much as the FRB hosts.
The one-sided MW test is run in the direction of an edge-on deficit. With 10,000 draws the smallest possible p_m is 10^{-4}.
Each block varies one criterion; the others are those of the adopted control ($0.4 < n < 1.5$, $r \leq 22$, $q > 0.2$).