Inclination tests in redshift bins

Variant	N_{FRB}	$N_{control}$	Quartile counts (face-on → edge-on)	AD p_m	MW two-sided p_m	MW one-sided p_m
All, control full	55	24,450	14,19,10,12	0.44	0.33	0.16
z < 0.1, control full	12	24,450	7,0,2,3	0.11	0.21	0.10
z < 0.1, control z_phot in [0.2, 0.3)	12	4,937	7,1,1,3	0.055	0.13	0.061
z < 0.2, control full	23	24,450	9,5,4,5	0.29	0.29	0.15
z < 0.2, control z_phot in [0.2, 0.3)	23	4,937	10,5,3,5	0.12	0.11	0.056
z ≥ 0.2, control full	32	24,450	5,14,6,7	0.52	0.70	0.35
z ≥ 0.2, control z_phot in [0.2, inf)	32	24,304	5,14,6,7	0.51	0.70	0.35
0.2 ≤ z < 0.4, control full	25	24,450	3,11,4,7	0.62	0.73	0.64
0.2 ≤ z < 0.4, control z_phot in [0.2, 0.4)	25	10,473	3,11,4,7	0.70	0.95	0.53
z ≥ 0.4, control –	7	–	–	–	–	–

p_m: fraction of 10,000 random control samples of the same size that differ from the control at least as much as the FRB hosts.
The one-sided MW test is run in the direction of an edge-on deficit. With 10,000 draws the smallest possible p_m is 10^{-4}.
Matched: control restricted to the same photometric-redshift range. The z ≥ 0.4 bin (7 hosts) is too small to test.