## Headline inclination tests

| Quantity | FRB hosts | Control (HSC disks) |
|---|---|---|
| Median $\cos i$ (median $i$) | 0.626 (51.3°) | 0.575 (54.9°) |
| Fraction with $i > 60°$ | 0.30 | 0.40 |
| Quartile counts, face-on → edge-on (expected 14 each) | 14,19,10,13 | – |
| AD / MW two-sided / MW one-sided $p_m$, 56 hosts | 0.54 / 0.46 / 0.23 | – |
| Same, with measurement errors (median, 16–84%) | 0.50 (0.29–0.74) / 0.43 (0.26–0.66) / 0.21 (0.13–0.33) | – |
| Realizations with $p_m < 0.05$ (AD / MW two-sided / one-sided) | 0.0% / 0.0% / 0.4% | – |
| AD / MW one-sided $p_m$, the 23 hosts of Bhardwaj et al. (2024) remeasured | 0.93 / 0.42 | – |
| Power: AD (MW) detects the published deficit at $p < 0.05$, samples of 56 | 99.95% (99.85%) | – |
| Power: same, samples of 23 | 90.85% (88.55%) | – |

$p_m$: fraction of 10,000 random control samples of the same size that differ from the control at least as much as the FRB hosts.
The one-sided MW test is run in the direction of an edge-on deficit. With 10,000 draws the smallest possible $p_m$ is $10^{-4}$.
Power: 2,000 samples drawn with replacement from the 23 published inclinations of Bhardwaj et al. (2024).