Fits with and without the PSF

Axis ratios	N_{FRB}	Median i	Quartile counts	AD p_m	MW two-sided p_m	MW one-sided p_m
GALFIT, PSF-convolved (adopted)	56	51.3°	14,19,10,13	0.54	0.46	0.23
AstroPhot, PSF-convolved	56	53.2°	13,18,8,17	0.44	0.99	0.50
GALFIT, no PSF	56	38.0°	32,13,6,5	$\leq 10^{-4}$	$\leq 10^{-4}$	$\leq 10^{-4}$
AstroPhot, no PSF	56	38.1°	31,14,6,5	$\leq 10^{-4}$	$\leq 10^{-4}$	$\leq 10^{-4}$
Hosts shared with Bhardwaj et al. (2024)						
Shared with Bhardwaj et al. (2024): this work	22	52.8°	5,7,4,6	0.96	0.85	0.43
Shared with Bhardwaj et al. (2024): their AutoProf values	22	42.5°	11,7,2,2	0.004	0.004	0.003

p_m: fraction of 10,000 random control samples of the same size that differ from the control at least as much as the FRB hosts.
The one-sided MW test is run in the direction of an edge-on deficit. With 10,000 draws the smallest possible p_m is 10^{-4}.
Without the PSF, seeing circularizes marginally resolved hosts and makes them look face-on; this reproduces the published deficit.