
Every inclination test: FRB hosts vs field disks (grey: no-PSF fits and published values, for contrast)
AD
MW two-sided
MW one-sided (edge-on deficit)
Adopted (GALFIT, PSF-convolved)
56 hosts vs HSC disks
Hosts of Bhardwaj et al. (2024), remeasured
23 hosts, this work
22 shared hosts, this work
Measurement errors (2,000 realizations; median, 16–84%)
q and 5° systematic redrawn
Control-sample variants
0.4 < n < 2.0
0.4 < n < 2.5
n < 2.5
all n (no morphology cut)
control r ≤ 21, all 56 hosts
both samples r ≤ 21
control r ≤ 20, all 56 hosts
both samples r ≤ 20
star-forming only (quiescent_flag = 0)
blue, g − i < 1.2
red, g − i ≥ 1.2
Intrinsic thickness q0
q0 = 0.10
q0 = 0.15
q0 = 0.20
q0 = 0.25
Host morphology subsets
Without the early-type host
Without early-type and ambiguous hosts
Reliable Sb–Irr or literature disks only
Redshift bins (full control; control matched in z_phot)
All (full)
z < 0.1 (full)
z < 0.1 (matched)
z < 0.2 (full)
z < 0.2 (matched)
z >= 0.2 (full)
z >= 0.2 (matched)
0.2 <= z < 0.4 (full)
0.2 <= z < 0.4 (matched)
For contrast: fits without the PSF, and the published inclinations
GALFIT, no PSF
AstroPhot, no PSF
Bhardwaj et al. (2024) published, 22 shared
p = 0.05
10^-4
10^-3
10^-2
10^-1
10^0
p_m (fraction of random control samples at least as different)