Correlations within tertiles: z, M*, SFR, DM_host

	low tertile	mid tertile	high tertile	all	pooled	partial τ
SFR vs M*, binned by z (N = 46)	+0.27 (0.31) AD 0.52 MC +0.29	+0.07 (0.82) AD 0.36 MC +0.20	+0.22 (0.41) AD 0.77 MC +0.15	+0.25 (0.097) AD 0.43 MC +0.23	+0.20 (0.24)	+0.15 (0.11) MC +0.14
DM_host vs M*, binned by z (N = 51)	-0.10 (0.70) AD 0.57 MC -0.13	-0.41 (0.10) AD 0.090 MC -0.25	+0.43 (0.082) AD 0.37 MC +0.27	-0.01 (0.96) AD 0.91 MC -0.04	-0.02 (0.88)	-0.01 (0.91) MC -0.03
DM_host vs SFR, binned by z (N = 46)	+0.13 (0.64) AD 0.34 MC +0.04	-0.12 (0.69) AD 0.84 MC -0.01	**+0.82 (< 0.001)** **AD 0.009** **MC +0.57**	+0.24 (0.10) AD 0.020 MC +0.17	**+0.40 (0.013)**	+0.19 (0.073) MC +0.13
SFR vs z, binned by M* (N = 46)	+0.33 (0.25) AD 0.11 MC +0.35	+0.17 (0.53) AD 0.59 MC +0.26	+0.31 (0.24) AD 1.00 MC +0.30	**+0.34 (0.020)** **AD 0.10** **MC +0.35**	+0.27 (0.10)	+0.20 (0.089) MC +0.21
DM_host vs z, binned by M* (N = 51)	-0.05 (0.85) AD 0.38 MC -0.05	+0.00 (1.00) AD 0.95 MC -0.09	+0.35 (0.15) AD 0.71 MC +0.23	+0.06 (0.66) AD 0.95 MC +0.01	+0.11 (0.47)	+0.04 (0.69) MC +0.02
DM_host vs SFR, binned by M* (N = 46)	-0.00 (0.99) AD 0.33 MC -0.10	**+0.70 (0.002)** **AD 0.009** **MC +0.42**	+0.31 (0.24) AD 0.44 MC +0.31	+0.24 (0.10) AD 0.023 MC +0.17	**+0.40 (0.013)**	+0.19 (0.068) MC +0.12
M* vs z, binned by SFR (N = 46)	+0.25 (0.36) AD 0.40 MC +0.27	+0.07 (0.81) AD 0.90 MC +0.16	+0.28 (0.31) AD 0.45 MC +0.19	+0.23 (0.12) AD 0.057 MC +0.23	+0.20 (0.23)	+0.12 (0.20) MC +0.12
DM_host vs z, binned by SFR (N = 46)	+0.11 (0.69) AD 0.85 MC +0.00	-0.48 (0.071) AD 0.070 MC -0.32	+0.37 (0.17) AD 0.81 MC +0.34	+0.00 (0.98) AD 0.99 MC -0.02	-0.00 (0.99)	-0.05 (0.62) MC -0.05
DM_host vs M*, binned by SFR (N = 46)	-0.30 (0.26) AD 0.15 MC -0.21	-0.02 (0.95) AD 0.70 MC -0.15	+0.14 (0.63) AD 0.71 MC +0.10	-0.04 (0.78) AD 0.96 MC -0.05	-0.07 (0.68)	-0.06 (0.59) MC -0.06
M* vs z, binned by DM_host (N = 51)	+0.11 (0.66) AD 0.77 MC +0.23	**+0.59 (0.013)** **AD 0.002** **MC +0.27**	+0.38 (0.13) AD 0.14 MC +0.47	**+0.32 (0.024)** **AD 0.023** **MC +0.31**	**+0.38 (0.012)**	**+0.22 (0.034)** **MC +0.21**
SFR vs z, binned by DM_host (N = 46)	**+0.56 (0.030)** **AD 0.13** **MC +0.39**	-0.16 (0.55) AD 0.72 MC +0.11	**+0.60 (0.019)** **AD 0.15** **MC +0.56**	**+0.34 (0.020)** **AD 0.097** **MC +0.35**	**+0.36 (0.028)**	**+0.23 (0.025)** **MC +0.23**
SFR vs M*, binned by DM_host (N = 46)	+0.11 (0.69) AD 0.45 MC +0.19	+0.25 (0.36) AD 0.64 MC +0.12	+0.32 (0.25) AD 0.80 MC +0.41	+0.25 (0.097) AD 0.41 MC +0.23	+0.23 (0.17)	+0.19 (0.096) MC +0.17

Cells: Spearman r_S (p) for the tertiles, all hosts and the Fisher-pooled within-tertile value; Kendall partial τ given the binning quantity.
AD: two-sample Anderson–Darling p that the y quantity has the same distribution below and above the median of the x quantity.
MC: median r_S over 2,000 draws through the measurement errors with tertiles re-formed. Bold: Spearman p < 0.05. Black frame: Benjamini–Hochberg q < 0.05 over the tertiles of the grid.