## Correlations with inclination within tertiles of z, M*, SFR, DM_host

| | low tertile | mid tertile | high tertile | all | pooled | partial τ |
|---|---|---|---|---|---|---|
| z vs i, binned by z (N = 55) | +0.14 (0.57)<br>AD 0.83<br>MC +0.12 | +0.36 (0.14)<br>AD 0.46<br>MC +0.32 | -0.24 (0.35)<br>AD 0.49<br>MC -0.21 | -0.02 (0.86)<br>AD 0.53<br>MC -0.04 | +0.09 (0.54) | n/a |
| M* vs i, binned by z (N = 51) | +0.10 (0.72)<br>AD 0.73<br>MC +0.08 | +0.21 (0.41)<br>AD 0.77<br>MC +0.14 | +0.15 (0.57)<br>AD 0.79<br>MC +0.20 | +0.14 (0.31)<br>AD 0.99<br>MC +0.13 | +0.15 (0.33) | +0.11 (0.21)<br>MC +0.09 |
| SFR vs i, binned by z (N = 46) | -0.39 (0.14)<br>AD 0.24<br>MC -0.35 | +0.38 (0.19)<br>AD 0.13<br>MC +0.31 | -0.12 (0.65)<br>AD 0.66<br>MC -0.19 | -0.12 (0.42)<br>AD 0.46<br>MC -0.14 | -0.07 (0.68) | -0.06 (0.56)<br>MC -0.09 |
| DM_host vs i, binned by z (N = 55) | +0.16 (0.50)<br>AD 0.27<br>MC +0.14 | -0.19 (0.44)<br>AD 0.56<br>MC -0.13 | +0.02 (0.95)<br>AD 0.70<br>MC -0.05 | +0.06 (0.64)<br>AD 0.19<br>MC +0.02 | -0.00 (0.99) | +0.05 (0.64)<br>MC +0.01 |
| z vs i, binned by M* (N = 51) | -0.05 (0.86)<br>AD 0.96<br>MC -0.06 | -0.06 (0.83)<br>AD 0.84<br>MC -0.10 | -0.24 (0.34)<br>AD 0.89<br>MC -0.10 | -0.04 (0.76)<br>AD 0.57<br>MC -0.05 | -0.12 (0.45) | -0.05 (0.61)<br>MC -0.05 |
| M* vs i, binned by M* (N = 52) | +0.06 (0.81)<br>AD 0.91<br>MC -0.10 | +0.18 (0.47)<br>AD 0.40<br>MC +0.16 | -0.16 (0.52)<br>AD 0.95<br>MC -0.04 | +0.11 (0.43)<br>AD 1.00<br>MC +0.10 | +0.03 (0.86) | n/a |
| SFR vs i, binned by M* (N = 46) | **-0.64 (0.013)**<br>**AD 0.22**<br>**MC -0.58** | +0.24 (0.37)<br>AD 0.071<br>MC +0.13 | -0.11 (0.69)<br>AD 0.35<br>MC -0.03 | -0.12 (0.42)<br>AD 0.46<br>MC -0.14 | -0.18 (0.29) | -0.08 (0.36)<br>MC -0.11 |
| DM_host vs i, binned by M* (N = 51) | -0.06 (0.82)<br>AD 0.64<br>MC -0.02 | +0.28 (0.29)<br>AD 0.10<br>MC +0.29 | +0.07 (0.78)<br>AD 0.54<br>MC -0.07 | +0.12 (0.39)<br>AD 0.043<br>MC +0.06 | +0.09 (0.55) | +0.09 (0.38)<br>MC +0.05 |
| z vs i, binned by SFR (N = 46) | -0.05 (0.84)<br>AD 0.85<br>MC -0.16 | +0.03 (0.91)<br>AD 0.86<br>MC +0.14 | -0.03 (0.92)<br>AD 0.73<br>MC -0.04 | -0.07 (0.65)<br>AD 0.75<br>MC -0.07 | -0.02 (0.92) | -0.03 (0.75)<br>MC -0.02 |
| M* vs i, binned by SFR (N = 46) | -0.08 (0.77)<br>AD 0.84<br>MC -0.13 | -0.14 (0.62)<br>AD 0.92<br>MC +0.05 | +0.32 (0.24)<br>AD 0.42<br>MC +0.32 | +0.10 (0.52)<br>AD 0.99<br>MC +0.09 | +0.03 (0.84) | +0.09 (0.38)<br>MC +0.08 |
| SFR vs i, binned by SFR (N = 46) | -0.44 (0.087)<br>AD 0.042<br>MC -0.27 | +0.10 (0.72)<br>AD 0.73<br>MC +0.04 | -0.29 (0.29)<br>AD 0.35<br>MC -0.35 | -0.12 (0.42)<br>AD 0.46<br>MC -0.14 | -0.23 (0.17) | n/a |
| DM_host vs i, binned by SFR (N = 46) | +0.03 (0.91)<br>AD 0.53<br>MC +0.10 | +0.37 (0.18)<br>AD 0.42<br>MC +0.11 | +0.07 (0.81)<br>AD 0.63<br>MC -0.02 | +0.12 (0.42)<br>AD 0.079<br>MC +0.07 | +0.16 (0.35) | +0.10 (0.28)<br>MC +0.06 |
| z vs i, binned by DM_host (N = 55) | +0.05 (0.83)<br>AD 0.41<br>MC +0.01 | -0.11 (0.67)<br>AD 0.28<br>MC +0.02 | -0.14 (0.58)<br>AD 0.47<br>MC -0.19 | -0.02 (0.86)<br>AD 0.53<br>MC -0.04 | -0.06 (0.67) | -0.02 (0.86)<br>MC -0.02 |
| M* vs i, binned by DM_host (N = 51) | -0.06 (0.83)<br>AD 0.84<br>MC +0.05 | **+0.53 (0.027)**<br>**AD 0.034**<br>**MC +0.39** | -0.04 (0.89)<br>AD 0.56<br>MC -0.01 | +0.14 (0.31)<br>AD 0.99<br>MC +0.13 | +0.17 (0.29) | +0.10 (0.27)<br>MC +0.09 |
| SFR vs i, binned by DM_host (N = 46) | -0.41 (0.13)<br>AD 0.28<br>MC -0.16 | +0.18 (0.51)<br>AD 0.56<br>MC -0.13 | -0.35 (0.20)<br>AD 0.73<br>MC -0.25 | -0.12 (0.42)<br>AD 0.46<br>MC -0.14 | -0.19 (0.25) | -0.09 (0.44)<br>MC -0.10 |
| DM_host vs i, binned by DM_host (N = 55) | -0.08 (0.76)<br>AD 0.67<br>MC -0.01 | +0.28 (0.24)<br>AD 0.34<br>MC +0.04 | -0.22 (0.39)<br>AD 0.76<br>MC -0.19 | +0.06 (0.64)<br>AD 0.20<br>MC +0.02 | +0.00 (0.98) | n/a |

Cells: Spearman $r_S$ (p) for the tertiles, all hosts and the Fisher-pooled within-tertile value; Kendall partial $\tau$ given the binning quantity.
AD: two-sample Anderson–Darling p that the y quantity has the same distribution below and above the median of the x quantity.
MC: median $r_S$ over 2,000 draws through the measurement errors with tertiles re-formed. Bold: Spearman p < 0.05. Black frame: Benjamini–Hochberg q < 0.05 over the tertiles of the grid.