## Sensitivity to the control-sample selection

| Variant | $N_{FRB}$ | $N_{control}$ | Quartile counts (face-on → edge-on) | A-Dr $p_m$ | MW two-sided $p_m$ | MW one-sided $p_m$ |
|---|---|---|---|---|---|---|
| ***Sérsic-index range*** | | | | | | |
| $0.4 < n < 1.5$ (adopted) | 56 | 24,450 | 14,19,10,13 | 0.54 | 0.46 | 0.23 |
| $0.4 < n < 2.0$ | 56 | 31,212 | 14,18,10,14 | 0.64 | 0.55 | 0.27 |
| $0.4 < n < 2.5$ | 56 | 36,700 | 14,18,10,14 | 0.68 | 0.66 | 0.32 |
| $n < 2.5$ | 56 | 37,670 | 14,18,10,14 | 0.69 | 0.63 | 0.31 |
| all $n$ (no morphology cut) | 56 | 52,682 | 12,16,14,14 | 0.74 | 0.76 | 0.62 |
| ***Magnitude limit*** | | | | | | |
| control $r \leq 22$, all 56 hosts | 56 | 24,450 | 14,19,10,13 | 0.54 | 0.46 | 0.23 |
| control $r \leq 21$, all 56 hosts | 56 | 7,423 | 12,20,10,14 | 0.74 | 0.69 | 0.35 |
| both samples $r \leq 21$ | 45 | 7,423 | 11,17,7,10 | 0.44 | 0.34 | 0.17 |
| control $r \leq 20$, all 56 hosts | 56 | 1,426 | 12,20,10,14 | 0.84 | 0.88 | 0.44 |
| both samples $r \leq 20$ | 32 | 1,426 | 9,10,6,7 | 0.75 | 0.49 | 0.25 |
| ***Colour and star formation*** | | | | | | |
| star-forming only (quiescent_flag = 0) | 56 | 23,331 | 14,18,10,14 | 0.57 | 0.49 | 0.25 |
| blue, $g - i < 1.2$ | 56 | 11,720 | 14,18,10,14 | 0.63 | 0.52 | 0.26 |
| red, $g - i \geq 1.2$ | 56 | 12,730 | 14,19,10,13 | 0.45 | 0.39 | 0.20 |

$p_m$: fraction of 10,000 random control samples of the same size that differ from the control at least as much as the FRB hosts.
The one-sided MW test is run in the direction of an edge-on deficit. With 10,000 draws the smallest possible $p_m$ is $10^{-4}$.
Each block varies one criterion; the others are those of the adopted control ($0.4 < n < 1.5$, $r \leq 22$, $q > 0.2$).