Correlations with inclination within tertiles of z, M*, SFR, DM_host

	low tertile	mid tertile	high tertile	all	pooled	partial τ
z vs i, binned by z (N = 55)	+0.14 (0.57) AD 0.83 MC +0.12	+0.36 (0.14) AD 0.46 MC +0.32	-0.24 (0.35) AD 0.49 MC -0.21	-0.02 (0.86) AD 0.53 MC -0.04	+0.09 (0.54)	n/a
M* vs i, binned by z (N = 51)	+0.10 (0.72) AD 0.73 MC +0.08	+0.21 (0.41) AD 0.77 MC +0.14	+0.15 (0.57) AD 0.79 MC +0.20	+0.14 (0.31) AD 0.99 MC +0.13	+0.15 (0.33)	+0.11 (0.21) MC +0.09
SFR vs i, binned by z (N = 46)	-0.39 (0.14) AD 0.24 MC -0.35	+0.38 (0.19) AD 0.13 MC +0.31	-0.12 (0.65) AD 0.66 MC -0.19	-0.12 (0.42) AD 0.46 MC -0.14	-0.07 (0.68)	-0.06 (0.56) MC -0.09
DM_host vs i, binned by z (N = 55)	+0.16 (0.50) AD 0.27 MC +0.14	-0.19 (0.44) AD 0.56 MC -0.13	+0.02 (0.95) AD 0.70 MC -0.05	+0.06 (0.64) AD 0.19 MC +0.02	-0.00 (0.99)	+0.05 (0.64) MC +0.01
z vs i, binned by M* (N = 51)	-0.05 (0.86) AD 0.96 MC -0.06	-0.06 (0.83) AD 0.84 MC -0.10	-0.24 (0.34) AD 0.89 MC -0.10	-0.04 (0.76) AD 0.57 MC -0.05	-0.12 (0.45)	-0.05 (0.61) MC -0.05
M* vs i, binned by M* (N = 52)	+0.06 (0.81) AD 0.91 MC -0.10	+0.18 (0.47) AD 0.40 MC +0.16	-0.16 (0.52) AD 0.95 MC -0.04	+0.11 (0.43) AD 1.00 MC +0.10	+0.03 (0.86)	n/a
SFR vs i, binned by M* (N = 46)	**-0.64 (0.013)** **AD 0.22** **MC -0.58**	+0.24 (0.37) AD 0.071 MC +0.13	-0.11 (0.69) AD 0.35 MC -0.03	-0.12 (0.42) AD 0.46 MC -0.14	-0.18 (0.29)	-0.08 (0.36) MC -0.11
DM_host vs i, binned by M* (N = 51)	-0.06 (0.82) AD 0.64 MC -0.02	+0.28 (0.29) AD 0.10 MC +0.29	+0.07 (0.78) AD 0.54 MC -0.07	+0.12 (0.39) AD 0.043 MC +0.06	+0.09 (0.55)	+0.09 (0.38) MC +0.05
z vs i, binned by SFR (N = 46)	-0.05 (0.84) AD 0.85 MC -0.16	+0.03 (0.91) AD 0.86 MC +0.14	-0.03 (0.92) AD 0.73 MC -0.04	-0.07 (0.65) AD 0.75 MC -0.07	-0.02 (0.92)	-0.03 (0.75) MC -0.02
M* vs i, binned by SFR (N = 46)	-0.08 (0.77) AD 0.84 MC -0.13	-0.14 (0.62) AD 0.92 MC +0.05	+0.32 (0.24) AD 0.42 MC +0.32	+0.10 (0.52) AD 0.99 MC +0.09	+0.03 (0.84)	+0.09 (0.38) MC +0.08
SFR vs i, binned by SFR (N = 46)	-0.44 (0.087) AD 0.042 MC -0.27	+0.10 (0.72) AD 0.73 MC +0.04	-0.29 (0.29) AD 0.35 MC -0.35	-0.12 (0.42) AD 0.46 MC -0.14	-0.23 (0.17)	n/a
DM_host vs i, binned by SFR (N = 46)	+0.03 (0.91) AD 0.53 MC +0.10	+0.37 (0.18) AD 0.42 MC +0.11	+0.07 (0.81) AD 0.63 MC -0.02	+0.12 (0.42) AD 0.079 MC +0.07	+0.16 (0.35)	+0.10 (0.28) MC +0.06
z vs i, binned by DM_host (N = 55)	+0.05 (0.83) AD 0.41 MC +0.01	-0.11 (0.67) AD 0.28 MC +0.02	-0.14 (0.58) AD 0.47 MC -0.19	-0.02 (0.86) AD 0.53 MC -0.04	-0.06 (0.67)	-0.02 (0.86) MC -0.02
M* vs i, binned by DM_host (N = 51)	-0.06 (0.83) AD 0.84 MC +0.05	**+0.53 (0.027)** **AD 0.034** **MC +0.39**	-0.04 (0.89) AD 0.56 MC -0.01	+0.14 (0.31) AD 0.99 MC +0.13	+0.17 (0.29)	+0.10 (0.27) MC +0.09
SFR vs i, binned by DM_host (N = 46)	-0.41 (0.13) AD 0.28 MC -0.16	+0.18 (0.51) AD 0.56 MC -0.13	-0.35 (0.20) AD 0.73 MC -0.25	-0.12 (0.42) AD 0.46 MC -0.14	-0.19 (0.25)	-0.09 (0.44) MC -0.10
DM_host vs i, binned by DM_host (N = 55)	-0.08 (0.76) AD 0.67 MC -0.01	+0.28 (0.24) AD 0.34 MC +0.04	-0.22 (0.39) AD 0.76 MC -0.19	+0.06 (0.64) AD 0.20 MC +0.02	+0.00 (0.98)	n/a

Cells: Spearman r_S (p) for the tertiles, all hosts and the Fisher-pooled within-tertile value; Kendall partial τ given the binning quantity.
AD: two-sample Anderson–Darling p that the y quantity has the same distribution below and above the median of the x quantity.
MC: median r_S over 2,000 draws through the measurement errors with tertiles re-formed. Bold: Spearman p < 0.05. Black frame: Benjamini–Hochberg q < 0.05 over the tertiles of the grid.