The Data
 Correlation
 SLR
 Inference for SLR

 000000
 000000000
 0000000000
 00000000000

### Correlation and Simple Linear Regression

Shannon Pileggi

STAT 217

1 / 44

The Data

**OUTLINE** 

STAT 217: Unit 3 Deck 3

The Data

Correlation

Simple Linear Regression

Inference for SLR

Inference for SLR

2 / 44

 The Data
 Correlation
 SLR
 Inference for SLR

 ●000000
 0000000000
 0000000000
 0000000000

# Chocolate Consumption, Cognitive Function, and Nobel Laureates by Franz H. Messerli, M.D.

New England Journal of Medicine 367(16), Oct 18, 2012

STAT 217: Unit 3 Deck 3

- chocolate contains flavanols, which are loosely linked to improved cognitive function
- ▶ Is there a relationship between country's level of chocolate consumption and its population's cognitive function?
- explanatory variable: per capita yearly chocolate consumption (in kg)
- response variable: total number of Nobel laureates per 10 million person (through 2011)


 The Data
 Correlation
 SLR
 Inference for SLR

 0 ● 00000
 0000000000
 0000000000
 00000000000

## Chocolate Consumption, Cognitive Function, and Nobel Laureates by Franz H. Messerli, M.D.

New England Journal of Medicine 367(16), Oct 18, 2012

Correlation



4□ ► 4□ ► 4□ ► 4□ ► 4□ ► 9 Q C 3 / 44

 The Data
 Correlation
 SLR
 Inference for SLR

 ○○●○○○○
 ○○○○○○○○○
 ○○○○○○○○○○

#### Discussion

- ▶ Is this study observational or experimental?
- What is the unit of observation?
- ► Describe the sample:
- ▶ Describe the population:
- ► What are other variables may affect the relationship between chocolate consumption and nobel laureates?

 STAT 217: Unit 3 Deck 3
 5 / 44

#### In the media

- ▶ Time: Secret to Winning a Nobel Prize? Eat more Chocolate
- ▶ MSN: Want to win a Nobel Prize? Eat more chocolate
- ► Forbes: Chocolate and Nobel Prizes Linked in Study
- ▶ USA Today: Study links eating chocolate to winning Nobels
- Reuters: Eat chocolate, win the Nobel Prize?
- ▶ NPR: The Secret to Genius? It Might Be More Chocolate

 The Data
 Correlation
 SLR
 Inference for SLR


 000●000
 000000000
 0000000000
 00000000000

### Discussion, continued

#### Other oddities:

- ▶ Does it make sense to use nobel laureates as a measure of a country's cognitive function?
- ▶ Do we even know if the nobel laureates themselves ate chocolate?
- ▶ Were the variables measured on the same temporal scale?
- ▶ How are nobel prize winners identified with a specific country?

Stats.org: Cacao or Caca? How the media bit into chocolate Nobel prize link



 The Data
 Correlation
 SLR
 Inference for SLR

 00000●0
 000000000
 0000000000
 00000000000

### From the author (Messerli)


#### Reuters:

"I started plotting this in a hotel room in Kathmandu, because I had nothing else to do, and I could not believe my eyes," he told Reuters Health. All the countries lined up neatly on a graph, with higher chocolate intake tied to more laureates. The link was so strong it made a joke out of a statistic that virtually all studies in medical journals hinge on - the so-called p-value.


#### NPR:

"I have published about 800 papers in peer-reviewed journals," he says, "and every single one of them stands and falls with the p-value. And now here I find a p-value of 0.0001, and this is, to my way of thinking, a completely nonsensical relation. Unless you or anybody else can come up with an explanation. I've presented it to a few of my colleagues, and nobody has any thoughts."

STAT 217: Unit 3 Deck 3 8 / 4









4 中 7 4 周 7 4 夏 7 4 夏 7

11 / 44

increase



### Correlation examples



#### Rules of thumb:

- ightharpoonup 0 < |r| < 0.3 weak correlation
- ightharpoonup 0.3 < |r| < 0.7 moderate correlation
- ightharpoonup 0.7 < |r| < 1.0 strong correlation

 ★ロト ◆ラト ◆恵ト ◆恵ト ◆恵 ◆つへで

 STAT 217: Unit 3 Deck 3

 13 / 44

 The Data
 Correlation
 SLR
 Inference for SLR

 000000
 000000000
 0000000000
 0000000000


#### Which scenario illustrates the strongest linear association?

- 1. College GPA and high school GPA, r = 0.46
- 2. price and quality rating of bike helmets, r = 0.30
- 3. years of education and years in jail, r = -0.53
- 4. New York City marathon, age and finish time, r = 0.04
- 5. college GPA and a measure of tendency to procrastinate, r=-0.36

### Correlation estimate

> plot(NEJM\$chocolate,NEJM\$nobel\_rate)
> cor(NEJM\$chocolate,NEJM\$nobel\_rate)

Γ17 0.8010949




4ロ > 4酉 > 4 喜 > 喜 めの(や

15 / 44

 The Data
 Correlation
 SLR
 Inference for SLR

 000000
 0000000000
 0000000000
 00000000000

### **Group Exercise**



## What do you think the correlation between *x* and *y* is?

1. r = 1


2. r = 0.5


3. r = 0

4. r = -1


 ★ロト ◆ 母 ト ◆ 恵 ト ◆ 恵 ト ◆ 恵 ト 恵 ト 恵 ・ ◇ Q ○

 STAT 217: Unit 3 Deck 3





STAT 217: Unit 3 Deck 3






18 / 44

 The Data
 Correlation
 SLR
 Inference for SLR

 000000
 000000000
 0000000000
 00000000000

### The idea



Estimate the line of best fit:

$$\hat{y}=b_0+b_1x$$

STAT 217: Unit 3 Deck 3

STAT 217: Unit 3 Deck 3

Correlation allows us to:

- 1. identify the direction of the association between *x* and *y*
- 2. quantify the strength of the association between *x* and *y*

A line of best fit additionally allows us to:

- 1. make predictions of y based on x
- 2. further describe the relationship between *x* and *y* with the slope *b*<sub>1</sub>



The Data Correlation SLR Inference for SLR

### Identifying the line

> lm(NEJM\$nobel\_rate~NEJM\$chocolate)

Call:

lm(formula = NEJM\$nobel\_rate ~ NEJM\$chocolate)

Coefficients:

(Intercept) NEJM\$chocolate -3.400 2.496

General form:  $\hat{y} = b_0 + b_1 x$ 


Here, this is:  $\hat{y} = b_0 + b_1 \times chocolate$ 

Estimated line:  $\hat{y} = -3.4 + 2.5x$ 

Here, this is:  $\hat{y} = -3.4 + 2.5 \times chocolate$ 

STAT 217: Unit 3 Deck 3 22 / 44

### The slope $(b_1)$



 $\hat{v} = -3.4 + 2.5x$ 


- ▶ the slope of this line is 2.5
- the slope is the amount that  $\hat{y}$  changes when x increases by one unit
- for each additional 1kg/yr/capita of chocolate that a country consumes, the rate of nobel laureates per 10 million persons increases by 2.5
- the sign of the slope (positive or negative) indicates the direction of the association

□ → ◆ □ → ◆ □ → ◆ □ → □ □ → ○ ○
 23 / 44

 The Data
 Correlation
 SLR
 Inference for SLR

 000000
 000000000
 000000000
 0000000000

### The y-intercept $(b_0)$



 $\hat{y} = -3.4 + 2.5x$ 

- ▶ the intercept of this line is -3.4
- the intercept is the predicted value of y when x = 0
- when a country consumes no chocolate, its predicted number of Nobel laureates is -3.4
- intercepts aren't always interpretable

←□ → ←□ → ← □ → ← □ → ← □ → ← □
 24 / 44

 The Data
 Correlation
 SLR
 Inference for SLR

 000000
 000000000
 000000000
 0000000000

Suppose we use literacy rate of a country (measured as a percent) to predict the life expectancy of that country (measured in years), and we estimate the regression line to be  $\hat{y} = 11.8 + 0.70x$ .

#### What is the interpretation of the slope?

- 1. For each one percent increase in literacy rate, life expectancy increases by 0.7 years.
- 2. For each one year increase in life expectancy, literacy rate increases by 0.7 percentage points.
- 3. For each one percent increase in literacy rate, life expectancy increases by 11.8 years.
- 4. For each one year increase in life expectancy, literacy rate increases by 11.8 percentage points.
- 5. The predicted life expectancy of a country with a literacy rate of 0 is 11.8 years.
- 6. The predicted literacy rate of a country with a life expectancy of 0 is 11.8%.

STAT 217: Unit 3 Deck 3

✓) Q (~25 / 44


27 / 44

STAT 217: Unit 3 Deck 3

 The Data
 Correlation
 SLR

 0000000
 000000000
 000000●00000

### Prediction $(\hat{y})$



$$\hat{v} = -3.4 + 2.5x$$

- ▶ when x = 7,  $\hat{y} = -3.4 + 2.5 \times 7 = 14.1$
- ► the predicted number of Nobel Laureates for a country that consumes 7 kg/yr/capita of chocolate is 14.1

Inference for SLR

Using length of pregnancy in days (the gestation period) to predict babies' birth weight in pounds in the babies data set:

What is the correct formula to predict the birth weight of a baby with a gestation period of 300 days?

1. 
$$\hat{y} = 0.03 - 1.31 \times 300$$

2. 
$$\hat{y} = 0.03 + 0.42 \times 300$$


3. 
$$\hat{\mathbf{v}} = -1.31 + 0.03 \times 300$$

4. 
$$\hat{y} = -1.31 + 0.42 \times 300$$

 The Data
 Correlation
 SLR
 Inference for SLR

 000000
 00000000
 00000000
 000000000

### Don't extrapolate!



- Extrapolation is using a regression line to predict y-values for x-values outside the observed range of the data
- ► Extrapolation gets riskier the farther we move from the range of the given *x*-values.
- ► There is no guarantee that the relationship given by the regression equation holds outside the range of sampled *x*-values.

< □ > < ② > < 호 > < 호 > < 호 > < 호 · 28 / 44

The Data Correlation SLR Inference for SLR

### Cautions in Regression

- ▶ Don't extrapolate
- ► The analysis is not robust to outliers (can affect estimates of correlation, slope, and intercept)
- ► Correlation/association does not imply causation
- ▶ Other variables can influence the analysis through confounding and interaction - you can control for this in multivariable regression where you use more than one variable in your model to predict y

4□ ▷ ◀률 ▷ ◀ 볼 ▷ ◀ 볼 ▷ 및
 29 / 44

STAT 217: Unit 3 Deck 3

 The Data
 Correlation
 SLR
 Inference for SLR

 0000000
 000000000
 0000000000
 0000000000

When using a simple linear regression model, which of the following allows us to assess the association between x and y?

1. the intercept  $(b_0)$ 

Correlation

- 2. the slope  $(b_1)$
- 3. both

The Data

4. neither


Inference for SLR

•00000000000

 The Data
 Correlation
 SLR
 Inference for SLR

 000000
 000000000
 0000000000
 0000000000

Using MMSE score to predict whole brain volume in the ADNI data set: r = 0.19,  $\hat{y} = 796239 + 8302 \times MMSE$ 



STAT 217: Unit 3 Deck 3

What can we conclude about the strength of the association between MMSE and whole brain volume?

- 1. there is a strong association because the slope is large
- 2. there is a strong association because the correlation is large
- 3. there is a weak association because the slope is small
- 4. there is a weak association because the correlation is small

Inference for SLR

 ✓□ > ✓□ > ✓□ > ✓□ > ✓□ > ✓□ >
 № № № № №
 № №
 № №
 № №
 № №
 № №
 № №
 № №
 № №
 № №
 № №
 № №
 № №
 № №
 № №
 № №
 № №
 №
 №
 №
 №
 №
 №
 №
 №
 №
 №
 №
 №
 №
 №
 №
 №
 №
 №
 №
 №
 №
 №
 №
 №
 №
 №
 №
 №
 №
 №
 №
 №
 №
 №
 №
 №
 №
 №
 №
 №
 №
 №
 №
 №
 №
 №
 №
 №
 №
 №
 №
 №
 №
 №
 №
 №
 №
 №
 №
 №
 №
 №
 №
 №
 №
 №
 №
 №
 №
 №
 №
 №
 №
 №
 №
 №
 №
 №
 №
 №
 №
 №
 №
 №
 №
 №
 №
 №
 №
 №
 №
 №
 №
 №
 №
 №
 №

4□ > 4酉 > 4Ē > 4Ē > Ē

31 / 44

 The Data
 Correlation
 SLR
 Inference for SLR

 000000
 000000000
 0000000000
 0●000000000

#### The idea

- ▶ the *slope* of the line allows us to assess whether or not there is an association between *x* and *y*
- ▶ a slope of 0 indicates no association between x and y; a non-zero slope indicates some sort of association between x and y
- we are estimating a slope  $b_1$  based on sample data, but we want to draw conclusions about the slope in the population  $(\beta_1)$
- ► We can do this with:
  - 1. a hypothesis test of  $H_0$ :  $\beta_1 = 0$  vs  $H_a$ :  $\beta_1 \neq 0$
  - 2. a confidence interval for  $\beta_1$



STAT 217: Unit 3 Deck 3


The Data

Correlation

SLR 00000000000 Inference for SLR

STAT 217: Unit 3 Deck 3

### **Example Figures**



STAT 217: Unit 3 Deck 3

- possum data
- ▶ the spread in y (head length) is about the same for all values of x (total length)
- this satisfies "constant variability in y about the regression line"
- car data
- ▶ the relationship between x (weight) and y (miles per gallon) follows a curve rather than a straight line
- ► this violates "linear relationship between x and y"

4 □ → 4 ₱ → 4 ≣ → 4 ≣ → 9 Q (~) 35 / 44 Correlation

The Data

SLR 000000000000 Inference for SLR

4□ > 4問 > 4 = > 4 = > = 900

34 / 44

### Conditions for inference in SLR

- 1. observations are independent
- 2. linear relationship between x and y
- 3. "normality" not addressed in STAT 217 (requires residuals)
- 4. constant variability in y about the regression line





Figure (A/B/C/D) violates the independence condition. Figure (A/B/C/D) violates the linear condition. Figure (A/B/C/D) violates the constant variability condition.

4 □ ▶ 4 畳 ▶ 4 差 ▶ 4 差 ▶ 5 € か Q © nit 3 Deck 3 36 / 44

The Data Correlation SLR Inference for SLR

### Assessing conditions



Which condition may be violated? Select the *best* answer.


- 1. independence of observations
- 2. linear relationship between *x* and *y*
- 3. constant variability in *y* about the regression line
- 4. no conditions are violated

200

39 / 44

The Data Correlation SLR Inference for SLR

### Assessing conditions, continued



Which condition may be violated? Select the *best* answer.

- independence of observations
- 2. linear relationship between *x* and *y*
- 3. constant variability in *y* about the regression line
- 4. no conditions are violated

#### R results

STAT 217: Unit 3 Deck 3

- > m1<-lm(NEJM\$nobel\_rate~NEJM\$chocolate)</pre>
- > summary(m1)

#### Call:

lm(formula = NEJM\$nobel rate ~ NEJM\$chocolate)

#### Residuals:

Min 1Q Median 3Q Max -12.888 -2.953 -0.213 1.992 19.279

#### Coefficients:

Estimate Std. Error t value Pr(>|t|)

(Intercept) -3.400 2.699 -1.260 0.222 NEJM\$chocolate 2.496 0.407 6.133 4.37e-06 \*\*

---

Signif. codes: 0 \*\*\* 0.001 \*\* 0.01 \* 0.05 . 0.1 1

Residual standard error: 6.26 on 21 degrees of freedom

Multiple R-squared: 0.6418, Adjusted R-squared: 0.6247

r t value Pr(>|t|)
9 -1.260 0.222
7 6.133 4.37e-06 \*\*\*
01 \* 0.05 . 0.1 1
21 degrees of freedom

 The Data
 Correlation
 SLR
 Inference for SLR

 0000000
 000000000
 0000000000
 0000000●0000

#### Focus on the coefficients box

#### Coefficients:

STAT 217: Unit 3 Deck 3

Estimate Std. Error t value Pr(>|t|)
(Intercept) -3.400 2.699 -1.260 0.222
NEJM\$chocolate 2.496 0.407 6.133 4.37e-06 \*\*\*

On the NEJM\$chocolate line:

Hypotheses:  $H_0$ :  $\beta_1 = 0$  vs  $H_a$ :  $\beta_1 \neq 0$ 

Test statistic:  $t = \frac{\hat{\beta}_1 - 0}{se_{\hat{\beta}_1}} = \frac{2.496 - 0}{0.407} = 6.133$ 

*p*-value: two-tailed area from t distribution with df = n - 2;

p-value= 0.00000437

Conclusion: At  $\alpha = 0.05$  reject  $H_0$ ; we have evidence of a positive

association between chocolate consumption and rate of nobel laureates.

The Data Correlation

Inference for SLR 00000000000000

Correlation

> cor(babies\$bwt\_lbs,babies\$gestation)

-1.309203

babies\$gestation 0.031433 0.001931 16.274 <2e-16 \*\*\*

Inference for SLR 00000000000000

### Confidence interval for the slope

> confint(m1) 2.5 % 97.5 % (Intercept) -9.013147 2.212415 NEJM\$chocolate 1.649869 3.342646

 $\hat{\beta}_1 \pm t_{df=n-2}^* \times se_{\hat{\beta}_1}$  $2.496 + 2.080 \times 0.407$ 

- ▶ The 95% CI for  $\beta_1$  is (1.65, 3.34)
- ▶ Because this CI is entirely positive, there is a *positive* association between rate of nobel laureates and chocolate consumption, such that as chocolate consumption increases so does the rate of nobel laureates.
- ► At the 95% confidence level, for each 1kg/yr/capita increase in chocolate consumption the rate of nobel laureates per 10 million persons increases by as few as 1.65 or as many as 3.34

41 / 44

STAT 217: Unit 3 Deck 3

0.42

Coefficients:

(Intercept)

The Data

0.42

Coefficients:

(Intercept)

Identify a number that represents between birth weight and gestation days:

- 1. the magnitude of the association
- 2. the strength of the evidence

Estimate Std. Error t value Pr(>|t|)

0.540712 - 2.421

the strength of the association

<ロ > < 部 > < 重 > < 重 > の < で 42 / 44

<2e-16 \*\*\*

0.1560

STAT 217: Unit 3 Deck 3

The Data

Inference for SLR 000000000000

The Data

Correlation

babies\$gestation 0.031433

> cor(babies\$bwt\_lbs,babies\$gestation)

-1.309203

Estimate Std. Error t value Pr(>|t|)

0.540712 -2.421

0.001931 16.274

Inference for SLR 000000000000

44 / 44

> cor(babies\$bwt\_lbs,babies\$gestation) 0.42


Coefficients:

Estimate Std. Error t value Pr(>|t|) (Intercept) -1.309203 0.540712 -2.421 0.1560 babies\$gestation 0.031433 0.001931 16.274 <2e-16 \*\*\*

STAT 217: Unit 3 Deck 3

At  $\alpha = 0.05$ , what can we conclude? We (do/do not) have evidence that  $\beta_1$  differs from zero; so we (do/do not) have evidence of an association between gestation days and birth weight.

- 1. do: do
- 2. do not: do not
- 3. do: do not
- 4. do not; do



- 1. Yes, because the *p*-value for  $H_0$ :  $\beta_1 = 0$  is less than 0.05.
- 2. No. Although the *p*-value for  $H_0$ :  $\beta_1 = 0$  is less than 0.05, the estimated slope is small.
- 3. No, because the *p*-value for  $H_0$ :  $\beta_1 = 0$  is greater than 0.05.
- 4. Yes, the correlation is positive.
- 5. No, the correlation isn't that big.

