

## Cox regression models with multiple predictors

Shannon Pileggi

STAT 417

## OUTLINE

## Multiple predictors

Inference ( $\beta_i$ )

Inference  $(\beta_i + \beta_j)$

## Overall tests

## Dummy variables

## Cox regression with multiple predictors

When there are several predictors  $X_1, X_2, \dots, X_p$  that we believe are associated with hazard, then the CR model becomes:

Parameters are estimated by maximizing the partial (log) likelihood function and the fitted CR model is given by:

## Time invariant predictors

We will assume that the values of each predictor are measured on each individual at the beginning of the study and remain fixed over time - these are called *time invariant predictors*.

For which of the following predictors is it reasonable to assume that they are *time invariant*? In a study of time to college graduation...

1. high school GPA
2. college GPA
3. gender
4. illegal drug use (yes/no)
5. weight

Multiple predictors	Inference ( $\beta_i$ )	Inference ( $\beta_i + \beta_j$ )	Overall tests	Dummy variables
○○●○○○○○○○	○○○○○	○○○○○	○○○○○○○	○○○○○○○○○○○○○

## Proportionality assumption

The proportionality assumption of the Cox regression model implies that for **two sets of values** of predictors,  $\{x_1, \dots, x_p\}$  and  $\{x_1^*, \dots, x_p^*\}$ , ...

1. the hazard ratio remains constant over time
2. the difference between the hazards remains constant over time
3. the ratio of log hazards remains constant over time
4. the difference between the log hazards remains constant over time

STAT 417: Set 10 5 / 47

Multiple predictors	Inference ( $\beta_i$ )	Inference ( $\beta_i + \beta_j$ )	Overall tests	Dummy variables
○○○●○○○○○○○	○○○○○	○○○○○	○○○○○○○	○○○○○○○○○○○○○

## Interpretation of $\beta_j$ 's and $e^{c\beta_j}$ 's

$\beta_j$

$e^{c\beta_j}$

STAT 417: Set 10 6 / 47

Multiple predictors	Inference ( $\beta_i$ )	Inference ( $\beta_i + \beta_j$ )	Overall tests	Dummy variables
○○○○●○○○○○	○○○○○	○○○○○	○○○○○○○	○○○○○○○○○○○○○

## VALCG study with multiple predictors

Recall the lung cancer study, and consider the predictors:

- ▶  $X_1$  = Karnofsky score (quantitative)
- ▶  $X_2$  = Cancer treatment (0 = standard, 1 = test) (categorical)

1. Write the form of the CR model with the two explanatory variables.
2. Provide an interpretation for  $\beta_1$ .

STAT 417: Set 10 7 / 47

Multiple predictors	Inference ( $\beta_i$ )	Inference ( $\beta_i + \beta_j$ )	Overall tests	Dummy variables
○○○○○●○○○○○	○○○○○	○○○○○	○○○○○○○	○○○○○○○○○○○○○

## VALCG study with multiple predictors, cont.

3. Provide an interpretation for  $e^{\beta_2}$ .

STAT 417: Set 10 8 / 47

## VALCG study with multiple predictors: hazard ratios

Set up the hazard ratios (in terms of  $\beta$ 's) for:

1. Patients taking the test treatment to patients taking the standard treatment (fixing Karnofsky score).
2. A ten point increase in Karnofsky score, fixing the treatment.

## VALCG study with multiple predictors: hazard ratios

Set up the hazard ratios (in terms of  $\beta$ 's) for:

3. Patients taking the test treatment whose Karnofsky score is 10 points higher than patients taking the standard treatment.

## VALCG study with multiple predictors: R output

R Code

```
CR_mod1 <- coxph(Surv(time, status) ~ karno + trt, data = veteran)
summary(CR_mod1)
```

R Code

R Output

```
      coef exp(coef) se(coef)      z Pr(>|z|)
karno -0.033954  0.966616  0.005084 -6.679  2.4e-11 ***
trt    0.177322  1.194016  0.183149  0.968   0.333
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
      exp(coef) exp(-coef) lower .95 upper .95
karno   0.9666    1.0345    0.9570  0.9763
trt     1.1940    0.8375    0.8339  1.7096
```

R Output

## VALCG study w/ mult. predictors: estimated hazard ratios

Compute and interpret estimated hazard ratios for:

1. Patients taking the test treatment to patients taking the standard treatment (fixing Karnofsky score).
2. A ten point increase in Karnofsky score, fixing the treatment.

## VALCG study w/ mult. predictors: estimated hazard ratios

Compute and interpret estimated hazard ratios for:

- Patients taking the test treatment whose Karnofsky score is 10 points higher than patients taking the standard treatment.

Multiple predictors

Inference ( $\beta_i$ )

Inference ( $\beta_i + \beta_j$ )

Overall tests

Dummy variables

## Full R output

R Output

```

      coef exp(coef) se(coef)      z Pr(>|z|)
karno -0.033954  0.966616  0.005084 -6.679  2.4e-11 ***
trt    0.177322  1.194016  0.183149  0.968   0.333
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

      exp(coef) exp(-coef) lower .95 upper .95
karno   0.9666    1.0345    0.9570  0.9763
trt     1.1940    0.8375    0.8339  1.7096

Concordance= 0.712 (se = 0.03 )
Rsquare= 0.269 (max possible= 0.999 )
Likelihood ratio test= 42.97 on 2 df,  p=4.676e-10
Wald test               = 44.66 on 2 df,  p=2.001e-10
Score (logrank) test = 46.78 on 2 df,  p=6.933e-11

```

R Output

## Hypothesis test for $\beta_j$

Test whether the predictor  $X_j$  has a significant effect on hazard  
**when all other predictors are included in the model with:**

The Wald test statistic is:

Multiple predictors  
○○○○○○○○○○
Inference ( $\beta_i$ )  
○○●○○
Inference ( $\beta_i + \beta_j$ )  
○○○○○
Overall tests  
○○○○○○○
Dummy variables  
○○○○○○○○○○○○○○

## VALCG study: Wald tests

Are treatment and Karnofsky score significant predictors of hazard?

R Output

	coef	exp(coef)	se(coef)	z	Pr(> z )
karno	-0.033954	0.966616	0.005084	-6.679	2.4e-11 ***
trt	0.177322	1.194016	0.183149	0.968	0.333

R Output

STAT 417: Set 10
17 / 47

Multiple predictors  
○○○○○○○○○○
Inference ( $\beta_i$ )  
○○○○●
Inference ( $\beta_i + \beta_j$ )  
○○○○○
Overall tests  
○○○○○○○
Dummy variables  
○○○○○○○○○○○○○○

## VALCG study: CI for HR

Construct and interpret a 95% confidence interval for the population hazard ratio of patients on the test treatment to those on the standard treatment (for fixed Karnofsky score).

R Output

	coef	exp(coef)	se(coef)	z	Pr(> z )
karno	-0.033954	0.966616	0.005084	-6.679	2.4e-11 ***
trt	0.177322	1.194016	0.183149	0.968	0.333

R Output

STAT 417: Set 10
18 / 47

Multiple predictors  
○○○○○○○○○○
Inference ( $\beta_i$ )  
○○○○●
Inference ( $\beta_i + \beta_j$ )  
○○○○○
Overall tests  
○○○○○○○
Dummy variables  
○○○○○○○○○○○○○○

## VALCG study: CI for HR

Interpret the interval associated with Karnofsky score.

R Output

	exp(coef)	exp(-coef)	lower .95	upper .95
karno	0.9666	1.0345	0.9570	0.9763
trt	1.1940	0.8375	0.8339	1.7096

R Output

STAT 417: Set 10
19 / 47

Multiple predictors  
○○○○○○○○○○
Inference ( $\beta_i$ )  
○○○○○
Inference ( $\beta_i + \beta_j$ )  
●○○○○
Overall tests  
○○○○○○○
Dummy variables  
○○○○○○○○○○○○○○

Multiple predictors  
Inference ( $\beta_i$ )  
Inference ( $\beta_i + \beta_j$ )  
Overall tests  
Dummy variables

STAT 417: Set 10
20 / 47

## Inference for a linear combination of $\beta$ 's

The general form of the CI:

$$\exp \left[ c\hat{\beta}_j \pm z_{\alpha/2} |c| SE(\hat{\beta}_j) \right]$$

The form of the true hazard ratio for patients taking the test treatment whose Karnofsky score is 10 points higher than patients taking the standard treatment:

$$HR = \exp[10\beta_1 + \beta_2]$$

How could we extend the above expression to this linear combination?

1. Which parts are straightforward?
2. Which parts need care?

## CI for a linear combination of $\beta$ 's

The  $100(1 - \alpha)\%$  CI for the  $HR$  of the general form  $e^{(a\beta_i + b\beta_j)}$  is given by:

## Estimated variance-covariance matrix

R Code

```
CR_mod1$var
```

R Code

R Output

```
      [,1]      [,2]
[1,] 2.584253e-05 -0.0001026675
[2,] -1.026675e-04 0.0335433798
```

R Output

## Estimated variance-covariance matrix

- ▶ The estimated covariance between  $\hat{\beta}_1$  and  $\hat{\beta}_2$  is:
- ▶ The estimated variance of  $\hat{\beta}_1$  is:
- ▶ The estimated variance of  $\hat{\beta}_2$  is:
- ▶ The standard errors of  $\hat{\beta}_1$  and  $\hat{\beta}_2$  are:

Multiple predictors ○○○○○○○○○○	Inference ( $\beta_i$ ) ○○○○○	Inference ( $\beta_i + \beta_j$ ) ○○○○●	Overall tests ○○○○○○○	Dummy variables ○○○○○○○○○○○○○○
-----------------------------------	----------------------------------	--	--------------------------	-----------------------------------

## VALCG study: CI for HR for linear combination of $\beta$ 's

Compute the confidence interval for the population hazard ratio for patients taking the test treatment whose Karnofsky score is 10 points higher than patients taking the standard treatment.

HR =  $\exp[10\beta_1 + \beta_2]$

STAT 417: Set 10 25 / 47

Multiple predictors ○○○○○○○○○○	Inference ( $\beta_i$ ) ○○○○○	Inference ( $\beta_i + \beta_j$ ) ○○○○○	Overall tests ●○○○○○○	Dummy variables ○○○○○○○○○○○○○○
-----------------------------------	----------------------------------	--	--------------------------	-----------------------------------

Multiple predictors

Inference ( $\beta_i$ )

Inference ( $\beta_i + \beta_j$ )

Overall tests

Dummy variables

STAT 417: Set 10 26 / 47

Multiple predictors ○○○○○○○○○○	Inference ( $\beta_i$ ) ○○○○○	Inference ( $\beta_i + \beta_j$ ) ○○○○○	Overall tests ●○○○○○○	Dummy variables ○○○○○○○○○○○○○○
-----------------------------------	----------------------------------	--	--------------------------	-----------------------------------

## Full R output

```

R Output

      coef exp(coef)  se(coef)      z Pr(>|z|)
karno -0.033954  0.966616  0.005084 -6.679  2.4e-11 ***
trt    0.177322  1.194016  0.183149  0.968   0.333
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

      exp(coef) exp(-coef) lower .95 upper .95
karno   0.9666    1.0345    0.9570    0.9763
trt     1.1940    0.8375    0.8339    1.7096

Concordance= 0.712 (se = 0.03 )
Rsquare= 0.269 (max possible= 0.999 )
Likelihood ratio test= 42.97 on 2 df,  p=4.676e-10
Wald test              = 44.66 on 2 df,  p=2.001e-10
Score (logrank) test = 46.78 on 2 df,  p=6.933e-11
  
```

STAT 417: Set 10 27 / 47

Multiple predictors ○○○○○○○○○○	Inference ( $\beta_i$ ) ○○○○○	Inference ( $\beta_i + \beta_j$ ) ○○○○○	Overall tests ●○○○○○○	Dummy variables ○○○○○○○○○○○○○○
-----------------------------------	----------------------------------	--	--------------------------	-----------------------------------

## Overall tests

What is different about the two highlighted lines?

STAT 417: Set 10 28 / 47

## Three tests

### 1. Partial Likelihood Ratio Test:

$$G_I = 2 \left[ l_p(\hat{\beta}) - l_p(0) \right]$$

### 2. Wald Test:

$$G_W = \hat{\beta}^T \mathbf{I}(\hat{\beta}) \hat{\beta}$$

### 3. Score Test:

$$G_S = \mathbf{u}^T(\mathbf{0})[\mathbf{I}(\mathbf{0})]^{-1}\mathbf{u}(\mathbf{0})$$

All three test statistics ( $G_I$ ,  $G_W$ ,  $G_S$ ) follow a  $\chi^2$ -distribution with  $p$  degrees of freedom.

### Details:

- ▶  $l_p(\beta)$  is the log partial likelihood function
- ▶  $\mathbf{I}(\beta)$  is the observed information matrix
- ▶  $\mathbf{0} = (0, 0, \dots, 0)^T$  is a  $p$ -vector of 0's
- ▶  $\mathbf{u}^T(\mathbf{0})$  is the vector of partial derivatives of the log partial likelihood function (also called a vector of scores) evaluated at  $\beta = \mathbf{0}$

## VALCG study: interpret the results

### R Output

```
Likelihood ratio test= 42.97 on 2 df, p=4.676e-10
Wald test              = 44.66 on 2 df, p=2.001e-10
Score (logrank) test = 46.78 on 2 df, p=6.933e-11
```

### R Output

## Partial likelihood ratio test

$l_p(\hat{\beta})$  = log partial likelihood function evaluated at the parameter estimates (measures of goodness-of-fit of the CR model to the data when the predictors  $X_1, X_2, \dots, X_p$  are included)

$l_p(0)$  = log partial likelihood function evaluated at 0 values (measures of the fit of the *null model*, i.e. a CR model with no predictors and consisting of only the baseline hazard function)

- ▶ If  $l_p(\hat{\beta})$  is “much larger” than  $l_p(0)$ , then:
- ▶ The partial likelihood ratio test statistic compares  $l_p(\hat{\beta})$  to  $l_p(0)$ :

## Log partial likelihoods in R

### R Code

```
CR_mod1$loglik
```

### R Code

### R Output

```
[1] -505.4491 -483.9657
```

### R Output

$l_p(0)$  = left value (-505.4491)  
 $l_p(\hat{\beta})$  = right value (-483.9657)

Verify that the partial likelihood ratio statistic is  $G_I = 42.97$ .



Multiple predictors ○○○○○○○○○○	Inference ( $\beta_i$ ) ○○○○○	Inference ( $\beta_i + \beta_j$ ) ○○○○○	Overall tests ○○○○○○●	Dummy variables ○○○○○○○○○○○○○○
-----------------------------------	----------------------------------	--	--------------------------	-----------------------------------

## $p$ -values from $\chi^2$ distribution in R

R Code

```
pchisq(q = 42.97, df = 2, lower.tail = F)
```

R Code

R Output

```
4.668561e-10
```

R Output

STAT 417: Set 10

33 / 47

Multiple predictors ○○○○○○○○○○	Inference ( $\beta_i$ ) ○○○○○	Inference ( $\beta_i + \beta_j$ ) ○○○○○	Overall tests ○○○○○○○	Dummy variables ●○○○○○○○○○○○○
-----------------------------------	----------------------------------	--	--------------------------	----------------------------------

## Multiple predictors

## Inference ( $\beta_i$ )

## Inference ( $\beta_i + \beta_j$ )

## Overall tests

## Dummy variables

STAT 417: Set 10

34 / 47

Multiple predictors ○○○○○○○○○○	Inference ( $\beta_i$ ) ○○○○○	Inference ( $\beta_i + \beta_j$ ) ○○○○○	Overall tests ○○○○○○○	Dummy variables ●○○○○○○○○○○○○
-----------------------------------	----------------------------------	--	--------------------------	----------------------------------

## Categorical predictors with $> 2$ levels

- To include a categorical predictor with  $k$  levels (i.e.  $k$  different possible values) into the CR model, a set of  $k$  **dummy variables**,  $D_1, D_2, \dots, D_k$ , must be created to “represent” the different values that  $X$  can take.

STAT 417: Set 10

35 / 47

Multiple predictors ○○○○○○○○○○	Inference ( $\beta_i$ ) ○○○○○	Inference ( $\beta_i + \beta_j$ ) ○○○○○	Overall tests ○○○○○○○	Dummy variables ○○●○○○○○○○○○○
-----------------------------------	----------------------------------	--	--------------------------	----------------------------------

## Categorical predictors with $> 2$ levels

Write out the dummy variables required for a CR model for the VALCG lung cancer study with the categorical predictor  $X = \text{cancer cell type}$  (**small cell**, **squamous**, **large cell**, and **adenocarcinoma**).

STAT 417: Set 10

36 / 47



Multiple predictors  
○○○○○○○○○○○

Inference ( $\beta_i$ )  
○○○○○

Inference ( $\beta_i + \beta_j$ )  
○○○○○

Overall tests  
○○○○○○○

Dummy variables  
○○○○○○○●○○○○○

## VALCG study - CR model with celltype

What is the interpretation of  $\hat{\beta}_1 = -0.9176$ ? The 1 of death for patients with 2 lung cancer is estimated to be 0.92 3 than the 4 of death for patients with 5.

- hazard, log hazard, hazard ratio
- adeno, large, smallcell, squamous
- lower, higher, times
- hazard, log hazard, hazard ratio
- adeno, large, smallcell, squamous

STAT 417: Set 10

41 / 47

Multiple predictors  
○○○○○○○○○○○

Inference ( $\beta_i$ )  
○○○○○

Inference ( $\beta_i + \beta_j$ )  
○○○○○

Overall tests  
○○○○○○○

Dummy variables  
○○○○○○○●○○○○○

## VALCG study - CR model with celltype

R Output

	coef	exp(coef)	se(coef)	z	Pr(> z )	
celltypelarge	-0.9176	0.3995	0.2880	-3.186	0.00144	**
celltypesmallcell	-0.1465	0.8638	0.2493	-0.587	0.55687	
celltypesquamous	-1.1477	0.3174	0.2929	-3.919	8.9e-05	***

R Output

Identify two sets of two groups of cancers that appear to have a similar effect on hazard. Classify these sets as “better off” or “worse off”.

Set 1:    adeno    large    smallcell    squamous

Set 2:    adeno    large    smallcell    squamous

STAT 417: Set 10

42 / 47

Multiple predictors  
○○○○○○○○○○○

Inference ( $\beta_i$ )  
○○○○○

Inference ( $\beta_i + \beta_j$ )  
○○○○○

Overall tests  
○○○○○○○

Dummy variables  
○○○○○○○●○○○○○

## VALCG study - CR model with celltype

R Output

	exp(coef)	exp(-coef)	lower .95	upper .95
celltypelarge	0.3995	2.503	0.2272	0.7025
celltypesmallcell	0.8638	1.158	0.5299	1.4080
celltypesquamous	0.3174	3.151	0.1788	0.5634

R Output

What is the interpretation of 0.3995?

STAT 417: Set 10

43 / 47

Multiple predictors  
○○○○○○○○○○○

Inference ( $\beta_i$ )  
○○○○○

Inference ( $\beta_i + \beta_j$ )  
○○○○○

Overall tests  
○○○○○○○

Dummy variables  
○○○○○○○●○○○○○

## VALCG study - CR model with celltype

R Output

	exp(coef)	exp(-coef)	lower .95	upper .95
celltypelarge	0.3995	2.503	0.2272	0.7025
celltypesmallcell	0.8638	1.158	0.5299	1.4080
celltypesquamous	0.3174	3.151	0.1788	0.5634

R Output

What is the interpretation of the interval 0.5299 - 1.4080?

STAT 417: Set 10

44 / 47

Multiple predictors  
○○○○○○○○○○○

Inference ( $\beta_i$ )  
○○○○○

Inference ( $\beta_i + \beta_j$ )  
○○○○○

Overall tests  
○○○○○○○

Dummy variables  
○○○○○○○○○○●○○

## VALCG study - CR model with celltype

R Output

	coef	exp(coef)	se(coef)	z	Pr(> z )	
celltypelarge	-0.9176	0.3995	0.2880	-3.186	0.00144	**
celltypesmallcell	-0.1465	0.8638	0.2493	-0.587	0.55687	
celltypesquamous	-1.1477	0.3174	0.2929	-3.919	8.9e-05	***

R Output

Estimate how many times higher (or percentage points lower) the hazard rate is for patients with small cell cancer than patients with large cell cancer.

STAT 417: Set 10

45 / 47

Multiple predictors  
○○○○○○○○○○○

Inference ( $\beta_i$ )  
○○○○○

Inference ( $\beta_i + \beta_j$ )  
○○○○○

Overall tests  
○○○○○○○

Dummy variables  
○○○○○○○○○○●○○

## VALCG study - CR model with celltype

R Output

	coef	exp(coef)	se(coef)	z	Pr(> z )	
celltypelarge	-0.9176	0.3995	0.2880	-3.186	0.00144	**
celltypesmallcell	-0.1465	0.8638	0.2493	-0.587	0.55687	
celltypesquamous	-1.1477	0.3174	0.2929	-3.919	8.9e-05	***

R Output

What is the general approach to construct a CI for the population hazard ratio for patients with small cell cancer relative to patients large cell cancer?

STAT 417: Set 10

46 / 47

Multiple predictors  
○○○○○○○○○○○

Inference ( $\beta_i$ )  
○○○○○

Inference ( $\beta_i + \beta_j$ )  
○○○○○

Overall tests  
○○○○○○○

Dummy variables  
○○○○○○○○○○●○○

## VALCG study - CR model with celltype

R Output

	coef	exp(coef)	se(coef)	z	Pr(> z )	
celltypelarge	-0.9176	0.3995	0.2880	-3.186	0.00144	**
celltypesmallcell	-0.1465	0.8638	0.2493	-0.587	0.55687	
celltypesquamous	-1.1477	0.3174	0.2929	-3.919	8.9e-05	***
...						
Likelihood ratio test=	24.85	on 3 df,	p=1.661e-05			
Wald test	= 24.09	on 3 df,	p=2.387e-05			
Score (logrank) test =	25.51	on 3 df,	p=1.208e-05			

R Output

Is type of cancer associated with hazard of death?

STAT 417: Set 10

47 / 47