

Summary characteristics ○○○○○○○○○	Evaluating fit ○○○○○○○	Parameter estimation ○○○○○○○○○○○○○	Minitab ○○○○○	Summary ○○○
--------------------------------------	---------------------------	---------------------------------------	------------------	----------------

# Parametric models continued: summary characteristics, fit, and estimation

Shannon Pileggi

STAT 417

STAT 417: Set 3
1 / 39

Summary characteristics ○○○○○○○○○	Evaluating fit ○○○○○○○	Parameter estimation ○○○○○○○○○○○○○	Minitab ○○○○○	Summary ○○○
--------------------------------------	---------------------------	---------------------------------------	------------------	----------------

# OUTLINE

- Summary characteristics
- Evaluating fit
- Parameter estimation
- Minitab
- Summary

STAT 417: Set 3
2 / 39

Summary characteristics ●○○○○○○○	Evaluating fit ○○○○○○○	Parameter estimation ○○○○○○○○○○○○○	Minitab ○○○○○	Summary ○○○
-------------------------------------	---------------------------	---------------------------------------	------------------	----------------

# Summary characteristics of $T$

With a specified probability distribution for  $T$ , we can answer:

- ▶ What is the average time it takes for a motorist to react aggressively?
- ▶ What is the median age that an individual has his/her first drink?
- ▶ For patients who have survived 100 days after being diagnosed with lung cancer, what is the average time they have left to live?

STAT 417: Set 3
3 / 39

Summary characteristics ●○○○○○○○	Evaluating fit ○○○○○○○	Parameter estimation ○○○○○○○○○○○○○	Minitab ○○○○○	Summary ○○○
-------------------------------------	---------------------------	---------------------------------------	------------------	----------------

# Mean survival time

The **mean** (or **expected value**) of the time-to-event random variable  $T$  ( $T \geq 0$ ), denoted by  $E(T)$  or  $\mu$ , is given by:


STAT 417: Set 3
4 / 39

## Motorist reaction time: compute $E(T)$

Suppose that the time until a motorist reacts aggressively is exponentially distributed with parameter  $\lambda = 5.77$ . Find the expected time it takes for a motorist to react aggressively.

## Group Exercise

Another measure of the “center” of the distribution for  $T$  is the **median survival time**.



The median survival time is \_\_\_\_\_ the mean survival time of 13.5.

1. greater than
2. less than
3. equal to
4. not enough information to determine

## Percentiles of $T$

- ▶ The **median survival time** is a special case of a *percentile* of the distribution of  $T$  (the 50<sup>th</sup> percentile). This is the time at which 50% of the individuals have experienced the event.
- ▶ The  $p^{\text{th}}$  percentile of the distribution of the time-to-event random variable  $T$  is the value of  $T$ , denoted  $t_p$  such that:
- ▶ **Interpretation:** The  $p^{\text{th}}$  percentile,  $t_p$ , can be interpreted as:

## Graphical representation of percentiles

Summary characteristics ○○○○○●○○	Evaluating fit ○○○○○○○	Parameter estimation ○○○○○○○○○○○○○	Minitab ○○○○○	Summary ○○○
-------------------------------------	---------------------------	---------------------------------------	------------------	----------------

## Age at first drink: median survival time

Suppose that the age at first drink of alcohol follows an exponential distribution with parameter  $\lambda = 13.5$ . Find the median age at first drink. Recall that  $S(t) = \exp(-t/\lambda)$ .

STAT 417: Set 3 9 / 39

Summary characteristics ○○○○○●○○	Evaluating fit ○○○○○○○	Parameter estimation ○○○○○○○○○○○○○	Minitab ○○○○○	Summary ○○○
-------------------------------------	---------------------------	---------------------------------------	------------------	----------------

## Mean residual lifetime

- ▶ **Example: Lung Cancer.** Suppose an individual with lung cancer has survived 100 months since being diagnosed. Then how much time (on average) does this individual have left to live? We can set this problem up as a *conditional expectation*:
- ▶ This is an example of quantity known as the **mean residual lifetime (mrl)**, denoted  $mrl(t^*)$ , and is defined to be the average remaining lifetime of all individuals given that the individuals have survived to time  $t^*$ , i.e:

STAT 417: Set 3 10 / 39

Summary characteristics ○○○○○○○●	Evaluating fit ○○○○○○○	Parameter estimation ○○○○○○○○○○○○○	Minitab ○○○○○	Summary ○○○
-------------------------------------	---------------------------	---------------------------------------	------------------	----------------

## Lung cancer example: mean residual lifetime

If the time until death from cancer follows an exponential distribution with parameter  $\lambda = 130.18$ , then find the average remaining lifetime for those patients who have survived at least 100 days.

STAT 417: Set 3 11 / 39

Summary characteristics ○○○○○○○○○	Evaluating fit ●○○○○○○○	Parameter estimation ○○○○○○○○○○○○○	Minitab ○○○○○	Summary ○○○
--------------------------------------	----------------------------	---------------------------------------	------------------	----------------

Summary characteristics

Evaluating fit

Parameter estimation

Minitab

Summary

STAT 417: Set 3 12 / 39

Summary characteristics ○○○○○○○○○	Evaluating fit ●○○○○○	Parameter estimation ○○○○○○○○○○○	Minitab ○○○○○	Summary ○○○
--------------------------------------	--------------------------	-------------------------------------	------------------	----------------

## Considerations for parametric models


- 1.
- 2.
- 3.

STAT 417: Set 3 13 / 39

Summary characteristics ○○○○○○○○○	Evaluating fit ○○●○○○	Parameter estimation ○○○○○○○○○○○	Minitab ○○○○○	Summary ○○○
--------------------------------------	--------------------------	-------------------------------------	------------------	----------------

## Veteran's Administration Lung Cancer Study (VALCS)

Lung cancer example



The figure is a histogram showing the distribution of time until death from lung cancer. The x-axis is labeled 'Time until death from lung cancer (days)' and ranges from 0 to 1000. The y-axis represents frequency. Two probability density curves are superimposed: a red line for an Exponential distribution with  $\lambda = 130.2$ , and a teal line for a Normal distribution with mean = 130.7 and standard deviation = 162.2. The exponential curve starts high at day 0 and decays rapidly, while the normal curve is more spread out and centered around 130 days.

Lifetimes (since treatment) of 137 lung cancer patients taken from another lung cancer study with two different probability density curves superimposed (exponential and normal).

STAT 417: Set 3 14 / 39

Summary characteristics ○○○○○○○○○	Evaluating fit ○○●○○○	Parameter estimation ○○○○○○○○○○○	Minitab ○○○○○	Summary ○○○
--------------------------------------	--------------------------	-------------------------------------	------------------	----------------

## Strategies for fitting parametric models

- 1.
- 2.
- 3.

STAT 417: Set 3 15 / 39

Summary characteristics ○○○○○○○○○	Evaluating fit ○○●○○○	Parameter estimation ○○○○○○○○○○○	Minitab ○○○○○	Summary ○○○
--------------------------------------	--------------------------	-------------------------------------	------------------	----------------

## Probability plots for any distribution

- ▶ Can construct a probability plot for any distribution
- ▶ With incomplete data, special techniques are used in the probability plot to account for censoring
- ▶ Assess if an exponential distribution fits the data well


$H_0$ :

$H_a$ :

- 1.
- 2.

STAT 417: Set 3 16 / 39


## VALCS study: probability plots



Which distribution appears to fit the data better, and why?

1. normal
2. exponential

## Motorist reaction times: probability plots



Summary characteristics

Evaluating fit

Parameter estimation

Minitab

Summary

## Parameter estimation

- ▶ We have worked through many examples of parametric models for  $T$ , and in each example, the value(s) of the parameter(s) were given. For example, the time until death for the lung cancer data was assumed to follow an exponential distribution with parameter  $\lambda = 130.18$ .
- ▶ Also, in the previous probability plots (observe the box on the right side), values of the parameters were provided. How are these values computed?

## Maximum likelihood estimation

Suppose we have a random sample of observations  $T_1, T_2, \dots, T_n$  on  $n$  individuals, each with the same pdf  $f(t_i)$ ,  $i = 1, \dots, n$  and parameter  $\theta$  (or possibly more than one parameter).

- ▶ Then **likelihood function** is given by:
- ▶ The likelihood function tells how likely the observed sample is as a function of the parameter values.
- ▶ By *maximizing* the likelihood function, we get the parameter value(s) that most likely would generate our given observed sample.

## Maximum likelihood estimation, cont.

**Goal:** Find an estimate of  $\theta$  that maximizes the likelihood function,  $L(\theta)$ .

**Process:** 1.

2.

3.

## MLE example: exponential distribution

Suppose we obtain a random sample of 5 observations,  $T_1, T_2, \dots, T_5$ , from an exponential distribution with parameter  $\lambda$  and pdf:


$$f(t) = \frac{1}{\lambda} e^{-t/\lambda}$$

1. Construct the likelihood function  $L(\lambda)$ .

## MLE example: exponential distribution, cont.

2. Suppose the observed values of  $T_1, T_2, \dots, T_5$  are 5.3, 4.8, 0.4, 2.3, and 0.4. Plug these into  $L(\lambda)$  and examine the graph of  $L(\lambda)$  as a function of  $\lambda$ . At which value of  $\lambda$  does the likelihood function appear to be maximized?

Exponential likelihood function



Summary characteristics

Evaluating fit

Parameter estimation

Minitab

Summary

MLE example: exponential distribution, cont.

3. Find the value of  $\lambda$  that maximizes  $L(\lambda)$ .

STAT 417: Set 3

25 / 39

Summary characteristics

Evaluating fit

Parameter estimation

Minitab

Summary

STAT 417: Set 3

26 / 39

Summary characteristics

Evaluating fit

Parameter estimation

Minitab

Summary

Likelihood function for right censored data

If there are right censored event times, then can't simply compute the likelihood function  $L$  as the product of the pdf's. Contributions to the likelihood function differ:

- ▶ If the event time for observation  $i$  is complete, then that observation contributes:
- ▶ If the event time for observation  $i$  is (right) censored, then that observation contributes:
- ▶ Putting the contributions of all observation together, the likelihood function is given by:

STAT 417: Set 3

27 / 39

Summary characteristics

Evaluating fit

Parameter estimation

Minitab

Summary

MLE for right censored data

The following data consist of the times (in months) until relapse of 10 bone marrow transplant patients. Patients 7-10 were free of relapse at the end of the study. Suppose time to relapse follows an exponential distribution with parameter  $\lambda$ .

Patient	1	2	3	4	5	6	7	8	9	10
Relapse Time	5	8	12	24	32	17	16 <sup>+</sup>	17 <sup>+</sup>	19 <sup>+</sup>	30 <sup>+</sup>

- Construct the likelihood function for the parameter  $\lambda$  and find the MLE for  $\lambda$ .
- Assuming that time until relapse follows an exponential probability distribution with  $\lambda$  found in Part (1), what is the mean time until relapse?

STAT 417: Set 3

28 / 39

Summary characteristics	Evaluating fit	Parameter estimation	Minitab	Summary
○○○○○○○○	○○○○○○	○○○○○○○○●	○○○○○	○○○

## MLE for right censored data, cont.

STAT 417: Set 3

29 / 39

Summary characteristics	Evaluating fit	Parameter estimation	Minitab	Summary
○○○○○○○○	○○○○○○	○○○○○○○○●	○○○○○	○○○

## MLE for right censored data, cont.

- Assuming that time until relapse follows an exponential probability distribution with  $\lambda$  found in Part (1), what is the mean time until relapse?

STAT 417: Set 3

30 / 39

Summary characteristics	Evaluating fit	Parameter estimation	Minitab	Summary
○○○○○○○○	○○○○○○	○○○○○○○○○○	●○○○○○	○○○

Summary characteristics

Evaluating fit

Parameter estimation

**Minitab**

Summary

STAT 417: Set 3

31 / 39

Summary characteristics	Evaluating fit	Parameter estimation	Minitab	Summary
○○○○○○○○	○○○○○○	○○○○○○○○○○	●○○○○○	○○○

## Parametric distribution analysis in Minitab

- Along with the probability plots, the MLE's, and summary characteristics about the distribution of  $T$  are also computed in Minitab when Parametric Distribution Analysis is performed.
- The MLE's are provided in the box to the right of the plot, as well as in the output in the Minitab Session Window.
- The mean, median, quartiles, and several other percentiles are displayed in the Session Window.

STAT 417: Set 3

32 / 39



Summary characteristics  
○○○○○○○○○

Evaluating fit  
○○○○○○○

Parameter estimation  
○○○○○○○○○○○

Minitab  
○○●○○○

Summary  
○○○

## Motorist reaction times in Minitab (lognormal model)

Variable: Seconds until Horn Honk

Censoring Information    Count

Uncensored value	42
Right censored value	15

Censoring value: Censor Horn = 0

STAT 417: Set 3

33 / 39

Summary characteristics  
○○○○○○○○○

Evaluating fit  
○○○○○○○

Parameter estimation  
○○○○○○○○○○○

Minitab  
○○○●○○

Summary  
○○○

## Parametric distribution analysis in Minitab, cont.

Estimation Method: Maximum Likelihood

Distribution:    Lognormal

Parameter Estimates

		Standard	95.0% Normal CI	
Parameter	Estimate	Error	Lower	Upper
Location	1.43054	0.0851775	1.26360	1.59749
Scale	0.599958	0.0665560	0.482718	0.745673

Log-Likelihood = -100.729

Goodness-of-Fit

Anderson-Darling (adjusted) = 6.163

...

STAT 417: Set 3

34 / 39

Summary characteristics  
○○○○○○○○○

Evaluating fit  
○○○○○○○

Parameter estimation  
○○○○○○○○○○○

Minitab  
○○○○●○

Summary  
○○○

## Parametric distribution analysis in Minitab, cont.

Characteristics of Distribution

	Estimate	Standard Error	95.0% Normal CI	
			Lower	Upper
Mean(MTTF)	5.00540	0.500060	4.11529	6.08804
Standard Deviation	3.29467	0.672961	2.20774	4.91673
Median	4.18097	0.356124	3.53813	4.94060
First Quartile(Q1)	2.78954	0.249485	2.34102	3.32400
Third Quartile(Q3)	6.26644	0.643405	5.12418	7.66334
Interquartile Range(IQR)	3.47690	0.541312	2.56254	4.71752

STAT 417: Set 3

35 / 39

Summary characteristics  
○○○○○○○○○

Evaluating fit  
○○○○○○○

Parameter estimation  
○○○○○○○○○○○

Minitab  
○○○○○●

Summary  
○○○

## Parametric distribution analysis in Minitab, cont.

Table of Percentiles

Percent	Percentile	Standard Error	95.0% Normal CI	
			Lower	Upper
1	1.03545	0.169643	0.751052	1.42753
2	1.21943	0.181146	0.911401	1.63155
3	1.35276	0.188314	1.02974	1.77711

...

What do you think the first line means?

- $Pr(T < 1.03545) = 0.01$
- $Pr(T < 1) = 0.0103545$
- $Pr(T > 1.03545) = 0.01$
- $Pr(T > 1) = 0.0103545$

STAT 417: Set 3

36 / 39

Summary characteristics ○○○○○○○○○	Evaluating fit ○○○○○○○	Parameter estimation ○○○○○○○○○○○○○	Minitab ○○○○○	Summary ●○○
Summary characteristics				
Evaluating fit				
Parameter estimation				
Minitab				
Summary				

STAT 417: Set 3

37 / 39

Summary characteristics ○○○○○○○○○	Evaluating fit ○○○○○○○	Parameter estimation ○○○○○○○○○○○○○	Minitab ○○○○○	Summary ○○●
Recap				
<ul style="list-style-type: none"> <li>▶ Features of time-to-event data and survival studies (e.g. censoring, truncation, etc.)</li> <li>▶ Parametric models for the time-to-event random variable <math>T</math>: <math>f(t)</math>, <math>F(t)</math>, <math>S(t)</math>, <math>h(t)</math>, <math>H(t)</math></li> <li>▶ Descriptive measures for <math>T</math>: <math>E(T)</math>, <math>t_p</math>, and, <math>\text{mrl}(t^*)</math></li> <li>▶ Strategies for selecting a probability model: probability plots, AD test statistic</li> <li>▶ Parameter estimation: maximum likelihood estimation</li> </ul>				

STAT 417: Set 3

38 / 39

Summary characteristics ○○○○○○○○○	Evaluating fit ○○○○○○○	Parameter estimation ○○○○○○○○○○○○○	Minitab ○○○○○	Summary ○○●
Future				
<p><b>Nonparametric</b> methods for summarizing and describing samples of time-to-event data:</p> <ul style="list-style-type: none"> <li>▶ Estimators for <math>S(t)</math>, <math>h(t)</math>, and <math>H(t)</math>.</li> <li>▶ Estimates of <math>E(T)</math> and <math>t_p</math></li> <li>▶ Inference methods for the estimator of <math>S(t)</math></li> <li>▶ Inference procedures for comparing two or more survival experiences.</li> <li>▶ Survival analysis in R</li> </ul>				

STAT 417: Set 3

39 / 39