PBL_BTech

by Pbl_btech Pbl_btech

Submission date: 02-Nov-2026 09:30AM (UTC+0530)
Submission ID: 3077430891
File name: PBL_BTech_Project_Report_Maharashtra_Sugarcane.docx (6.05M)
Word count: 8509
Character count: 54874

SYMBIOSIS INSTITUTE OF TECHNOLOGY, NAGPUR
Symbiosis International (Deemed University)
(Established under section 3 of the UGC Act, 1956)
Re-accredited by NAAC with 'A++' Grade | Awarded Category – I by UGC
Founder: Prof. Dr. S. B. Mujumdar, M. Sc., Ph. D. (Awarded Padma Bhushan and Padma Shri by President of India)

A PROJECT REPORT

ON

MAHARASHTRA SUGARCANE YIELD PREDICTION AND AGRO-RESOURCE OPTIMIZATION USING EXPLAINABLE MACHINE LEARNING

A project report submitted in partial fulfillment of the requirements for the degree of Bachelor of Technology in Computer Science and Engineering

BACHELOR OF TECHNOLOGY COMPUTER SCIENCE AND ENGINEERING

Submitted By

Shantanu Jumale (PRN: 24070521096)
Piyush Agrawal (PRN: 24070521073)
Om Shendre(PRN: 24070521195)

UNDER THE GUIDANCE OF

Dr. Snehalata wankhade
Assistant Professor / PBL Guide

SYMBIOSIS INSTITUTE OF TECHNOLOGY (SIT), NAGPUR /
PUNSYMBIOSIS INTERNATIONAL (DEEMED UNIVERSITY), PUNE

AY 2026-27

SYMBIOSIS INSTITUTE OF TECHNOLOGY, NAGPUR
Symbiosis International (Deemed University)
(Established under section 3 of the UGC Act, 1956)
Re-accredited by NAAC with 'A++' Grade | Awarded Category – I by UGC
Founder: Prof. Dr. S. B. Mujumdar, M. Sc., Ph. D. (Awarded Padma Bhushan and Padma Shri by President of India)

DEPARTMENT OF COMPUTER SCIENCE AND ENGINEERING
SYMBIOSIS INSTITUTE OF TECHNOLOGY, NAGPUR

CERTIFICATE

This is to certify that the Project work entitled **"Maharashtra Sugarcane Yield Prediction and Agro-Resource Optimization Using Explainable Machine Learning"** is carried out by **Shantanu jumale, Om Shendre, piyush Agarawal** in partial fulfillment for the award of the degree of **Bachelor of Technology in Computer Science and Engineering**, Symbiosis International (Deemed University), Pune during the academic year 2025-2026.

Dr. Snehalata Wankhade **PBL Guide**	**Dr. Giridhar Urkude** **PBL Coordinator**
Dr. Sagarkumar S. Badhiye **Head of Department, CSE**	**Dr. Nitin Rakesh** **Director, SIT Nagpur**

DECLARATION

I hereby declare that the project titled **"Maharashtra Sugarcane Yield Prediction and Agro-Resource Optimization Using Explainable Machine Learning"** submitted to Symbiosis Institute of Technology, Constituent of Symbiosis International (Deemed University) Pune for the award of the degree of Bachelor of Technology in Computer Science and Engineering is a result of original research carried out by me. I understand that my report may be made electronically available to the public. It is further declared that the project report or any part thereof has not been previously submitted to any University or Institute for the award of any degree or diploma.

Name of Student	:	Shantanu Jumale, Piyush Agrawal, Om Shendre
PRN	:	24070521096, 24070521073, 24070521195
Degree	:	Bachelor of Technology in CSE
Department	:	Computer Science and Engineering
Title of Project	:	Maharashtra Sugarcane Yield Prediction and Agro-Resource Optimization Using Explainable Machine Learning

Shantanu Jumale –

Om Shendre –

Piyush Agarawal -

_____

(Signature of the Student)

Date:

ACKNOWLEDGEMENT

I wish to express my profound gratitude to all individuals and institutions whose intellectual guidance, moral support, and infrastructural resources contributed to the successful completion of this Project-Based Learning (PBL) research endeavor.

First and foremost, I express my deepest sense of gratitude and indebtedness to my project guide and mentor, Dr. Giridhar Urkude, Assistant Professor, Department of Computer Science and Engineering, Symbiosis Institute of Technology. His profound technical acumen, constructive critiques, and meticulous supervision guided the architectural formulation and experimental evaluation of this research. His constant encouragement inspired me to pursue scientific rigor across machine learning modeling, explainability, and agro-resource optimization.

I am profoundly grateful to Prof. (Dr.) Nitin Rakesh, Director, Symbiosis Institute of Technology, Nagpur, for providing access to high-performance computing facilities, institutional resources, and a vibrant research atmosphere that made this multi-disciplinary project feasible.

I extend my sincere thanks to Dr. Sagarkumar S. Badhiye, Head of the Department of Computer Science and Engineering, for his administrative leadership, encouraging discussions, and insightful recommendations that enriched the scope of this investigation.

Special acknowledgments are due to the agronomic scientists and research staff at Vasantdada Sugar Institute (VSI), Manjari, Pune, and the Central Sugarcane Research Station (CSRS), Padegaon, Maharashtra, whose published technical reports, vertisol soil surveys, and cane physiological benchmarks provided vital ground-truth domain knowledge for our feature engineering framework.

I also thank my fellow classmates, lab colleagues, and faculty members in the Department of Computer Science and Engineering for their continuous peer feedback and stimulating technical exchanges.

Finally, I owe my deepest gratitude to my parents and family members for their unconditional love, moral support, patience, and encouragement throughout my academic journey.

ABSTRACT

Sugarcane (Saccharum officinarum) is an indispensable commercial cash crop underpinning the agrarian economy, rural cooperative industrial base, and bio-energy infrastructure of Maharashtra, India. The state represents India's second-largest producer of refined white sugar, cultivating over 1.4 million hectares across 14 principal districts and sustaining over 200 operational sugar mills alongside ethanol distilleries vital to the national E20 fuel blending mandate. However, agricultural productivity across Maharashtra displays severe spatial and temporal disparities. While the canal-command vertisol soils of Western Maharashtra (Kolhapur, Sangli, Satara) regularly achieve yields exceeding 400 Quintals/Acre (98.8 Tonnes/Hectare), semi-arid and rain-shadow districts in the Marathwada division (Beed, Osmanabad, Jalna, Latur) suffer drastic harvest depressions below 280 Q/Acre due to recurring meteorological droughts, acute groundwater depletion, and intense summer thermal stress (>42°C). Furthermore, uncalibrated farming practices characterized by excessive urea broadcasting without commensurate potassium application distort soil nutrient stoichiometry and trigger severe agrochemical runoff.

To address these critical vulnerabilities, this research formulates, develops, and validates a unified, end-to-end explainable machine learning architecture coupled with a constrained agro-resource optimization engine tailored specifically to Maharashtra's agro-climatic zones. A comprehensive empirical dataset comprising 1,000 multi-seasonal farm records (expanded across cropping regimes to 3,000 observational instances) was curated, cleansed of target leakage variables, and processed through an iterative 3-Stage Evolutionary Lifecycle Feature Engineering framework. This engineering protocol synthesized 12 specialized bio-climatic and physiological indicators, including the Crop Water Stress Index (CWSI), Cane Growth Index (CGI), Soil Water Retention Index (SWRI), and stoichiometric macronutrient ratios (N:P, N:K, P:K). Four distinct machine learning algorithms—Ordinary Least Squares (OLS) Linear Regression, Classification and Regression Trees (CART), Random Forest Regressor, and Extreme Gradient Boosting (XGBoost)—were trained and rigorously benchmarked under a stratified 5-fold cross-validation scheme. Hyperparameter optimization via RandomizedSearchCV elevated the tuned XGBoost regressor to state-of-the-art predictive performance, achieving a Coefficient of Determination (R^2) of 0.8716 ± 0.018, Mean Absolute Error (MAE) of 29.33 Q/Acre, and Root Mean Squared Error (RMSE) of 39.42 Q/Acre, substantially outperforming traditional Linear Regression ($R^2 = 0.5420$) and Random Forest ($R^2 = 0.8142$).

To dismantle algorithmic black-box opacity and establish actionable farmer trust, the model was interpreted using TreeSHAP (SHapley Additive exPlanations) grounded in cooperative game theory. Global attribution analysis revealed Cane Growth Index (mean |SHAP| = 14.82 Q/Acre), Soil Nitrogen (11.45 Q/Acre), and Soil Moisture (9.87 Q/Acre) as the primary positive drivers of cane tonnage, while CWSI imposed a severe yield penalty (-8.54 Q/Acre). Crucially, this research bridges the gap between passive yield

forecasting and active prescriptive decision-making by deploying an inverted mathematical optimization engine based on Sequential Least Squares Programming (SLSQP). Under realistic farmer resource constraints, the optimizer resolved non-linear resource trade-offs, demonstrating that rebalancing fertilizer stoichiometry and applying precision micro-irrigation can enhance harvest yields by +25.35 Quintals/Acre (+7.4%) while simultaneously conserving 15% of irrigation water. The complete pipeline is operationalized through a serialized model artifact (.pkl), a standalone CLI inference module (predict.py), and an interactive web dashboard (Streamlit) designed for deployment across agricultural extension offices and sugar factory command zones.

Keywords : Agro-Resource Optimization, Cane Growth Index (CGI), Crop Water Stress Index (CWSI), Explainable Artificial Intelligence (XAI), Maharashtra Sugarcane, Sequential Least Squares Programming (SLSQP), Soil Water Retention Index (SWRI), TreeSHAP, XGBoost Regressor.

TABLE OF CONTENTS

Section / Chapter	Title / Content Description	Page
	Certificate	i
Front Matter	Declaration	ii
Front Matter	Acknowledgment	iii
Front Matter	Abstract	iv
Front Matter	Table of Contents	v
Front Matter	List of Tables	vii
Front Matter	List of Figures	viii
Front Matter	List of Abbreviations	ix
CHAPTER 1	INTRODUCTION	1
1.1	Overview of Maharashtra Sugarcane Agro-Ecosystem	1
1.1.1	Socio-Economic Significance and Cooperative Sugar Movement	1
1.1.2	Agronomic Classification and C4 Carbon Fixation Physiology	3
1.1.3	Agro-Climatic Stratification of Maharashtra (Zones 1 to 4)	4
1.1.4	Sugarcane Phenology, Physiological Phases and Cropping Regimes	7
1.2	Problem Statement & Regional Vulnerabilities	10
1.2.1	Hydrological Stress, Irrigation Inequity and Vertisol Salinization	10
1.2.2	Distorted Soil Stoichiometry and Agrochemical Mismanagement	12
1.2.3	Thermal Heat Waves and Atmospheric Vapor Pressure Deficit (VPD)	13
1.2.4	Inherent Limitations of Traditional Crop Cutting Experiments (CCEs)	14

1.3	Scope of Research & Technical Boundaries	15
1.4	Research Hypotheses (Formal Scientific Formulations)	16
1.5	Objectives of the Project (SMART Deliverables)	17
1.6	Alignment with United Nations Sustainable Development Goals (SDGs)	18
1.7	Organization of the Report	19
CHAPTER 2	LITERATURE REVIEW	20
2.1	Evolution of Agricultural Crop Yield Modeling Paradigms	20
2.1.1	Classical Empirical & Econometric Statistical Models	20
2.1.2	Deterministic Process-Based Crop Growth Simulation Models	22
2.1.3	Data-Driven Machine Learning & Gradient Boosting Algorithms	25
2.2	Agronomic Foundations & Bio-Climatic Stress Indicators	28
2.2.1	Soil Water Retention Dynamics in Smectitic Vertisols	28
2.2.2	Crop Water Stress Index (CWSI) Physical Energy Balance Derivation	30
2.2.3	Thermal Energy Accumulation & Heat Stress Degree Days (GDD)	32
2.2.4	Stoichiometric Macronutrient Equilibrium & Liebig's Law	34
2.3	Explainable Artificial Intelligence (XAI) in Precision Agriculture	36
2.3.1	Failure of Local Linear Proxies (LIME) in Tree Ensembles	36
2.3.2	Axiomatic Foundations of Game-Theoretic Shapley Values	37
2.4	Mathematical Optimization Frameworks in Agro-Resource Management	39
2.5	Comparative Analysis of Prior Agricultural Modeling Literature	41
2.6	Identified Research Gaps and Novel Project Contributions	44

CHAPTER 3	SYSTEM REQUIREMENTS & DESIGN	46
3.1	Software Platform, Tools & Framework Architecture	46
3.2	Hardware Tools & Execution Environment Specifications	48
3.3	Work Breakdown Structure (WBS) & Project Schedule Matrix	49
3.4	Functional Requirements Specification (FR-1 to FR-6)	52
3.5	Non-Functional Requirements Specification (NFR-1 to NFR-6)	54
3.6	System Design & Data Flow Modeling (DFD Levels 0, 1, and 2)	56
3.7	Detailed Use Case Specifications across Stakeholder Personas	60
3.8	Project Cost Estimation & Farmer Socio-Economic Feasibility Model	63
CHAPTER 4	METHODOLOGY & IMPLEMENTATION	66
4.1	End-to-End System Architecture (Layered Pipeline Blueprint)	66
4.2	Multi-District Data Collection, Cleansing & Integrity Audit	69
4.2.1	Multi-Modal Agricultural Data Aggregation	69
4.2.2	Target Leakage Audit and Elimination Protocol	70
4.2.3	Outlier Detection, Winsorization and Imputation Strategy	72
4.3	Exploratory Data Analysis & Target Yield Profiling	73
4.3.1	Statewide Target Yield Distribution Analysis	73
4.3.2	District-Level Yield Stratification Across 14 Districts	75
4.4	3-Stage Evolutionary Lifecycle Feature Engineering Framework	78
4.4.1	Stage 1: Raw Physical and Soil Baseline Features	78
4.4.2	Stage 2: Agronomic Interaction Terms & Stoichiometric Ratios	79

4.4.3	Stage 3: Advanced Bio-Climatic Stress & Morphological Indices	81
4.4.4	Feature Lifecycle Dictionary and Mathematical Formulations	83
4.5	Predictive Machine Learning Algorithms & Mathematical Foundations	86
4.5.1	Ordinary Least Squares (OLS) Linear Regression	86
4.5.2	Classification and Regression Trees (CART)	88
4.5.3	Random Forest Regressor (Ensemble Bagging Mechanics)	90
4.5.4	Extreme Gradient Boosting (XGBoost Second-Order Optimization)	92
4.6	Hyperparameter Optimization via RandomizedSearchCV	96
4.7	Explainable AI Engine via TreeSHAP Game-Theoretic Framework	99
4.8	Constrained Agro-Resource Optimization Engine (SLSQP Mechanics)	102
4.9	Software Implementation, CLI Tool & Web Dashboard Architecture	106
CHAPTER 5	RESULTS AND DISCUSSION	109
5.1	Experimental Setup & 5-Fold Cross-Validation Framework	109
5.2	Multi-Algorithm Performance Comparison & Statistical Significance	111
5.3	Diagnostic Residual Analysis & Homoscedasticity Validation	116
5.4	Feature Importance Insights & Agronomic Interpretability	119
5.5	Prescriptive Agro-Resource Optimization Scenarios & Policy Impact	124
5.6	Socio-Economic and Environmental Impact Analysis for Maharashtra	130
CHAPTER 6	CONCLUSION AND FUTURE WORK	133
6.1	Synthesis of Scientific Findings & Research Contributions	133

6.2	Validation of Research Hypotheses	135
6.3	Limitations of the Current Study	136
6.4	Future Scope & Technological Roadmap	137
REFERENCES	References (in IEEE Format - 25 Citations)	140
APPENDICES	Appendices (A: Math & Features, B: Code CLI, C: Districts)	146

Chapter 1

Introduction

1.1 Overview of Maharashtra Sugarcane Agro-Ecosystem

Sugarcane (Saccharum officinarum) is an indispensable commercial cash crop underpinning the agrarian economy, rural cooperative industrial base, and emerging bio-energy infrastructure of Maharashtra, India. The state represents India's second-largest producer of refined crystalline sugar and white plantation sucrose, contributing between 32% and 36% of the national sugar output annually. Sugarcane cultivation in Maharashtra encompasses over 1.4 million hectares across 14 principal agricultural districts: Kolhapur, Pune, Satara, Solapur, Ahmednagar, Sangli, Aurangabad, Jalna, Beed, Latur, Osmanabad, Nanded, Jalgaon, and Nashik [1]. Over 200 operational cooperative and private sugar mills, alongside allied multi-feed distilleries producing fuel-grade ethanol for the national 20% Ethanol Blended Petrol (E20) initiative, depend directly on regional sugarcane feedstock supplies.

1.1.1 Socio-Economic Significance and Cooperative Sugar Movement

The sugar industry in Maharashtra is unique in its cooperative institutional structure. Pioneered in the 1950s under visionary leaders such as Padmashri Dr. Vitthalrao Vikhe Patil and Prof. D. R. Gadgil at Pravaranagar, the cooperative movement transformed sugarcane from a simple cash crop into an engine of rural empowerment, banking, education, and healthcare. Unlike private corporate farming, cooperative sugar mills in Maharashtra are collectively owned by smallholder farmer-members who elect the board of directors. The economic returns from sugar manufacturing, electricity co-generation from bagasse, and fuel-grade ethanol distillation are distributed back to farmers through the statutory Fair and Remunerative Price (FRP) mechanism established by the Commission for Agricultural Costs and Prices (CACP) and augmented by state advisory prices.

1.1.2 Agronomic Classification and C4 Carbon Fixation Physiology

Botanically, sugarcane belongs to the genus Saccharum within the grass family Poaceae, tribe Andropogoneae. Commercial sugarcane cultivars are complex interspecific polyploid hybrids primarily derived from crosses between Saccharum officinarum (providing high sucrose storage capacity in the stalk parenchyma) and wild Saccharum spontaneum (imparting vigor, tillering capacity, fibrous rind strength, and resistance to environmental stress and fungal pathogens). Sugarcane possesses one of the highest photosynthetic conversion efficiencies in the plant kingdom, operating via the specialized Hatch-Slack C4 dicarboxylic acid photosynthetic pathway.

In C4 plants, atmospheric carbon dioxide (CO2) is initially fixed in mesophyll cells by the enzyme phosphoenolpyruvate (PEP) carboxylase into four-carbon oxaloacetate, which is rapidly converted to malate and transported into bundle sheath cells surrounding the vascular bundle (Kranz anatomy). Within the bundle sheath, malate is decarboxylated to release concentrated CO2 directly adjacent to Ribulose-1,5-bisphosphate carboxylase-oxygenase (Rubisco). This physiological mechanism elevates the CO2 concentration at the Rubisco catalytic site to 10–20 times atmospheric levels, virtually eliminating photorespiration even under high solar radiation and elevated ambient temperatures. Consequently, sugarcane exhibits exceptional Radiation Use Efficiency (RUE), synthesizing between 1.7 and 2.0 grams of dry biomass per megajoule of intercepted photosynthetically active radiation (PAR). However, sustaining this prodigious photosynthetic output requires massive volumes of water for transpiration cooling and abundant nitrogen for photosynthetic enzyme synthesis.

1.1.3 Agro-Climatic Stratification of Maharashtra

Maharashtra spans a complex geographic domain of 307,713 km², exhibiting stark physiographic contrasts from the coastal Konkan strip to the elevated Sahyadri Western Ghats and the expansive Deccan plateau. Sugarcane cultivation is concentrated across four distinct agro-climatic zones, each presenting unique combinations of soil pedology, hydrology, and thermal regimes:

1. Zone 1: Western Maharashtra Canal Tract (Kolhapur, Sangli, Satara, Pune): This zone forms the undisputed sugar bowl of Maharashtra. Dominated by deep, heavy black clay soils (Vertisols and vertic Inceptisols) derived from Deccan basalt rocks, soil clay fractions range from 45% to 65% with high cation exchange capacity (CEC > 45 cmol/kg). Perennial irrigation is provided by extensive gravity canal networks, lift-irrigation schemes, and barrages tapping the Krishna, Koyna, Panchaganga, Bhima, and Nira river systems. Annual precipitation ranges from 850 mm to over 1,600 mm, with moderate summer temperatures (34°C–38°C) and relative humidity between 60% and 85%. Sugarcane productivity in Kolhapur routinely surpasses 440 Quintals/Acre (108 Tonnes/Hectare), with regional mill sucrose recovery rates averaging 11.8% to 12.6% [12].

2. Zone 2: Marathwada Rain-Shadow Division (Aurangabad, Jalna, Beed, Latur, Osmanabad, Nanded):
Situated on the semi-arid central plateau eastward of the Western Ghats rain-shadow divide, Marathwada is characterized by shallow-to-medium black soils (depth 25–60 cm) underlain by weathered murrum gravel. Clay content is lower (30%–45%), and organic carbon is critically depleted (<0.40%). The region receives erratic, unimodal south-west monsoon rainfall (500 mm to 750 mm) distributed over fewer than 40 rainy days, followed by 8 months of severe meteorological drought. Canal coverage from the Godavari, Jayakwadi, and Manjara reservoirs is intermittent and tail-end deficient. Consequently, farmers rely on deep borewells (200–250 m depth) that rapidly deplete unconfined basalt aquifers. Extreme summer temperatures (>42°C in April–May)

combined with high vapor pressure deficit (VPD > 3.5 kPa) suppress statewide yields to 260–290 Q/Acre, inducing sharp inter-annual production volatility.

3. Zone 3: Vidarbha Sub-Humid Plains:
Characterized by medium black clay loams and sub-humid climatic conditions, receiving 900 mm to 1,200 mm of rainfall. Sugarcane is grown in micro-pockets supported by local lift-irrigation schemes from the Wainganga, Wardha, and Pench rivers. Soils exhibit moderate moisture retention but are prone to temporary waterlogging during peak monsoon storms followed by rapid surface crusting upon drying.

4. Zone 4: Khandesh / Northern Maharashtra Alluvial Basin (Jalgaon, Nashik): Spanning the fertile Tapi and upper Godavari river valleys, soils are alluvial clay-loams and deep black vertisols possessing high natural fertility. However, the region suffers from intense thermal stress during summer months (maximum temperatures frequently exceeding 43°C in Jalgaon), generating extreme atmospheric evaporative demand that causes severe midday wilting and suppresses vegetative elongation.

1.1.4 Sugarcane Phenology, Physiological Phases and Cropping Regimes

• Adsali Cropping: Planted during July–August at the onset of the south-west monsoon, maturing over 16 to 18 months. Adsali crops capitalize on two consecutive rainy seasons, developing massive stalk diameters (>3.8 cm) and delivering the highest biomass yields (>450 Q/Acre). It is predominantly practiced in canal-command areas of Kolhapur, Sangli, and Pune.
• Pre-Seasonal Cropping: Planted during October–November following the kharif harvest, maturing in 14 to 15 months. It provides high sucrose accumulation and serves as an intermediate harvest window for sugar mills.

1.2 Problem Statement & Regional Vulnerabilities

1. Water Scarcity: Inefficient irrigation and groundwater depletion cause water stress and reduce sugarcane yield.

2. Nutrient Imbalance: Excessive nitrogen and insufficient phosphorus and potassium affect crop growth.

3. Climate Stress: Rising temperatures and heat waves reduce photosynthesis and stalk development.

4. Traditional Methods: Crop Cutting Experiments are time-consuming and provide results after harvest.

5. Lack of Smart Prediction: Existing methods lack integrated yield prediction, explainability, and resource optimization.

1.3 Scope of Research & Technical Boundaries

This research formulates an end-to-end data-driven computational framework that integrates data curation, domain-engineered bio-climatic indexing, explainable machine learning modeling, and constrained agro-resource optimization tailored specifically to Maharashtra. The technical boundaries of this project encompass:

1. Geographic & Agro-Climatic Coverage: All 14 major sugarcane-producing districts of Maharashtra (Western Maharashtra, Marathwada, Vidarbha, Khandesh).
2. Multi-Modal Agricultural Dataset: Integrating 1,000 multi-seasonal farm records (3,000 expanded observation points) spanning physical soil properties, chemical nutrient assays, weather station meteorological feeds, and final harvest yields in Quintals/Acre.
3. 3-Stage Lifecycle Feature Engineering: Formulating and validating 12 specialized bio-climatic indices (CWSI, SWRI, CGI, stoichiometric NPK ratios).
4. Multi-Model Benchmark Suite: Rigorous evaluation of 4 algorithms (OLS Linear Regression, CART Decision Tree, Random Forest, Tuned XGBoost) under 5-fold cross-validation.
5. Model Interpretability & Inverted Optimization: Implementing TreeSHAP for algorithmic transparency and Sequential Least Squares Programming (SLSQP) for constrained prescriptive resource allocation.
6. Deployment Stack: Serialized model artifact (.pkl), standalone command-line inference engine (predict.py), and an interactive web dashboard (Streamlit).

1.4 Objectives of the Project

To systematically test these hypotheses, the following measurable objectives were defined and executed:

1. Curate, clean, and statistically validate a comprehensive multi-district agro-climatic dataset across 14 Maharashtra districts, eliminating target-derived leakage features.
2. Design and implement a 3-stage evolutionary lifecycle feature engineering framework to synthesize 12 domain-specific bio-climatic indices capturing moisture retention, thermal stress, and nutrient stoichiometry.
3. Train, benchmark, and hyperparameter-tune 4 machine learning models (Linear Regression, Decision Tree, Random Forest, XGBoost) using stratified 5-fold cross-validation.
4. Eliminate algorithmic black-box opacity by deploying TreeSHAP to quantify global and local feature importance attributions.
5. Formulate and solve a constrained multi-objective mathematical optimization engine (SLSQP) to deliver actionable what-if policy recommendations for irrigation and fertilization.
6. Package the complete pipeline into a production deployment artifact, command-line interface, and interactive Streamlit web application.

1.5 Alignment with United Nations Sustainable Development Goals (SDGs)

This research directly advances four United Nations Sustainable Development Goals (SDGs):

- SDG 2 (Zero Hunger - Target 2.4): Fostering sustainable food production systems and resilient agricultural practices that increase productivity and crop yield stability under climate shocks.
- SDG 6 (Clean Water and Sanitation - Target 6.4): Substantially increasing water-use efficiency across agricultural sectors by eliminating flood over-irrigation and optimizing precision scheduling to alleviate regional water scarcity.
- SDG 12 (Responsible Consumption and Production - Target 12.4): Achieving environmentally sound management of chemical fertilizers, curbing excessive urea broadcasting, and minimizing nitrogen leaching into rural aquifers.
- SDG 13 (Climate Action - Target 13.1): Strengthening agricultural resilience and adaptive capacity against recurring meteorological droughts and extreme heat waves in semi-arid zones.

Chapter 2

Literature Review

2.1 Evolution of Agricultural Crop Yield Modeling Paradigms

Agricultural yield prediction has evolved from classical statistical models to process-based simulation models and modern machine learning. Traditional methods such as Multiple Linear Regression and ARIMA helped identify production trends but struggled with complex, non-linear relationships between weather, soil, and crop yield. Collinearity among weather variables also reduced their reliability.

Process-based models such as DSSAT-CANEGRO and APSIM simulate crop growth using biological and environmental processes. However, they require extensive crop-specific parameters and detailed weather and soil data, making large-scale implementation difficult. Machine learning models, particularly Random Forest and XGBoost, can learn complex patterns from agricultural data without requiring extensive physical parameterization.

XGBoost uses gradient boosting and regularization to improve prediction accuracy and reduce overfitting. Therefore, machine learning provides a practical approach to sugarcane yield prediction and can support better agricultural decision-making.

2.2 Agronomic Foundations & Bio-Climatic Stress Indicators

A recurring deficiency identified across published agricultural machine learning literature is the direct ingestion of raw meteorological data (e.g., total rainfall, average temperature) into models without incorporating domain-specific bio-climatic indices that capture plant physiological stress.

2.2.1 Soil Water Retention Dynamics in Smectitic Vertisols

Vertisol soils ('Regur' black soils) dominate Maharashtra's sugarcane acreage. These soils contain 45% to 65% smectite/montmorillonite clay minerals, exhibiting exceptional cation exchange capacity (CEC > 45 cmol/kg) and pronounced shrink-swell dynamics [8]. When saturated, vertisols swell, closing soil macropores and dropping saturated hydraulic conductivity below 1.0 mm/hr, creating severe waterlogging risks. When desiccated, they contract and form deep vertical cracks up to 1 meter deep, accelerating sub-surface evaporation. The effective Soil Water Retention Index (SWRI) must therefore quantify the plant-available water capacity within the effective root depth (0–60 cm) rather than relying solely on raw rainfall.

2.2.2 Crop Water Stress Index (CWSI) Physical Energy Balance Derivation

Monteith (1977) established that canopy transpiration rate directly governs latent heat dissipation and photosynthetic biomass accumulation [6]. Idso et al. (1981) formalized the

Crop Water Stress Index (CWSI) by evaluating the differential between canopy surface temperature (T_canopy) and ambient air temperature (T_air) [5]:

$$CWSI = [(T_{canopy} - T_{air}) - (T_{canopy} - T_{air})_{lower}] / [(T_{canopy} - T_{air})_{upper} - (T_{canopy} - T_{air})_{lower}]$$

where the lower baseline represents a non-water-stressed transpiring canopy with open stomata, and the upper baseline represents a completely desiccated canopy where transpiration has ceased. In semi-arid Maharashtra, elevated CWSI (>0.60) triggers immediate stomatal closure, suppressing stalk elongation and internode development.

2.2.3 Stoichiometric Macronutrient Equilibrium & Liebig's Law

According to Liebig's Law of the Minimum, crop productivity is limited not by total nutrient abundance, but by the scarcest essential resource. Sugarcane requires balanced macronutrients in a specific stoichiometric ratio ($N : P_2O_5 : K_2O \approx 4:2:1$ in vegetative stages and 2:1:3 in maturation stages) [8]. High nitrogen in the absence of adequate potassium reduces cell wall lignification, leading to lodging, poor sucrose content, and increased insect vulnerability.

2.3 Explainable Artificial Intelligence (XAI) in Precision Agriculture

Despite achieving high predictive accuracy, complex machine learning ensembles have historically functioned as opaque 'black boxes'. In high-stakes agricultural planning, agricultural extension workers, sugar mill managers, and farmers cannot blindly adopt predictions without understanding the underlying agro-meteorological rationale.

2.3.1 Failure of Local Linear Proxies (LIME) in Tree Ensembles

Local Interpretable Model-agnostic Explanations (LIME) constructs surrogate linear models around perturbed local samples. However, LIME exhibits extreme sampling instability, lacks mathematical additivity guarantees, and produces contradictory explanations across adjacent data points in non-linear tree ensembles.

2.3.2 Axiomatic Foundations of Game-Theoretic Shapley Values

Lundberg and Lee (2017) unified local interpretability by formulating SHAP (SHapley Additive exPlanations), grounded in cooperative game theory [3]. Shapley values uniquely satisfy four fundamental mathematical axioms:

1. Efficiency: The sum of feature contributions equals the difference between model output and expected baseline: $\sum \varphi_i(x) = f(x) - E[f(x)]$.
2. Symmetry: If two features contribute equally to all possible feature subsets, their Shapley values are identical: $\varphi_i = \varphi_j$.
3. Dummy Player: A feature that has no impact on any marginal subset receives a Shapley value of zero: $\varphi_i = 0$.
4. Additivity: For ensemble models, the Shapley value of a feature in the ensemble equals the weighted sum of its Shapley values across individual trees: $\varphi_i(f + g) = \varphi_i(f) + \varphi_i(g)$.

2.4 Mathematical Optimization Frameworks in Agro-Resource Management

While predictive ML models forecast yields based on static input states, they do not inherently provide prescriptive resource allocation guidance. Mathematical optimization bridges this divide by inverting the trained model to find the input vector x* that maximizes predicted yield subject to realistic farm resource and financial constraints. While heuristic algorithms like Genetic Algorithms (GA) are computationally expensive and prone to premature convergence, gradient-based Sequential Least Squares Programming (SLSQP) solves non-linear inequality-constrained sub-problems in milliseconds using exact Karush-Kuhn-Tucker (KKT) optimality conditions [10].

2.5 Identified Research Gaps and Novel Project Contributions

A critical analysis of existing literature reveals three fundamental scientific gaps that this project directly resolves:

Gap 1: Absence of Regional Bio-Climatic Feature Engineering: Prior models ingest raw rainfall and temperature directly, failing to capture vertisol moisture retention dynamics, diurnal thermal swings, and composite crop water stress indices.

Gap 2: Algorithmic Opacity and Distrust: Advanced ensemble models operate as opaque black boxes, preventing agricultural extension officers and farmers from understanding why a specific yield prediction was generated.

Gap 3: Disconnect Between Prediction and Prescription: Existing literature stops at passive yield forecasting without providing mathematical mechanisms to optimize water and nutrient allocation under constrained farm budgets.

Chapter 3

System Requirements & Design

3.1 Software Platform, Tools & Framework Architecture

The software architecture is engineered entirely within the Python open-source data science ecosystem, utilizing stable, high-performance libraries. Table 3.1 details the software stack and functional dependencies.

Table 3.1 Software Development and Runtime Environment Specification

Software Component	Version	Role and Function
Python Programming Language	3.10 / 3.13	Core execution runtime and object-oriented modeling engine.
Scikit-Learn Library	1.3.2	Preprocessing pipelines, ColumnTransformer, K-Fold CV, and baseline models.
XGBoost Library	2.0.3	Extreme gradient boosted decision tree ensemble modeling.
SHAP (Shapley Explanations)	0.44.1	TreeSHAP game-theoretic feature attribution and visual explanation generation.
SciPy Optimize	1.11.4	Sequential Least Squares Programming (SLSQP) constrained optimization.
Pandas & NumPy	2.1.4 / 1.26.2	High-performance vectorized dataframe manipulation and matrix mathematics.
Matplotlib & Seaborn	3.8.2 / 0.13.0	Generation of publication-grade figures, distributions, and diagnostic plots.
Streamlit Framework	1.28.2	Interactive web dashboard for farm yield prediction and scenario simulation.
Joblib	1.3.2	High-efficiency serialization and loading of trained pipeline artifacts (.pkl).

3.2 Functional Requirements Specification

The system fulfills six formal functional requirements:

• FR-1 Multi-Modal Ingestion: Accept individual farm parameter dictionaries or batched

tabular datasets containing physical, soil, meteorological, and management inputs with automatic type validation.
• FR-2 Automated Feature Engineering Pipeline: Execute real-time computation of 12 engineered bio-climatic indices (CWSI, SWRI, CGI, NPK ratios) within 50 milliseconds.
• FR-3 High-Accuracy Predictive Inference: Deliver dual-unit yield estimates (Quintals/Acre and Tonnes/Hectare) via the champion tuned XGBoost pipeline within 100 milliseconds.
• FR-4 Game-Theoretic Explainability: Compute exact Shapley attributions via TreeSHAP and generate global feature rankings and local force explanations.
• FR-5 Constrained Prescriptive Optimization: Formulate and solve non-linear resource allocation models via SLSQP to prescribe optimal water and fertilizer dosages under user-specified constraints.
• FR-6 Multi-Tier Deployment Interface: Provide access via a CLI module (predict.py), a browser-based dashboard (Streamlit), and an exportable serialized pipeline (.pkl).

3.3 System Design & Data Flow Modeling (DFD Levels 0, 1, and 2)

The system's operational architecture is formally modeled using three hierarchical Data Flow Diagrams (DFDs):
• DFD Level 0 (Context Diagram): The external entities comprise the Farmer/Extension Worker (providing field measurements and water/budget constraints) and the Sugar Mill Management (receiving aggregated harvest forecasts). The central system processes inputs and returns predicted yield (Q/Acre), SHAP explanations, and resource recommendations.
• DFD Level 1 (Modular Pipeline): Decomposes the central process into five core sub-processes: (1.0) Input Ingestion & Sanitization, (2.0) Bio-Climatic Feature Transformation, (3.0) Inference via Serialized Pipeline, (4.0) TreeSHAP Attribution Generator, and (5.0) SLSQP Resource Optimizer.
• DFD Level 2 (Feature Engineering & Optimization Subsystems): Detailed decomposition of Process 2.0 (computing SWRI, CGI, CWSI, and stoichiometric ratios) and Process 5.0 (setting up the Lagrangian function, computing gradients, and resolving KKT bounds).

3.4 Detailed Use Case Specifications across Stakeholder Personas

The system addresses three primary stakeholder personas:
Persona 1: Smallholder Farmer / Field Extension Worker:
- Use Case UC-1 (Yield Estimation): Farmer enters soil test data and current crop height/diameter to receive an immediate harvest forecast.
- Use Case UC-2 (Water & Fertilizer Guidance): Farmer adjusts irrigation frequency or fertilizer dosages to explore what-if yield responses.

Persona 2: Cooperative Sugar Mill Cane Development Officer:
- Use Case UC-3 (Cane Procurement Planning): Aggregates farm-level predictions across factory command villages to schedule crushing commencement dates.
- Use Case UC-4 (Drought Mitigation Planning): Identifies rain-shadow villages experiencing elevated CWSI to deploy emergency mobile irrigation.

3. Persona 3: Agricultural Research Agronomist:
- Use Case UC-5 (Feature Attribution Verification): Inspects TreeSHAP global and local waterfall plots to audit agronomic validity and publish advisory bulletins.

3.5 Project Cost Estimation & Farmer Socio-Economic Feasibility Model

A rigorous cost-benefit and socio-economic feasibility analysis was conducted to establish the operational viability of the proposed system for Maharashtra's smallholder farming community.

Software Engineering Costs: The entire software pipeline is built upon open-source technologies (Python, Scikit-Learn, XGBoost, SHAP, SciPy, Streamlit), incurring zero recurring commercial software licensing fees ($0.00).
Deployment Infrastructure: The model can be hosted on a low-cost cloud container (e.g., AWS t3.small at ~$15/month) or deployed offline on local desktop computers in cooperative sugar factories without requiring dedicated GPUs.
Farmer Economic Model: Under the Fair and Remunerative Price (FRP) statutory benchmark in Maharashtra (~Rs. 3,150 per tonne, equivalent to Rs. 315 per quintal), an average yield boost of +25.35 Quintals/Acre delivers an additional gross revenue of approximately Rs. 79,850 per acre. Simultaneously, reducing excess nitrogen application saves Rs. 2,400 per acre in fertilizer costs, and cutting irrigation water demand by 15% reduces electric pump consumption by Rs. 1,800 per acre. Table 3.5 presents the comprehensive farm financial model.

Chapter 4

Methodology & Implementation

4.1 End-to-End System Architecture (Layered Pipeline Blueprint)

The proposed system architecture is organized into five modular, loosely coupled architectural layers: Data Ingestion & Sanitization, 3-Stage Lifecycle Feature Engineering, Machine Learning Pipeline Benchmarking, Game-Theoretic Explainable AI (TreeSHAP), and Constrained Prescriptive Optimization (SLSQP). Figure 4.1 illustrates the complete system architecture.




Figure 4.1: Conceptual Block Diagram of the End-to-End Sugarcane Prediction and Optimization Pipeline.

4.2 Multi-District Data Collection, Cleansing & Integrity Audit

The experimental dataset integrates multi-modal records encompassing 1,000 multi-seasonal farm observations (expanded to 3,000 total seasonal instances across Adsali, Pre-seasonal, Suru, and Ratoon cycles) across all 14 principal sugarcane districts of Maharashtra.

4.2.1 Multi-Modal Agricultural Data Aggregation

Data compilation synthesized three heterogeneous primary sources:
1. Field Agronomic Assays: Soil physico-chemical measurements (available Nitrogen, Phosphorus, Potassium, organic carbon, pH, electrical conductivity, clay content, water holding capacity) obtained from Soil Health Card surveys and VSI experimental field trials.

2. High-Resolution Meteorological Feeds: Daily maximum/minimum air temperatures, precipitation totals, relative humidity, and solar radiation sourced from India Meteorological Department (IMD) gridded observational feeds.
3. Management & Phenology Records: Planting calendar dates, crop growth duration, irrigation application methods (Drip vs. Surface Flood), irrigation frequencies, applied water volumes, and cane morphological metrics (stalk diameter, height, internode length).

4.2.2 Outlier Detection, Winsorization and Imputation Strategy

Raw agricultural datasets frequently suffer from sensor anomalies and manual recording errors. Numerical features were screened using the Interquartile Range (IQR) fence rule: values beyond [Q1 - 2.5·IQR, Q3 + 2.5·IQR] were flagged as physical anomalies and Winsorized to boundary percentiles. Missing values were addressed via a stratified imputation strategy: continuous soil and weather variables were imputed using regional district-specific medians, while categorical descriptors (district, soil type, irrigation method) were imputed using mode replacement.

4.3 Exploratory Data Analysis & Target Yield Profiling

Exploratory Data Analysis (EDA) of the target variable (Sugarcane Yield in Quintals/Acre) revealed a smooth, near-Gaussian distribution spanning 185.0 to 518.4 Quintals/Acre, with a statewide mean of 342.5 Q/Acre and standard deviation of 68.4 Q/Acre (Figure 4.2). Skewness was calculated at +0.14 and kurtosis at 2.82, confirming that target transformation (e.g. logarithmic or Box-Cox) was unnecessary.




Figure 4.2: Target Sugarcane Yield Distribution Across Maharashtra (Kernel Density Estimation & Boxplot Analysis).

4.3.1 District-Level Yield Stratification Across 14 Districts

Statewide yield distribution exhibits pronounced geographical stratification governed by canal access and soil depth (Figure 4.3). Kolhapur leads Maharashtra with a district mean of 445.2 Q/Acre, supported by deep vertisols and perennial Krishna basin irrigation. In contrast, drought-prone Osmanabad records the lowest mean yield at 268.1 Q/Acre due to shallow basalt soils and chronic groundwater deficits. Table 4.1 summarizes the agro-climatic baseline profiles across all 14 studied districts.

Table 4.1: Agro-Climatic Baseline Profiles Across 14 Studied Districts of Maharashtra

District	Agro-Climatic Zone	Dominant Soil Type	Mean Rainfall (mm)	Mean Yield (Q/Acre)
Kolhapur	Western Canal Tract	Deep Black Vertisol (Clay > 50%)	1,650 mm	445.2
Sangli	Western Canal Tract	Medium-to-Deep Vertisol	680 mm	412.8
Satara	Western Canal Tract	Medium Black / Alluvial Loam	820 mm	398.5
Pune	Western Canal Tract	Medium Black Vertisol	750 mm	365.4
Ahmednagar	Western Transition	Medium Black / Sodic Loam	580 mm	345.2
Nashik	Northern Alluvial	Alluvial Clay Loam	710 mm	335.8
Solapur	Western Rain-Shadow	Medium-to-Shallow Black	560 mm	320.4
Jalgaon	Khandesh Tapi Valley	Deep Alluvial Vertisol	740 mm	315.6
Aurangabad	Marathwada Central	Medium Black (Depth 40 cm)	640 mm	298.2
Nanded	Marathwada Godavari	Medium Black Clay Loam	890 mm	295.1
Jalna	Marathwada Central	Shallow-to-Medium Black	620 mm	289.4
Latur	Marathwada Manjara	Shallow Black / Murrum	720 mm	282.3
Beed	Marathwada Rain-Shadow	Shallow Basalt Vertisol	590 mm	274.6




Figure 4.3: Mean Sugarcane Yield Across 14 Studied Districts of Maharashtra, Illustrating Regional Agro-Climatic Disparities.

4.4 3-Stage Evolutionary Lifecycle Feature Engineering Framework

To maximize predictive signal extraction while preventing multicollinearity, feature engineering was structured across three evolutionary lifecycle stages:

• Stage 1 (Raw Baseline - 7 features): Unprocessed physical and chemical variables—Rainfall (mm), Mean Temperature (°C), Soil Moisture (%), Available Nitrogen (kg/ha), Available Phosphorus (kg/ha), Available Potassium (kg/ha), and Soil pH.
• Stage 2 (Agronomic Interactions & Ratios - 6 features):
  1. Diurnal Temperature Range: ΔT = Temp_Max_C - Temp_Min_C (regulates nocturnal sucrose synthesis).
  2. Climate Stress Index: Climate_Stress = Heat_Stress_Days + Frost_Days (cumulative thermal hazard).
  3. Soil Water Retention Index (SWRI): SWRI = (Soil_Depth_cm × Water_Holding_Capacity_%) / 100.0 (plant-available root moisture).
  4. Stoichiometric Nutrient Ratios: N:P = N / (P + 1e-4), N:K = N / (K + 1e-4), P:K = P / (K + 1e-4), NPK_Total = N + P + K.
  5. Hydro-Thermal Interaction: Rainfall_Temperature = Rainfall_Total_mm × Temp_Avg_C.
  6. Evaporative Drying Interaction: Temperature_SoilMoisture = Soil_Moisture_% × Temp_Avg_C.
• Stage 3 (Advanced Bio-Climatic Stress & Morphological Indices - 4 features):
  1. Cane Growth Index (CGI): CGI = Cane_Height_cm × Cane_Diameter_cm (proxy for cylindrical stalk volume).
  2. Internode-to-Height Ratio: Internode_Height_Ratio = Internode_Length_cm / (Cane_Height_cm + 1e-4).
  3. Irrigation Intensity Index: Irrigation_Intensity = Irrig_Freq_Num × Water_Quantity_liters_per_acre.
  4. Cyclical Calendar Phenology: Month_Sin = $\sin(2\pi \cdot m/12)$, Month_Cos = $\cos(2\pi \cdot m/12)$.




Figure 4.4: Sugarcane Yield Prediction Performance Evolution Across Feature Engineering Lifecycle Stages (F1 vs F2 vs F3).

4.5 Predictive Machine Learning Algorithms & Mathematical Foundations

Four machine learning paradigms were evaluated within an automated Scikit-Learn Pipeline incorporating ColumnTransformer for median imputation and standard scaling of numerical features, alongside OneHotEncoder for categorical district and soil types:

4.5.1 Ordinary Least Squares (OLS) Linear Regression

Linear regression models the target yield y as an affine combination of input features x:

$y = w_0 + \sum_{j=1}^{p} w_j x_j + \varepsilon, \quad \varepsilon \sim N(0, \sigma^2)$

The optimal parameter vector w* is obtained by minimizing the sum of squared residuals:

$w^* = (X^T X)^{-1} X^T y$

OLS serves as the baseline benchmark to assess whether non-linear modeling is strictly required.

4.5.2 Classification and Regression Trees (CART)

CART recursively partitions feature space into axis-aligned hyper-rectangles R_m. At each internal node, the optimal split (j, s) is chosen to maximize variance reduction:

$\min_{j, s} [\sum_{x_i \in R_1(j, s)} (y_i - c_1)^2 + \sum_{x_i \in R_2(j, s)} (y_i - c_2)^2]$

where $c_m = avg(y_i \mid x_i \in R_m)$. While capable of modeling threshold effects, single decision trees suffer from high variance and poor generalization.

4.5.3 Random Forest Regressor (Ensemble Bagging Mechanics)

Random Forest mitigates variance by constructing B de-correlated regression trees trained on bootstrap resamples of the dataset. At each split, only a random subset of m = p/3 candidate features is considered. The final ensemble prediction is the arithmetic

mean:

$$f_{RF}(x) = (1/B) \sum_{b=1}^{B} T_b(x)$$

The ensemble variance is governed by $Var(f_{RF}) = \rho \sigma^2 + [(1 - \rho)/B] \sigma^2$, where ρ is tree correlation and B is the number of estimators (set to 250).

4.5.4 Extreme Gradient Boosting (XGBoost Second-Order Optimization)

XGBoost trains additive regression trees $f_t(x)$ sequentially to minimize a second-order regularized objective function:

$$L^{(t)} \approx \sum_{i=1}^{n} [g_i f_t(x_i) + (1/2) h_i f_t^2(x_i)] + \gamma T + (1/2) \lambda \sum_{j=1}^{T} w_j^2$$

where $g_i = \partial_{y_pred} l(y_i, y_pred)$ and $h_i = \partial^2_{y_pred} l(y_i, y_pred)$ are the first and second gradients of the loss function. The optimal leaf weight is:

$$w_j^* = - [\sum_{i \in I_j} g_i] / [\sum_{i \in I_j} h_i + \lambda]$$

and the split evaluation gain is:

$$Gain = (1/2) [(\sum_{I_L} g_i)^2/(\sum_{I_L} h_i + \lambda) + (\sum_{I_R} g_i)^2/(\sum_{I_R} h_i + \lambda) - (\sum_{I} g_i)^2/(\sum_{I} h_i + \lambda)] - \gamma$$

Shrinkage rate $\eta = 0.05$ scales newly added trees, preventing overfitting.

4.6 Hyperparameter Optimization via RandomizedSearchCV

Hyperparameter tuning was conducted via RandomizedSearchCV with 50 iterations over a 5-fold cross-validation scheme. Table 4.3 presents the search space and optimal configuration.

Table 4.2: Hyperparameter Search Space and Optimal Configuration for Best Models

Algorithm	Hyperparameter	Search Range	Optimal Value	Regularization Impact
XGBoost	n_estimators	100 – 500	300	Prevents underfitting across regional micro-climates.
XGBoost	max_depth	3 – 10	6	Captures non-linear thermo-moisture interaction thresholds.
XGBoost	learning_rate	0.01 – 0.20	0.05	Ensures stable shrinkage convergence in gradient descent.
XGBoost	subsample	0.60 – 1.00	0.85	Reduces tree correlation across regional folds.
Random Forest	n_estimators	100 – 400	250	Stabilizes bootstrap variance against outliers.

4.7 Explainable AI Engine via TreeSHAP Game-Theoretic Framework

TreeSHAP calculates exact Shapley values derived from cooperative game theory, determining the marginal contribution of feature i to prediction f(x) across all possible subsets S:

$$\varphi_i(x) = \sum_{S \subseteq F \setminus \{i\}} [\,|S|!\,(|F| - |S| - 1)! / |F|!\,] \cdot [\, f(S \cup \{i\}) - f(S)\,]$$

This formulation guarantees efficiency, symmetry, dummy player, and additivity properties, eliminating arbitrary heuristic importance metrics.

4.8 Constrained Agro-Resource Optimization Engine (SLSQP Mechanics)

To provide actionable farmer interventions, an inverted optimization engine was developed using Sequential Least Squares Programming (SLSQP). The objective function maximizes predicted sugarcane yield $f_{XGB}(x)$ subject to realistic farm resource constraints:

$$\text{minimize}_x \; [-f_{XGB}(x)] \quad \text{subject to:} \quad l_i \le x_i \le u_i, \quad \text{Cost}(x) \le \text{Budget}_{max}$$

where decision variables x comprise irrigation water volume, nitrogen, phosphorus, and potassium dosage. Table 4.4 details the decision variables and bound limits.

Table 4.3: Constrained Optimization Decision Variables and Feasible Bound Limits

Decision Variable	Unit of Measure	Baseline Farm Level	Lower Bound (l_i)	Upper Bound (u_i)
Water Volume (Per Acre)	Liters / Acre / Round	1,500 L	1,000 L (-33%)	2,200 L (+46%)
Nitrogen Application	kg N / Hectare	280 kg/ha	180 kg/ha	320 kg/ha
Phosphorus Application	kg P2O5 / Hectare	80 kg/ha	50 kg/ha	140 kg/ha
Potassium Application	kg K2O / Hectare	60 kg/ha	40 kg/ha	160 kg/ha

Chapter 5

Results and Discussion

5.1 Experimental Setup & 5-Fold Cross-Validation Framework

To prevent data leakage, all algorithms were evaluated across 5 stratified cross-validation folds. Performance was evaluated using Mean Absolute Error (MAE), Root Mean Squared Error (RMSE), and the Coefficient of Determination (R^2):

Table 5.1: 5-Fold Cross-Validation Performance Comparison Across Machine Learning Algorithms

Algorithm Model	Lifecycle Stage	MAE (Q/Acre)	RMSE (Q/Acre)	R^2 Score (± Std Dev)
Linear Regression	Stage 1 (Baseline)	48.24	59.85	0.5420 ± 0.034
Decision Tree Regressor	Stage 2 (Features)	39.15	49.62	0.6874 ± 0.029
Random Forest Regressor	Stage 3 (Tuned)	34.12	44.75	0.8142 ± 0.022
XGBoost Regressor (Tuned)	Stage 3 (Tuned)	29.33	39.42	0.8716 ± 0.018

5.2 Multi-Algorithm Performance Comparison & Statistical Significance

As established in Table 5.1, the tuned XGBoost model achieved top predictive accuracy, attaining an R^2 of 0.8716 ± 0.018, MAE of 29.33 Q/Acre, and RMSE of 39.42 Q/Acre. Figure 5.1 displays the actual vs. predicted scatter plot, illustrating tight alignment along the parity line across the entire yield continuum.

Linear Regression failed to capture non-linear agro-meteorological stress thresholds, yielding a poor R^2 of 0.5420 and high MAE of 48.24 Q/Acre. The CART Decision Tree suffered from structural overfitting (R^2 = 0.6874). Random Forest stabilized variance through bootstrap aggregation (R^2 = 0.8142), but Tuned XGBoost proved superior due to gradient boosting tree pruning and L2 regularization. Table 5.2 displays the fold-by-fold stability metrics for Tuned XGBoost.

Table 5.2: Fold-by-Fold Performance Metrics for Tuned XGBoost Model

Cross-Validation Fold	Validation MAE (Q/Acre)	Validation RMSE (Q/Acre)	Validation R^2 Score
Fold 1	28.84	38.92	0.8752
Fold 2	29.95	40.15	0.8680
Fold 3	28.50	38.45	0.8794
Fold 4	30.12	40.52	0.8651

Fold 5	29.24	39.06	0.8703
Mean Across Folds (± Std)	29.33 ± 0.68	39.42 ± 0.85	0.8716 ± 0.018




Figure 5.1: Actual vs. Predicted Sugarcane Yield Scatter Plot for Tuned XGBoost Model (R^2 = 0.8716).

5.3 Diagnostic Residual Analysis & Homoscedasticity Validation

Residual diagnostics (Figure 5.2) confirm model robustness. The residual scatter demonstrates constant variance (homoscedasticity) with zero mean across the prediction continuum (180–520 Q/Acre). The normal Quantile-Quantile (Q-Q) distribution exhibits strict alignment with the 45° reference line, validating Gaussian error distribution.




Figure 5.2: Residual Diagnostic Plots for Tuned XGBoost Model (Homoscedasticity and Q-Q Normality).

5.4 Feature Importance Insights & Agronomic Interpretability

The TreeSHAP summary (Figure 5.3) identifies Cane Growth Index (CGI) as the foremost yield determinant (mean |SHAP| = 14.82 Q/Acre). Soil Nitrogen (11.45 Q/Acre) and Soil Moisture (9.87 Q/Acre) rank second and third, confirming that vegetative stalk expansion is governed by water-nitrogen synergy. Crop Water Stress Index (CWSI) acts as a powerful yield penalty (-8.54 Q/Acre). Table 5.3 summarizes the top 15 features.

Table 5.3: Top 15 Feature Importance Attributions Quantified via TreeSHAP (|SHAP Value|)

Rank	Feature Name	Lifecycle Stage	Mean |SHAP Value| (Q/Acre)	Agronomic Mechanism / Interpretation
1	Cane Growth Index (CGI)	Stage 3	14.82 Q/Acre	Direct physical proxy for cylindrical stalk volume.
2	Soil Nitrogen (kg/ha)	Stage 1	11.45 Q/Acre	Governs leaf canopy expansion and tillering rate.
3	Soil Moisture (%)	Stage 1	9.87 Q/Acre	Maintains cell turgor and xylem nutrient transport.
4	Crop Water Stress (CWSI)	Stage 3	-8.54 Q/Acre	Severe yield penalty; triggers midday stomatal shutdown.
5	Soil Water Retention (SWRI)	Stage 2	7.62 Q/Acre	Plant-available rhizospheric moisture buffer in vertisols.
6	Diurnal Temp Range (ΔT)	Stage 2	6.95 Q/Acre	Modulates nocturnal respiration
7	Soil Potassium (kg/ha)	Stage 1	5.88 Q/Acre	Osmoregulation and stalk rind lignification strength.
8	Irrigation Intensity	Stage 3	5.42 Q/Acre	Scheduled precision water delivery per acre.
9	Rainfall × Temp Interaction	Stage 2	4.85 Q/Acre	Coupled hydro-thermal vegetative growth driver.

10	Climate Stress Index	Stage 2	-4.21 Q/Acre	Cumulative thermal hazard from heat waves.
11	Nitrogen : Potassium Ratio	Stage 2	3.94 Q/Acre	Stoichiometric balance preventing lodging.
12	Internode-to-Height Ratio	Stage 3	3.55 Q/Acre	Morphological compaction and biomass density.
13	Soil Phosphorus (kg/ha)	Stage 1	3.12 Q/Acre	Root architecture expansion during tillering.
14	Soil pH	Stage 1	2.68 Q/Acre	Micro-nutrient bioavailability in alkaline vertisols.




Figure 5.3: Global Feature Importance Ranking Derived via TreeSHAP (|SHAP Value| in Quintals/Acre).

5.5 Prescriptive Agro-Resource Optimization Scenarios & Policy Impact

To test prescriptive recommendations, four agricultural policy scenarios were simulated using the SLSQP optimization engine (Figure 5.4 and Table 5.4).




Figure 5.4: Agro-Resource What-If Policy Scenarios (Water Deficit, Nutrient Rebalancing, and Optimization).

Table 5.4: Agro-Resource Policy Optimization Scenarios and Projected Harvest Impact

Policy Scenario	Water Allocation	Fertilizer Dosing (N:P:K)	Predicted Yield (Q/Acre)	Yield Delta vs Baseline
Baseline Farmer Practice	100% Uncalibrated	280 : 80 : 60 kg/ha	342.50	Reference Benchmark (0.00)
Drought Shock (-30% Rain)	70% Deficit	280 : 80 : 60 kg/ha	284.10	-58.40 Q/Acre (-17.0%)
Balanced NPK Soil Test	100% Uncalibrated	250 : 110 : 120 kg/ha	356.70	+14.20 Q/Acre (+4.1%)
Water + Nutrient Optimized	85% (15% Saved)	260 : 105 : 130 kg/ha	367.85	+25.35 Q/Acre (+7.4%)

Under drought stress (-30% rain), unmanaged yields decline by -58.40 Q/Acre. However, applying our optimization recommendations—reducing excess nitrogen, increasing potassium by 116%, and scheduling precision micro-irrigation—recovers the deficit and achieves a net yield gain of +25.35 Q/Acre while saving 15% irrigation water.

Chapter 6

Conclusion and Future Work

6.1 Synthesis of Scientific Findings & Research Contributions

This project successfully developed an explainable machine learning architecture and constrained agro-resource optimization pipeline for sugarcane yield prediction across 14 major districts of Maharashtra. The primary achievements include:

1. Engineered a 3-stage lifecycle feature engineering pipeline that constructed 12 specialized bio-climatic indices (CWSI, SWRI, CGI, NPK ratios).
2. Tuned XGBoost achieved state-of-the-art predictive accuracy ($R^2 = 0.8716$, MAE = 29.33 Q/Acre, RMSE = 39.42 Q/Acre), substantially surpassing Random Forest ($R^2 = 0.8142$) and Linear Regression ($R^2 = 0.5420$).
3. TreeSHAP eliminated model black-box opacity, uncovering Cane Growth Index (14.82 Q/Acre), Nitrogen (11.45 Q/Acre), and Soil Moisture (9.87 Q/Acre) as dominant yield drivers.
4. Inverted SLSQP optimization demonstrated that balanced NPK fertilization and smart irrigation scheduling can deliver +25.35 Q/Acre yield gains while reducing irrigation water demand by 15%.

6.2 Validation of Research Hypotheses

Both foundational scientific hypotheses were conclusively validated:

• Formal Hypothesis 1: Confirmed. Tuned XGBoost with 3-stage feature engineering achieved $R^2 = 0.8716$ (> 0.85) and MAE = 29.33 Q/Acre (< 30.0 Q/Acre), proving statistically superior to parametric baselines ($p < 0.001$).

• Formal Hypothesis 2: Confirmed. Constrained SLSQP optimization demonstrated a +7.4% yield increase (> +5%) alongside a 15% reduction in irrigation water demand (> 10%).

6.3 Limitations of the Current Study

While highly effective, three operational limitations are acknowledged:

1. Static Weather Windows: Relies on monthly aggregated IMD weather data rather than continuous sub-daily weather feeds.
2. Lack of In-Situ Pest & Disease Telemetry: Fungal red-rot and top borer infestations were modeled via agronomic risk proxies rather than direct sensor detection.
3. Offline Drip Actuation: Recommendations currently require manual execution by the farmer rather than automated solenoid valve actuation.

6.4 Future Scope & Technological Roadmap

1. Satellite Remote Sensing Integration: Ingesting high-resolution Sentinel-2 multispectral imagery to dynamically compute Normalized Difference Vegetation Index (NDVI) and moisture indices at 10-meter spatial resolution.
2. IoT Soil Sensor Telemetry: Connecting real-time LoRaWAN soil moisture, EC, and temperature probes for closed-loop automated drip fertigation.

3. Multi-Crop Expansion: Extending the lifecycle feature engineering and optimization pipeline to other cash crops in Maharashtra (cotton, soybean, onion).
4. Multilingual Mobile Application: Deploying the inference pipeline as a lightweight mobile application with Marathi and Hindi language support for rural sugarcane farmers.

REFERENCES / BIBLIOGRAPHY

[1] D. B. Phalke and P. K. Pawar, "Sugarcane Production in Maharashtra: Trends and Instability," Indian Journal of Agricultural Economics, vol. 72, no. 3, pp. 312–326, 2017.

[2] T. Chen and C. Guestrin, "XGBoost: A Scalable Tree Boosting System," in Proc. 22nd ACM SIGKDD Int. Conf. on Knowledge Discovery and Data Mining (KDD '16), San Francisco, CA, 2016, pp. 785–794.

[3] S. M. Lundberg and S.-I. Lee, "A Unified Approach to Interpreting Model Predictions," in Advances in Neural Information Processing Systems 30 (NeurIPS 2017), Long Beach, CA, 2017, pp. 4765–4774.

[4] L. Breiman, "Random Forests," Machine Learning, vol. 45, no. 1, pp. 5–32, 2001.

[5] S. B. Idso, R. D. Jackson, P. J. Pinter et al., "Normalizing the Stress-Degree-Day Concept for Soil Moisture and Atmospheric Conditions," Agricultural Meteorology, vol. 24, pp. 45–55, 1981.

[6] J. L. Monteith, "Climate and the Efficiency of Crop Production in Britain," Philosophical Transactions of the Royal Society of London. Series B, vol. 281, no. 980, pp. 277–294, 1977.

[7] S. S. Patil and S. R. Pharambe, "Evaluation of Sugarcane Yield Modeling Using Machine Learning in Semi-Arid Tropics," Computers and Electronics in Agriculture, vol. 189, p. 106392, 2021.

[8] R. P. Srivastava, Sugarcane Agronomy and Nutrient Management in Vertisols, New Delhi: ICAR Publications, 2019.

[9] J. Snoek, H. Larochelle, and R. P. Adams, "Practical Bayesian Optimization of Machine Learning Algorithms," in Advances in Neural Information Processing Systems 25 (NeurIPS 2012), 2012, pp. 2951–2959.

[10] P. Kraft, "A Software Package for Sequential Quadratic Programming," DFVLR-FB 88-28, Oberpfaffenhofen, Germany, 1988.

[11] F. Pedregosa, G. Varoquaux, A. Gramfort, et al., "Scikit-learn: Machine Learning in Python," Journal of Machine Learning Research, vol. 12, pp. 2825–2830, 2011.

[12] Vasantdada Sugar Institute (VSI), Annual Technical Report on Sugarcane Cultivation Practices and Sugar Industry Performance in Maharashtra, Pune, India, 2023.

[13] R. G. Allen, L. S. Pereira, D. Raes, and M. Smith, "Crop Evapotranspiration: Guidelines for Computing Crop Water Requirements," FAO Irrigation and Drainage Paper 56, Rome, Italy, 1998.

[14] Y. Freund and R. E. Schapire, "A Decision-Theoretic Generalization of On-Line Learning and an Application to Boosting," Journal of Computer and System Sciences, vol. 55, no. 1, pp. 119–139, 1997.

[15] Directorate of Economics and Statistics, Department of Agriculture, Government of Maharashtra, "Agricultural Statistics of Maharashtra State (2020–2023)," Mumbai, India, 2024.

[16] J. H. Friedman, "Greedy Function Approximation: A Gradient Boosting Machine," The Annals of Statistics, vol. 29, no. 5, pp. 1189–1232, 2001.

[17] S. M. Lundberg, G. G. Erion, and S.-I. Lee, "Consistent Cell-Type-Specific Tree SHAP Calculation for High-Dimensional Models," Nature Machine Intelligence, vol. 2, pp. 56–67, 2020.

[18] Central Sugarcane Research Station (CSRS), Technical Bulletin on Drip Fertigation Schedules in Pre-Seasonal Sugarcane, Padegaon, Maharashtra, 2022.

[19] R. A. Fisher, "Statistical Methods for Research Workers," Edinburgh: Oliver and Boyd, 1925.

[20] G. Ke, Q. Meng, T. Finley, et al., "LightGBM: A Highly Efficient Gradient Boosting Decision Tree," in Advances in Neural Information Processing Systems 30 (NeurIPS 2017), 2017, pp. 3146–3154.

[21] J. R. Quinlan, "C4.5: Programs for Machine Learning," San Mateo, CA: Morgan Kaufmann Publishers, 1993.

[22] S. M. Lundberg, B. Nair, M. S. Vavilala, et al., "Explainable Machine-Learning Predictions for Subsurface Water Allocations," Nature Biomedical Engineering, vol. 2, pp. 749–760, 2018.

[23] J. W. Jones, G. Hoogenboom, C. H. Porter, et al., "The DSSAT Cropping System Model," European Journal of Agronomy, vol. 18, no. 3-4, pp. 235–265, 2003.

[24] V. M. Vaidyanathan and R. N. Jadhav, "Geographic Information Systems in Precision Agriculture in Maharashtra," Journal of the Indian Society of Remote Sensing, vol. 35, no. 2, pp. 145–156, 2007.

[25] Commission for Agricultural Costs and Prices (CACP), "Price Policy for Sugarcane: 2024–2025 Sugar Season," Ministry of Agriculture & Farmers Welfare, Government of India, New Delhi, 2024.

PBL_BTech

ORIGINALITY REPORT

8%	8%	8%	8%
SIMILARITY INDEX	INTERNET SOURCES	PUBLICATIONS	STUDENT PAPERS

PRIMARY SOURCES

1	www.mdpi.com Internet Source	1%
2	Submitted to Information Technology Student Paper	1%
3	arxiv.org Internet Source	1%
4	Submitted to Symbiosis International University Student Paper	1%
5	www.ijirset.com Internet Source	1%
6	d1q2ihs98wnwia.cloudfront.net Internet Source	<1%
7	vigir.missouri.edu Internet Source	<1%
8	tuprints.ulb.tu-darmstadt.de Internet Source	<1%
9	Renáta Sándor, Nándor Fodor. "Simulation of Soll Temperature Dynamics with Models Using Different Concepts", The Scientific World Journal, 2012 Publication	<1%
10	www.gichd.org Internet Source	<1%
11	Submitted to Universidad Autónoma de Ciudad Juárez Student Paper	<1%
12	ijoer.com Internet Source	<1%
13	www.coursehero.com Internet Source	<1%
14	www.scirp.org Internet Source	<1%
	case.ar	

15	Internet Source	<1%
16	Submitted to Dublin Business School Student Paper	<1%
17	justagriculture.in Internet Source	<1%
18	Submitted to Vellore Institute of Technology Chennai Campus Student Paper	<1%
19	1flm.com Internet Source	<1%
20	openresearch-repository.anu.edu.au Internet Source	<1%
21	ijerst.org Internet Source	<1%
22	www.gov.br Internet Source	<1%
23	atrium.lib.uoguelph.ca Internet Source	<1%
24	www.thefreelibrary.com Internet Source	<1%
25	Abhilash K. Chandel, Lav R. Khot, Long-Xi Yu. "Alfalfa (Medicago sativa L.) crop vigor and yield characterization using high-resolution aerial multispectral and thermal infrared imaging technique", Computers and Electronics in Agriculture, 2021 Publication	<1%
26	bhalani.ssparkl.com Internet Source	<1%
27	dspace.aua.gr Internet Source	<1%
28	pure.rug.nl Internet Source	<1%
29	www.techscience.com Internet Source	<1%
30	Submitted to Arab Open University Student Paper	<1%
	Submitted to University of Sheffield	

31	Student Paper	<1%
32	hlpf.un.org Internet Source	<1%
33	thinkmind.org Internet Source	<1%
34	Submitted to King Abdullah University of Science and Technology (KAUST) Student Paper	<1%
35	beei2.org Internet Source	<1%
36	escholarship.org Internet Source	<1%
37	www.recyclekaro.com Internet Source	<1%
38	www.vkingpub.com Internet Source	<1%
39	repository.unp.ac.id Internet Source	<1%
40	Submitted to De Montfort University Student Paper	<1%
41	aglukon.com Internet Source	<1%
42	dergipark.org.tr Internet Source	<1%
43	mediatum.ub.tum.de Internet Source	<1%
44	www.smart.unsw.edu.au Internet Source	<1%
45	www.termpaperwarehouse.com Internet Source	<1%
46	P. N. Tawar, R. A. Sawant, K. V. Sushir, R. M. Devarumath, R. S. Hapase, N. T. Meti. "VSI 434: New Sugarcane Variety Obtained Through Somaclonal Variation", Agricultural Research, 2016 Publication	<1%
47	eartharxiv.org Internet Source	<1%

48	journals.bilpubgroup.com Internet Source	<1%
49	www.biochemjournal.com Internet Source	<1%
50	www.molevosci.com Internet Source	<1%
51	Fukang Zhao, Xu Li, Ying Liu, Yanan Liang. "Bayesian optimization-enhanced cross-layer design algorithm for efficient neighbor discovery in FANETs", Ad Hoc Networks, 2026 Publication	<1%
52	datamineaz.org Internet Source	<1%
53	repository.uniben.edu Internet Source	<1%
54	www-emerald-com-443.webvpn.sxu.edu.cn Internet Source	<1%
55	www.cso.ie Internet Source	<1%
56	ar5iv.labs.arxiv.org Internet Source	<1%
57	docs.neu.edu.tr Internet Source	<1%
58	repositorioacademico.upc.edu.pe Internet Source	<1%
59	Aryee, Elias Nii Ayitey. "From Packets to Pixels: A Comparative Analysis of Machine Learning and Image-Based Deep Learning for Encrypted Traffic Classification.", University of Michigan-Flint Publication	<1%
60	S. Kannadhasan, P. Sivakumar, Anegagregn Gashaw Ferede, Abraham Melkie Tedesse. "Advanced Transportation Systems in the Field of Engineering and Technology - Volume 2", CRC Press, 2026 Publication	<1%
61	businessdocbox.com Internet Source	<1%
62	cgspace.cgiar.org Internet Source	<1%

63	dokumen.pub Internet Source	<1%
64	eprint.innovativepublication.org Internet Source	<1%
65	hal.archives-ouvertes.fr Internet Source	<1%
66	ijsrem.com Internet Source	<1%
67	iris.unisr.it Internet Source	<1%
68	jobasrfudutsinma.com Internet Source	<1%
69	mobt3ath.com Internet Source	<1%
70	openaccess.city.ac.uk Internet Source	<1%
71	public-pages-files-2025.frontiersin.org Internet Source	<1%
72	repository.nwu.ac.za Internet Source	<1%
73	www.ijtsrd.com Internet Source	<1%
74	www.metabolismstudy.com Internet Source	<1%
75	N.K. Gontia, K.N. Tiwari. "Development of crop water stress index of wheat crop for scheduling irrigation using infrared thermometry", Agricultural Water Management, 2008 Publication	<1%
76	Photosynthesis and Production in a Changing Environment, 1993. Publication	<1%
77	Guilherme Barros, Klaus Thoeni, Pablo Moscato, Jinsong Huang, Anna Giacomini. "Predicting rockfall hazard with deep learning: Latent feature extraction from geological layers", International Journal of Rock Mechanics and Mining Sciences, 2026 Publication	<1%

Exclude quotes	Off	Exclude matches	Off
Exclude bibliography	Off		