

Introduction to Logic

Department of Science and Mathematics

IIIT Guwahati.

August 20, 2020

Translating from English into Logical Expressions

1. Express the statement "Every student in this class has studied calculus" using predicates and quantifiers.

Translating from English into Logical Expressions

1. Express the statement "Every student in this class has studied calculus" using predicates and quantifiers.
 - "For every student in this class, that student has studied calculus."

Translating from English into Logical Expressions

1. Express the statement "Every student in this class has studied calculus" using predicates and quantifiers.
 - "For every student in this class, that student has studied calculus."
 - Introduce a variable x so that our statement becomes "For every student x in this class, x has studied calculus".

Translating from English into Logical Expressions

1. Express the statement "Every student in this class has studied calculus" using predicates and quantifiers.

- "For every student in this class, that student has studied calculus."
- Introduce a variable x so that our statement becomes "For every student x in this class, x has studied calculus".
- Introduce $C(x)$: " x has studied calculus."

Translating from English into Logical Expressions

1. Express the statement "Every student in this class has studied calculus" using predicates and quantifiers.

- "For every student in this class, that student has studied calculus."
- Introduce a variable x so that our statement becomes "For every student x in this class, x has studied calculus".
- Introduce $C(x)$: " x has studied calculus."

If the domain for x consists of the students in the class, we can translate our statement as

Translating from English into Logical Expressions

1. Express the statement "Every student in this class has studied calculus" using predicates and quantifiers.

- "For every student in this class, that student has studied calculus."
- Introduce a variable x so that our statement becomes "For every student x in this class, x has studied calculus".
- Introduce $C(x)$: " x has studied calculus."

If the domain for x consists of the students in the class, we can translate our statement as $\forall x C(x)$.

Translating from English into Logical Expressions

1. Express the statement "Every student in this class has studied calculus" using predicates and quantifiers.

- "For every student in this class, that student has studied calculus."
- Introduce a variable x so that our statement becomes "For every student x in this class, x has studied calculus".
- Introduce $C(x)$: " x has studied calculus."

If the domain for x consists of the students in the class, we can translate our statement as $\forall x C(x)$.

If we change the domain to consist of all people, we will need to express our statement as

Translating from English into Logical Expressions

1. Express the statement "Every student in this class has studied calculus" using predicates and quantifiers.

- "For every student in this class, that student has studied calculus."
- Introduce a variable x so that our statement becomes "For every student x in this class, x has studied calculus".
- Introduce $C(x)$: " x has studied calculus."

If the domain for x consists of the students in the class, we can translate our statement as $\forall x C(x)$.

If we change the domain to consist of all people, we will need to express our statement as

- "For every person x , if person x is a student in this class then x has studied calculus."

Translating from English into Logical Expressions

1. Express the statement "Every student in this class has studied calculus" using predicates and quantifiers.

- "For every student in this class, that student has studied calculus."
- Introduce a variable x so that our statement becomes "For every student x in this class, x has studied calculus".
- Introduce $C(x)$: " x has studied calculus."

If the domain for x consists of the students in the class, we can translate our statement as $\forall x C(x)$.

If we change the domain to consist of all people, we will need to express our statement as

- "For every person x , if person x is a student in this class then x has studied calculus."

If $S(x)$ represents the statement that person x is in this class, we see that our statement can be expressed as

Translating from English into Logical Expressions

1. Express the statement "Every student in this class has studied calculus" using predicates and quantifiers.

- "For every student in this class, that student has studied calculus."
- Introduce a variable x so that our statement becomes "For every student x in this class, x has studied calculus".
- Introduce $C(x)$: " x has studied calculus."

If the domain for x consists of the students in the class, we can translate our statement as $\forall x C(x)$.

If we change the domain to consist of all people, we will need to express our statement as

- "For every person x , if person x is a student in this class then x has studied calculus."

If $S(x)$ represents the statement that person x is in this class, we see that our statement can be expressed as $\forall x (S(x) \rightarrow C(x))$.

Translating from English into Logical Expressions

1. Express the statement "Every student in this class has studied calculus" using predicates and quantifiers.

- "For every student in this class, that student has studied calculus."
- Introduce a variable x so that our statement becomes "For every student x in this class, x has studied calculus".
- Introduce $C(x)$: " x has studied calculus."

If the domain for x consists of the students in the class, we can translate our statement as $\forall x C(x)$.

If we change the domain to consist of all people, we will need to express our statement as

- "For every person x , if person x is a student in this class then x has studied calculus."

If $S(x)$ represents the statement that person x is in this class, we see that our statement can be expressed as $\forall x (S(x) \rightarrow C(x))$.

What does $\forall x (S(x) \wedge C(x))$ represent?

Using two or more variables

When we are interested in the background of people in subjects besides calculus, we may prefer to use the two-variable quantifier $Q(x, y)$ for the statement "student x has studied subject y ." Then we would replace $C(x)$ by $Q(x, \text{calculus})$ in both approaches to obtain $\forall x Q(x, \text{calculus})$ or $\forall x (S(x) \rightarrow Q(x, \text{calculus}))$.

2. Express the statements "Some student in this class has visited Delhi" using predicates and quantifiers when

2. Express the statements "Some student in this class has visited Delhi" using predicates and quantifiers when Domain : Students in the class

2. Express the statements "Some student in this class has visited Delhi" using predicates and quantifiers when Domain : Students in the class
Domain: When people other than students in the class
"There is a person x having the properties that x is a student in this class and x has visited Delhi."

3. "Every student in this class has visited either Guwahati or Delhi".

3. "Every student in this class has visited either Guwahati or Delhi".
Domain: Students in class

3. "Every student in this class has visited either Guwahati or Delhi".

Domain: Students in class

For every x in this class, x has the property that x has visited Guwahati or x has visited Delhi.

3. "Every student in this class has visited either Guwahati or Delhi".

Domain: Students in class

For every x in this class, x has the property that x has visited Guwahati or x has visited Delhi.

$G(x)$: x has visited Guwahati and $D(x)$: x has visited Delhi.

3. "Every student in this class has visited either Guwahati or Delhi".

Domain: Students in class

For every x in this class, x has the property that x has visited Guwahati or x has visited Delhi.

$G(x)$: x has visited Guwahati and $D(x)$: x has visited Delhi.

$\forall x (G(x) \vee D(x))$.

3. "Every student in this class has visited either Guwahati or Delhi".

Domain: Students in class

For every x in this class, x has the property that x has visited Guwahati or x has visited Delhi.

$G(x)$: x has visited Guwahati and $D(x)$: x has visited Delhi.

$\forall x (G(x) \vee D(x))$.

Domain: consists of all people.

3. "Every student in this class has visited either Guwahati or Delhi".

Domain: Students in class

For every x in this class, x has the property that x has visited Guwahati or x has visited Delhi.

$G(x)$: x has visited Guwahati and $D(x)$: x has visited Delhi.

$\forall x (G(x) \vee D(x))$.

Domain: consists of all people.

Our statement can be expressed as "For every person x , if x is a student in this class, then x has visited Guwahati or x has visited Delhi."

3. "Every student in this class has visited either Guwahati or Delhi".

Domain: Students in class

For every x in this class, x has the property that x has visited Guwahati or x has visited Delhi.

$G(x)$: x has visited Guwahati and $D(x)$: x has visited Delhi.

$\forall x (G(x) \vee D(x))$.

Domain: consists of all people.

Our statement can be expressed as "For every person x , if x is a student in this class, then x has visited Guwahati or x has visited Delhi."

$S(x)$: x is a student in this class

3. "Every student in this class has visited either Guwahati or Delhi".

Domain: Students in class

For every x in this class, x has the property that x has visited Guwahati or x has visited Delhi.

$G(x)$: x has visited Guwahati and $D(x)$: x has visited Delhi.

$\forall x (G(x) \vee D(x))$.

Domain: consists of all people.

Our statement can be expressed as "For every person x , if x is a student in this class, then x has visited Guwahati or x has visited Delhi."

$S(x)$: x is a student in this class

$\forall x (S(x) \rightarrow (G(x) \vee D(x)))$.

Note: We could use a two-place predicate $Q(x, y)$ to represent " x has visited y ." Useful when more number of places are used.

Quantifiers in system specifications

"Every mail message larger than one megabyte will be compressed"

Quantifiers in system specifications

"Every mail message larger than one megabyte will be compressed"

- Denote $S(m, y)$: Mail m is larger than y mega bytes and $C(m)$: Mail m will be compressed.

Quantifiers in system specifications

"Every mail message larger than one megabyte will be compressed"

- Denote $S(m, y)$: Mail m is larger than y mega bytes and $C(m)$: Mail m will be compressed.
- Domain of m is the collection of all mails and y is a positive real number.

Quantifiers in system specifications

"Every mail message larger than one megabyte will be compressed"

- Denote $S(m, y)$: Mail m is larger than y mega bytes and $C(m)$: Mail m will be compressed.
- Domain of m is the collection of all mails and y is a positive real number.
- $\forall m(S(m, 1) \rightarrow C(m))$

"If a user is active, at least one network link will be available."

"If a user is active, at least one network link will be available."

- Let $A(u)$ represent "User u is active" where u has the domain of all users.

"If a user is active, at least one network link will be available."

- Let $A(u)$ represent "User u is active" where u has the domain of all users.
- Let $S(n, x)$ denote "Network link n is in state x ," where n has the domain of all network links and x has the domain of all possible states for a network link.

"If a user is active, at least one network link will be available."

- Let $A(u)$ represent "User u is active" where u has the domain of all users.
- Let $S(n, x)$ denote "Network link n is in state x ," where n has the domain of all network links and x has the domain of all possible states for a network link.
- $\exists u A(u) \rightarrow \exists n S(n, \text{available})$.

Example from Lewis Carroll

Consider these statements. The first two are called premises and the third is called the conclusion. The entire set is called an argument.

Example from Lewis Carroll

Consider these statements. The first two are called premises and the third is called the conclusion. The entire set is called an argument.

- All lions are fierce.

Example from Lewis Carroll

Consider these statements. The first two are called premises and the third is called the conclusion. The entire set is called an argument.

- All lions are fierce.
- Some lions do not drink coffee.

Example from Lewis Carroll

Consider these statements. The first two are called premises and the third is called the conclusion. The entire set is called an argument.

- All lions are fierce.
- Some lions do not drink coffee.
- Some fierce creatures do not drink coffee.

Example from Lewis Carroll

Consider these statements. The first two are called premises and the third is called the conclusion. The entire set is called an argument.

- All lions are fierce.
- Some lions do not drink coffee.
- Some fierce creatures do not drink coffee.

Let the domain consists of all creatures. Express each statement using quantifiers and predicates.

Example from Lewis Carroll

Consider these statements. The first two are called premises and the third is called the conclusion. The entire set is called an argument.

- All lions are fierce.
- Some lions do not drink coffee.
- Some fierce creatures do not drink coffee.

Let the domain consists of all creatures. Express each statement using quantifiers and predicates.

Let $P(x)$: x is a lion,

Example from Lewis Carroll

Consider these statements. The first two are called premises and the third is called the conclusion. The entire set is called an argument.

- All lions are fierce.
- Some lions do not drink coffee.
- Some fierce creatures do not drink coffee.

Let the domain consists of all creatures. Express each statement using quantifiers and predicates.

Let $P(x)$: x is a lion, $Q(x)$: x is fierce

Example from Lewis Carroll

Consider these statements. The first two are called premises and the third is called the conclusion. The entire set is called an argument.

- All lions are fierce.
- Some lions do not drink coffee.
- Some fierce creatures do not drink coffee.

Let the domain consists of all creatures. Express each statement using quantifiers and predicates.

Let $P(x)$: x is a lion, $Q(x)$: x is fierce and $R(x)$: x drinks coffee, respectively.

- $\forall x (P(x) \rightarrow Q(x))$.

Example from Lewis Carroll

Consider these statements. The first two are called premises and the third is called the conclusion. The entire set is called an argument.

- All lions are fierce.
- Some lions do not drink coffee.
- Some fierce creatures do not drink coffee.

Let the domain consists of all creatures. Express each statement using quantifiers and predicates.

Let $P(x)$: x is a lion, $Q(x)$: x is fierce and $R(x)$: x drinks coffee, respectively.

- $\forall x (P(x) \rightarrow Q(x))$.
- $\exists x (P(x) \wedge \neg R(x))$.

Example from Lewis Carroll

Consider these statements. The first two are called premises and the third is called the conclusion. The entire set is called an argument.

- All lions are fierce.
- Some lions do not drink coffee.
- Some fierce creatures do not drink coffee.

Let the domain consists of all creatures. Express each statement using quantifiers and predicates.

Let $P(x)$: x is a lion, $Q(x)$: x is fierce and $R(x)$: x drinks coffee, respectively.

- $\forall x (P(x) \rightarrow Q(x))$.
- $\exists x (P(x) \wedge \neg R(x))$.
- $\exists (Q(x) \wedge \neg R(x))$.

Example from Lewis Carroll

Consider these statements. The first two are called premises and the third is called the conclusion. The entire set is called an argument.

- All lions are fierce.
- Some lions do not drink coffee.
- Some fierce creatures do not drink coffee.

Let the domain consists of all creatures. Express each statement using quantifiers and predicates.

Let $P(x)$: x is a lion, $Q(x)$: x is fierce and $R(x)$: x drinks coffee, respectively.

- $\forall x (P(x) \rightarrow Q(x))$.
- $\exists x (P(x) \wedge \neg R(x))$.
- $\exists (Q(x) \wedge \neg R(x))$.

Compare the statements $\exists x (P(x) \wedge \neg R(x))$ and $\exists x (P(x) \rightarrow \neg R(x))$. (If there exists any creature (not necessarily lion) that does not drink coffee)

Consider these statements

- All hummingbirds are richly colored.

Consider these statements

- All hummingbirds are richly colored.
- No large birds live on honey.

Consider these statements

- All hummingbirds are richly colored.
- No large birds live on honey.
- Birds that do not live on honey are dull in color.

Consider these statements

- All hummingbirds are richly colored.
- No large birds live on honey.
- Birds that do not live on honey are dull in color.
- Hummingbirds are small.

Consider these statements

- All hummingbirds are richly colored.
- No large birds live on honey.
- Birds that do not live on honey are dull in color.
- Hummingbirds are small.

Assuming that the domain consists of all birds, express each of them using quantifiers and predicates.

Consider these statements

- All hummingbirds are richly colored.
- No large birds live on honey.
- Birds that do not live on honey are dull in color.
- Hummingbirds are small.

Assuming that the domain consists of all birds, express each of them using quantifiers and predicates.

Let $P(x)$: x is a hummingbird, $Q(x)$: x is large, $R(x)$: x lives on honey, and $S(x)$: x is richly colored, respectively.

Consider these statements

- All hummingbirds are richly colored.
- No large birds live on honey.
- Birds that do not live on honey are dull in color.
- Hummingbirds are small.

Assuming that the domain consists of all birds, express each of them using quantifiers and predicates.

Let $P(x)$: x is a hummingbird, $Q(x)$: x is large, $R(x)$: x lives on honey, and $S(x)$: x is richly colored, respectively.

- $\forall x(P(x) \rightarrow R(x))$

Consider these statements

- All hummingbirds are richly colored.
- No large birds live on honey.
- Birds that do not live on honey are dull in color.
- Hummingbirds are small.

Assuming that the domain consists of all birds, express each of them using quantifiers and predicates.

Let $P(x)$: x is a hummingbird, $Q(x)$: x is large, $R(x)$: x lives on honey, and $S(x)$: x is richly colored, respectively.

- $\forall x(P(x) \rightarrow R(x))$
- $\forall x(Q(x) \rightarrow \neg R(x))$ or $\neg \exists x(Q(x) \wedge R(x))$

Consider these statements

- All hummingbirds are richly colored.
- No large birds live on honey.
- Birds that do not live on honey are dull in color.
- Hummingbirds are small.

Assuming that the domain consists of all birds, express each of them using quantifiers and predicates.

Let $P(x)$: x is a hummingbird, $Q(x)$: x is large, $R(x)$: x lives on honey, and $S(x)$: x is richly colored, respectively.

- $\forall x(P(x) \rightarrow R(x))$
- $\forall x(Q(x) \rightarrow \neg R(x))$ or $\neg \exists x(Q(x) \wedge R(x))$
- $\forall x(\neg R(x) \rightarrow \neg S(x))$

Consider these statements

- All hummingbirds are richly colored.
- No large birds live on honey.
- Birds that do not live on honey are dull in color.
- Hummingbirds are small.

Assuming that the domain consists of all birds, express each of them using quantifiers and predicates.

Let $P(x)$: x is a hummingbird, $Q(x)$: x is large, $R(x)$: x lives on honey, and $S(x)$: x is richly colored, respectively.

- $\forall x(P(x) \rightarrow R(x))$
- $\forall x(Q(x) \rightarrow \neg R(x))$ or $\neg \exists x(Q(x) \wedge R(x))$
- $\forall x(\neg R(x) \rightarrow \neg S(x))$
- $\forall x(P(x) \rightarrow \neg Q(x)).$

Consider these statements

- All hummingbirds are richly colored.
- No large birds live on honey.
- Birds that do not live on honey are dull in color.
- Hummingbirds are small.

Assuming that the domain consists of all birds, express each of them using quantifiers and predicates.

Let $P(x)$: x is a hummingbird, $Q(x)$: x is large, $R(x)$: x lives on honey, and $S(x)$: x is richly colored, respectively.

- $\forall x(P(x) \rightarrow R(x))$
- $\forall x(Q(x) \rightarrow \neg R(x))$ or $\neg \exists x(Q(x) \wedge R(x))$
- $\forall x(\neg R(x) \rightarrow \neg S(x))$
- $\forall x(P(x) \rightarrow \neg Q(x)).$

Understanding Statements Involving Nested Quantifiers

Assume that the domain for the variables x and y consists of all real numbers.

Understanding Statements Involving Nested Quantifiers

Assume that the domain for the variables x and y consists of all real numbers.

- $\forall x \forall y (x + y = y + x)$

Understanding Statements Involving Nested Quantifiers

Assume that the domain for the variables x and y consists of all real numbers.

- $\forall x \forall y (x + y = y + x)$
 $x + y = y + x$ for all real numbers x and y . (Commutative law for addition of real numbers).

Understanding Statements Involving Nested Quantifiers

Assume that the domain for the variables x and y consists of all real numbers.

- $\forall x \forall y (x + y = y + x)$
 $x + y = y + x$ for all real numbers x and y . (Commutative law for addition of real numbers).
- $\forall x \exists y (x + y = 0)$

Understanding Statements Involving Nested Quantifiers

Assume that the domain for the variables x and y consists of all real numbers.

- $\forall x \forall y (x + y = y + x)$
 $x + y = y + x$ for all real numbers x and y . (Commutative law for addition of real numbers).
- $\forall x \exists y (x + y = 0)$
For every real number x there is a real number y such that $x + y = 0$. (Existence of additive inverse of x)

Understanding Statements Involving Nested Quantifiers

Assume that the domain for the variables x and y consists of all real numbers.

- $\forall x \forall y (x + y = y + x)$
 $x + y = y + x$ for all real numbers x and y . (Commutative law for addition of real numbers).
- $\forall x \exists y (x + y = 0)$
For every real number x there is a real number y such that $x + y = 0$. (Existence of additive inverse of x)
- $\forall x \forall y \forall z (x + (y + z) = (x + y) + z)$ is the associative law for addition of real numbers

Translate into English the statement $\forall x \forall y ((x > 0) \wedge (y < 0) \rightarrow (xy < 0))$, where the domain for both variables consists of all real numbers.

Translate into English the statement $\forall x \forall y ((x > 0) \wedge (y < 0) \rightarrow (xy < 0))$, where the domain for both variables consists of all real numbers.

Solution: This statement says that for every real number x and for every real number y , if $x > 0$ and $y < 0$, then $xy < 0$.

Translate into English the statement $\forall x \forall y ((x > 0) \wedge (y < 0) \rightarrow (xy < 0))$, where the domain for both variables consists of all real numbers.

Solution: This statement says that for every real number x and for every real number y , if $x > 0$ and $y < 0$, then $xy < 0$. That is, this statement says that for real numbers x and y , if x is positive and y is negative, then xy is negative.

Translate into English the statement $\forall x \forall y ((x > 0) \wedge (y < 0) \rightarrow (xy < 0))$, where the domain for both variables consists of all real numbers.

Solution: This statement says that for every real number x and for every real number y , if $x > 0$ and $y < 0$, then $xy < 0$. That is, this statement says that for real numbers x and y , if x is positive and y is negative, then xy is negative. This can be stated more clearly as "The product of a positive real number and a negative real number is always a negative real number."

Quantification as loops

In working with quantifications of more than one variable, it is sometimes helpful to think in terms of nested loops.

Quantification as loops

In working with quantifications of more than one variable, it is sometimes helpful to think in terms of nested loops.

To see whether $\forall x \forall y P(x, y)$ is true, we loop through the values for x , and for each x we loop through the values for y . If we find that $P(x, y)$ is true for all values for x and y , we have determined that $\forall x \forall y P(x, y)$ is true. If we ever hit a value x for which we hit a value y for which $P(x, y)$ is false, we have shown that $\forall x \forall y P(x, y)$ is false.

Similarly, to determine whether $\forall x \exists y P(x, y)$ is true, we loop through the values for x . For each x we loop through the values for y until we find a y for which $P(x, y)$ is true. If for every x we hit such a y , then $\forall x \exists y P(x, y)$ is true; if for some x we never hit such a y , then $\forall x \exists y P(x, y)$ is false.

To see whether $\exists x \forall y P(x, y)$ is true, we loop through the values for x until we find an x for which $P(x, y)$ is always true when we loop through all values for y . Once we find such an x , we know that $\exists x \forall y P(x, y)$ is true. If we never hit such an x , then we know that $\exists x \forall y P(x, y)$ is false.

To see whether $\exists x \forall y P(x, y)$ is true, we loop through the values for x until we find an x for which $P(x, y)$ is always true when we loop through all values for y . Once we find such an x , we know that $\exists x \forall y P(x, y)$ is true. If we never hit such an x , then we know that $\exists x \forall y P(x, y)$ is false.

Finally, to see whether $\exists x \exists y P(x, y)$ is true, we loop through the values for x , where for each x we loop through the values for y until we hit an x for which we hit a y for which $P(x, y)$ is true. The statement $\exists x \exists y P(x, y)$ is false only if we never hit an x for which we hit a y such that $P(x, y)$ is true.

Order of Quantifiers

1. Let $P(x, y)$ be the statement $x + y = y + x$.
 $\forall x \forall y P(x, y)$ denotes the proposition "For all real numbers x , for all real numbers y , $x + y = y + x$."

Order of Quantifiers

1. Let $P(x, y)$ be the statement $x + y = y + x$.
 $\forall x \forall y P(x, y)$ denotes the proposition "For all real numbers x , for all real numbers y , $x + y = y + x$."
Is it same as $\forall y \forall x P(x, y)$?

Order of Quantifiers

1. Let $P(x, y)$ be the statement $x + y = y + x$.
 $\forall x \forall y P(x, y)$ denotes the proposition "For all real numbers x , for all real numbers y , $x + y = y + x$."
Is it same as $\forall y \forall x P(x, y)$?
2. Let $Q(x, y)$ denote $x + y = 0$. What are the truth values of the quantifications $\exists x \forall y Q(x, y)$ and $\forall y \exists x Q(x, y)$ where the domain for all variables consists of all real numbers?

1. Let $Q(x, y, z)$ be the statement $x + y = z$. What are the truth values of the statements $\forall x \forall y \exists z Q(x, y, z)$ and $\exists z \forall x \forall y Q(x, y, z)$, where the domain of all variables consists of all real numbers?

1. Let $Q(x, y, z)$ be the statement $x + y = z$. What are the truth values of the statements $\forall x \forall y \exists z Q(x, y, z)$ and $\exists z \forall x \forall y Q(x, y, z)$, where the domain of all variables consists of all real numbers?
2. Translate the statement "The sum of two positive integers is always positive" into a logical expression.

1. Let $Q(x, y, z)$ be the statement $x + y = z$. What are the truth values of the statements $\forall x \forall y \exists z Q(x, y, z)$ and $\exists z \forall x \forall y Q(x, y, z)$, where the domain of all variables consists of all real numbers?
2. Translate the statement "The sum of two positive integers is always positive" into a logical expression.
3. Translate the statement "Every real number except zero has a multiplicative inverse."

1. Let $Q(x, y, z)$ be the statement $x + y = z$. What are the truth values of the statements $\forall x \forall y \exists z Q(x, y, z)$ and $\exists z \forall x \forall y Q(x, y, z)$, where the domain of all variables consists of all real numbers?
2. Translate the statement "The sum of two positive integers is always positive" into a logical expression.
3. Translate the statement "Every real number except zero has a multiplicative inverse."
4. Use quantifiers to express the definition of the limit of a real-valued function $f(x)$ of a real variable x at a point a in its domain.

1. Let $Q(x, y, z)$ be the statement $x + y = z$. What are the truth values of the statements $\forall x \forall y \exists z Q(x, y, z)$ and $\exists z \forall x \forall y Q(x, y, z)$, where the domain of all variables consists of all real numbers?
2. Translate the statement "The sum of two positive integers is always positive" into a logical expression.
3. Translate the statement "Every real number except zero has a multiplicative inverse."
4. Use quantifiers to express the definition of the limit of a real-valued function $f(x)$ of a real variable x at a point a in its domain.
For every real number $\epsilon > 0$ there exists a real number $\delta > 0$ such that $|f(x) - L| < \epsilon$ whenever $0 < |x - a| < \delta$.

- Translate the statement $\forall x(C(x) \vee \exists y(C(y) \wedge F(x, y)))$ into English, where $C(x)$ is "x has a computer," $F(x, y)$ is "x and y are friends," and the domain for both x and y consists of all students in your school.

- Translate the statement $\forall x(C(x) \vee \exists y(C(y) \wedge F(x, y)))$ into English, where $C(x)$ is "x has a computer," $F(x, y)$ is "x and y are friends," and the domain for both x and y consists of all students in your school.

The statement says that for every student x in your school, x has a computer or there is a student y such that y has a computer and x and y are friends.

- Translate the statement $\forall x(C(x) \vee \exists y(C(y) \wedge F(x, y)))$ into English, where $C(x)$ is "x has a computer," $F(x, y)$ is "x and y are friends," and the domain for both x and y consists of all students in your school.

The statement says that for every student x in your school, x has a computer or there is a student y such that y has a computer and x and y are friends. In other words, every student in your school has a computer or has a friend who has a computer.

Translate the statement $\exists x \forall y \forall z ((F(x, y) \wedge F(x, z) \wedge (y \neq z)) \rightarrow \neg F(y, z))$ into English, where $F(a, b)$ means a and b are friends and the domain for x, y , and z consists of all students in your school.

Translate the statement $\exists x \forall y \forall z ((F(x, y) \wedge F(x, z) \wedge (y \neq z)) \rightarrow \neg F(y, z))$ into English, where $F(a, b)$ means a and b are friends and the domain for x, y , and z consists of all students in your school.

- Examine the expression $(F(x, y) \wedge F(x, z) \wedge (y \neq z)) \rightarrow \neg F(y, z)$.

Translate the statement $\exists x \forall y \forall z ((F(x, y) \wedge F(x, z) \wedge (y \neq z)) \rightarrow \neg F(y, z))$ into English, where $F(a, b)$ means a and b are friends and the domain for x, y , and z consists of all students in your school.

- Examine the expression $(F(x, y) \wedge F(x, z) \wedge (y \neq z)) \rightarrow \neg F(y, z)$.
- This expression says that if students x and y are friends, and students x and z are friends, and furthermore, if y and z are not the same student, then y and z are not friends.

Translate the statement $\exists x \forall y \forall z ((F(x, y) \wedge F(x, z) \wedge (y \neq z)) \rightarrow \neg F(y, z))$ into English, where $F(a, b)$ means a and b are friends and the domain for x, y , and z consists of all students in your school.

- Examine the expression $(F(x, y) \wedge F(x, z) \wedge (y \neq z)) \rightarrow \neg F(y, z)$.
- This expression says that if students x and y are friends, and students x and z are friends, and furthermore, if y and z are not the same student, then y and z are not friends.
- It follows that the original statement, says that there is a student x such that for all students y and all students z other than y , if x and y are friends and x and z are friends, then y and z are not friends.

Translate the statement $\exists x \forall y \forall z ((F(x, y) \wedge F(x, z) \wedge (y \neq z)) \rightarrow \neg F(y, z))$ into English, where $F(a, b)$ means a and b are friends and the domain for x, y , and z consists of all students in your school.

- Examine the expression $(F(x, y) \wedge F(x, z) \wedge (y \neq z)) \rightarrow \neg F(y, z)$.
- This expression says that if students x and y are friends, and students x and z are friends, and furthermore, if y and z are not the same student, then y and z are not friends.
- It follows that the original statement, says that there is a student x such that for all students y and all students z other than y , if x and y are friends and x and z are friends, then y and z are not friends.
- In other words, there is a student none of whose friends are also friends with each other.

Express the statement "If a person is female and is a parent, then this person is someones mother" as a logical expression involving predicates, quantifiers with a domain consisting of all people, and logical connectives.

Express the statement "If a person is female and is a parent, then this person is someones mother" as a logical expression involving predicates, quantifiers with a domain consisting of all people, and logical connectives. $F(x)$: represent "x is female," $P(x)$: represent "x is a parent," and $M(x, y)$ to represent "x is the mother of y." The original statement can be represented as

Express the statement "If a person is female and is a parent, then this person is someones mother" as a logical expression involving predicates, quantifiers with a domain consisting of all people, and logical connectives. $F(x)$: represent "x is female," $P(x)$: represent "x is a parent," and $M(x, y)$ to represent "x is the mother of y." The original statement can be represented as

$$\forall x((F(x) \wedge P(x)) \rightarrow \exists y M(x, y)) \equiv \forall x \exists y((F(x) \wedge P(x)) \rightarrow M(x, y))$$

Express the statement "Everyone has exactly one best friend" as a logical expression involving predicates, quantifiers with a domain consisting of all people, and logical connectives.

$\forall x(\text{person } x \text{ has exactly one best friend})$, where the domain consists of all people.

Express the statement "Everyone has exactly one best friend" as a logical expression involving predicates, quantifiers with a domain consisting of all people, and logical connectives.

$\forall x(\text{person } x \text{ has exactly one best friend})$, where the domain consists of all people.

To say that x has exactly one best friend means that there is a person y who is the best friend of x , and furthermore, that for every person z , if person z is not person y , then z is not the best friend of x .

Express the statement "Everyone has exactly one best friend" as a logical expression involving predicates, quantifiers with a domain consisting of all people, and logical connectives.

$\forall x(\text{person } x \text{ has exactly one best friend})$, where the domain consists of all people.

To say that x has exactly one best friend means that there is a person y who is the best friend of x , and furthermore, that for every person z , if person z is not person y , then z is not the best friend of x . When we introduce the predicate $B(x, y)$ to be the statement " y is the best friend of x ," the statement that x has exactly one best friend can be represented as

Express the statement "Everyone has exactly one best friend" as a logical expression involving predicates, quantifiers with a domain consisting of all people, and logical connectives.

$\forall x(\text{person } x \text{ has exactly one best friend})$, where the domain consists of all people.

To say that x has exactly one best friend means that there is a person y who is the best friend of x , and furthermore, that for every person z , if person z is not person y , then z is not the best friend of x . When we introduce the predicate $B(x, y)$ to be the statement " y is the best friend of x ," the statement that x has exactly one best friend can be represented as $\forall x \exists y (B(x, y) \wedge \forall z ((z \neq y) \rightarrow \neg B(x, z)))$.

Use quantifiers and predicates to express the fact that $\lim_{x \rightarrow a} f(x)$ does not exist where $f(x)$ is a real-valued function of a real variable x and a belongs to the domain of f .

Use quantifiers and predicates to express the fact that $\lim_{x \rightarrow a} f(x)$ does not exist where $f(x)$ is a real-valued function of a real variable x and a belongs to the domain of f .

- $\lim_{x \rightarrow a} f(x)$ does not exist means that for all real numbers L ,
 $\lim_{x \rightarrow a} f(x) \neq L$.

Use quantifiers and predicates to express the fact that $\lim_{x \rightarrow a} f(x)$ does not exist where $f(x)$ is a real-valued function of a real variable x and a belongs to the domain of f .

- $\lim_{x \rightarrow a} f(x)$ does not exist means that for all real numbers L , $\lim_{x \rightarrow a} f(x) \neq L$.
- $\lim_{x \rightarrow a} f(x) \neq L$ can be expressed as $\neg(\forall \epsilon > 0 \exists \delta > 0 \forall x (0 < |x - a| < \delta \rightarrow |f(x) - L| < \epsilon))$.

Use quantifiers and predicates to express the fact that $\lim_{x \rightarrow a} f(x)$ does not exist where $f(x)$ is a real-valued function of a real variable x and a belongs to the domain of f .

- $\lim_{x \rightarrow a} f(x)$ does not exist means that for all real numbers L , $\lim_{x \rightarrow a} f(x) \neq L$.
- $\lim_{x \rightarrow a} f(x) \neq L$ can be expressed as $\neg(\forall \epsilon > 0 \exists \delta > 0 \forall x (0 < |x - a| < \delta \rightarrow |f(x) - L| < \epsilon))$.
- $\exists \epsilon > 0 \forall \delta > 0 \exists x (0 < |x - a| < \delta \wedge |f(x) - L| \geq \epsilon)$.

Use quantifiers and predicates to express the fact that $\lim_{x \rightarrow a} f(x)$ does not exist where $f(x)$ is a real-valued function of a real variable x and a belongs to the domain of f .

- $\lim_{x \rightarrow a} f(x)$ does not exist means that for all real numbers L , $\lim_{x \rightarrow a} f(x) \neq L$.
- $\lim_{x \rightarrow a} f(x) \neq L$ can be expressed as $\neg(\forall \epsilon > 0 \exists \delta > 0 \forall x(0 < |x - a| < \delta \rightarrow |f(x) - L| < \epsilon))$.
- $\exists \epsilon > 0 \forall \delta > 0 \exists x(0 < |x - a| < \delta \wedge |f(x) - L| \geq \epsilon)$.
- $\forall L \exists \epsilon > 0 \forall \delta > 0 \exists x(0 < |x - a| < \delta \wedge |f(x) - L| \geq \epsilon)$.

Use quantifiers and predicates to express the fact that $\lim_{x \rightarrow a} f(x)$ does not exist where $f(x)$ is a real-valued function of a real variable x and a belongs to the domain of f .

- $\lim_{x \rightarrow a} f(x)$ does not exist means that for all real numbers L , $\lim_{x \rightarrow a} f(x) \neq L$.
- $\lim_{x \rightarrow a} f(x) \neq L$ can be expressed as $\neg(\forall \epsilon > 0 \exists \delta > 0 \forall x(0 < |x - a| < \delta \rightarrow |f(x) - L| < \epsilon))$.
- $\exists \epsilon > 0 \forall \delta > 0 \exists x(0 < |x - a| < \delta \wedge |f(x) - L| \geq \epsilon)$.
- $\forall L \exists \epsilon > 0 \forall \delta > 0 \exists x(0 < |x - a| < \delta \wedge |f(x) - L| \geq \epsilon)$.

This last statement says that for every real number L there is a real number $\epsilon > 0$ such that for every real number $\delta > 0$, there exists a real number x such that $0 < |x - a| < \delta$ and $|f(x) - L| \geq \epsilon$.

An argument in propositional logic is a sequence of propositions. All but the final proposition in the argument are called premises and the final proposition is called the conclusion.

An argument in propositional logic is a sequence of propositions. All but the final proposition in the argument are called premises and the final proposition is called the conclusion. An argument is valid if the truth of all its premises implies that the conclusion is true.

An argument in propositional logic is a sequence of propositions. All but the final proposition in the argument are called premises and the final proposition is called the conclusion. An argument is valid if the truth of all its premises implies that the conclusion is true.

Examples: 1. Suppose that the conditional statement "If it snows today, then we will go skiing" and its hypothesis, "It is snowing today," are true.

An argument in propositional logic is a sequence of propositions. All but the final proposition in the argument are called premises and the final proposition is called the conclusion. An argument is valid if the truth of all its premises implies that the conclusion is true.

Examples: 1. Suppose that the conditional statement "If it snows today, then we will go skiing" and its hypothesis, "It is snowing today," are true. Then, it follows that the conclusion of the conditional statement, "We will go skiing," is true.

An argument in propositional logic is a sequence of propositions. All but the final proposition in the argument are called premises and the final proposition is called the conclusion. An argument is valid if the truth of all its premises implies that the conclusion is true.

Examples: 1. Suppose that the conditional statement "If it snows today, then we will go skiing" and its hypothesis, "It is snowing today," are true. Then, it follows that the conclusion of the conditional statement, "We will go skiing," is true.

2. Determine whether the argument given here is valid and determine whether its conclusion must be true because of the validity of the argument.

An argument in propositional logic is a sequence of propositions. All but the final proposition in the argument are called premises and the final proposition is called the conclusion. An argument is valid if the truth of all its premises implies that the conclusion is true.

Examples: 1. Suppose that the conditional statement "If it snows today, then we will go skiing" and its hypothesis, "It is snowing today," are true. Then, it follows that the conclusion of the conditional statement, "We will go skiing," is true.

2. Determine whether the argument given here is valid and determine whether its conclusion must be true because of the validity of the argument.

If $\sqrt{2} > \frac{3}{2}$, then $2 > (\frac{3}{2})^2$.

Table of rules of inference

p	$p \rightarrow q$	$\therefore q$
$\neg q$	$p \rightarrow q$	$\therefore \neg p$
$p \rightarrow q$	$q \rightarrow r$	$\therefore p \rightarrow r$
$p \vee q$	$\neg p$	$\therefore q$
p	$\therefore p \vee q$	
$p \wedge q$	$\therefore p$	
p	q	$\therefore p \wedge q$
$p \vee q$	$\neg p \vee r$	$\therefore q \vee r$

Table of rules of inference

p	$p \rightarrow q$	$\therefore q$
$\neg q$	$p \rightarrow q$	$\therefore \neg p$
$p \rightarrow q$	$q \rightarrow r$	$\therefore p \rightarrow r$
$p \vee q$	$\neg p$	$\therefore q$
p	$\therefore p \vee q$	
$p \wedge q$	$\therefore p$	
p	q	$\therefore p \wedge q$
$p \vee q$	$\neg p \vee r$	$\therefore q \vee r$

1. Show that the premises "It is not sunny this afternoon and it is colder than yesterday," "We will go swimming only if it is sunny," "If we do not go swimming, then we will take a boat trip," and "If we take a boat trip, then we will be home by sunset" lead to the conclusion "We will be home by sunset."

2. Show that the premises "If you send me an e-mail message, then I will finish writing the program," "If you do not send me an e-mail message, then I will go to sleep early," and "If I go to sleep early, then I will wake up feeling refreshed" lead to the conclusion "If I do not finish writing the program, then I will wake up feeling refreshed."

2. Show that the premises "If you send me an e-mail message, then I will finish writing the program," "If you do not send me an e-mail message, then I will go to sleep early," and "If I go to sleep early, then I will wake up feeling refreshed" lead to the conclusion "If I do not finish writing the program, then I will wake up feeling refreshed."
3. Use the rules of inference to show that the hypotheses "Jasmine is skiing or it is not snowing" and "It is snowing or Bart is playing hockey" imply that "Jasmine is skiing or Bart is playing hockey."

2. Show that the premises "If you send me an e-mail message, then I will finish writing the program," "If you do not send me an e-mail message, then I will go to sleep early," and "If I go to sleep early, then I will wake up feeling refreshed" lead to the conclusion "If I do not finish writing the program, then I will wake up feeling refreshed."
3. Use the rules of inference to show that the hypotheses "Jasmine is skiing or it is not snowing" and "It is snowing or Bart is playing hockey" imply that "Jasmine is skiing or Bart is playing hockey."
4. Show that the premises $(p \wedge q) \vee r$ and $r \rightarrow s$ imply the conclusion $p \vee s$.

2. Show that the premises "If you send me an e-mail message, then I will finish writing the program," "If you do not send me an e-mail message, then I will go to sleep early," and "If I go to sleep early, then I will wake up feeling refreshed" lead to the conclusion "If I do not finish writing the program, then I will wake up feeling refreshed."

3. Use the rules of inference to show that the hypotheses "Jasmine is skiing or it is not snowing" and "It is snowing or Bart is playing hockey" imply that "Jasmine is skiing or Bart is playing hockey."

4. Show that the premises $(p \wedge q) \vee r$ and $r \rightarrow s$ imply the conclusion $p \vee s$.

We can rewrite the premises $(p \wedge q) \vee r$ as two clauses (clause is a disjunction of variables or negations of these variables), $p \vee r$ and $q \vee r$.

We can also replace $r \rightarrow s$ by the equivalent clause $\neg r \vee s$. Using the two clauses $p \vee r$ and $\neg r \vee s$, we can use resolution to conclude $p \vee s$.

Rules of Inference for Quantified Statements

- $\forall x P(x)$
 $\therefore P(c).$

Rules of Inference for Quantified Statements

- $\forall xP(x)$
 $\therefore P(c).$
- $P(c)$ for an arbitrary c
 $\therefore \forall xP(x).$

Rules of Inference for Quantified Statements

- $\forall x P(x)$
 $\therefore P(c).$
- $P(c)$ for an arbitrary c
 $\therefore \forall x P(x).$
- $\exists x P(x)$
 $\therefore P(c)$ for some element $c.$

Rules of Inference for Quantified Statements

- $\forall xP(x)$
 $\therefore P(c)$.
- $P(c)$ for an arbitrary c
 $\therefore \forall xP(x)$.
- $\exists xP(x)$
 $\therefore P(c)$ for some element c .
- $P(c)$ for some element c
 $\exists xP(x)$.

Rules of Inference for Quantified Statements

- $\forall x P(x)$
 $\therefore P(c).$
- $P(c)$ for an arbitrary c
 $\therefore \forall x P(x).$
- $\exists x P(x)$
 $\therefore P(c)$ for some element $c.$
- $P(c)$ for some element c
 $\exists x P(x).$
- If $\forall x (P(x) \rightarrow Q(x))$ is true, and if $P(a)$ is true for a particular element a in the domain of the universal quantifier, then $Q(a)$ must also be true