

Introduction to Logic

Department of Science and Mathematics

IIIT Guwahati.

August 21, 2020

Introduction to proofs

- A proof is a valid argument that establishes the truth of a mathematical statement.

Introduction to proofs

- A proof is a valid argument that establishes the truth of a mathematical statement. A proof can use the hypotheses of the theorem, if any, axioms assumed to be true, and previously proven theorems.

Introduction to proofs

- A proof is a valid argument that establishes the truth of a mathematical statement. A proof can use the hypotheses of the theorem, if any, axioms assumed to be true, and previously proven theorems. Using these ingredients and rules of inference, the final step of the proof establishes the truth of the statement being proved.

Introduction to proofs

- A proof is a valid argument that establishes the truth of a mathematical statement. A proof can use the hypotheses of the theorem, if any, axioms assumed to be true, and previously proven theorems. Using these ingredients and rules of inference, the final step of the proof establishes the truth of the statement being proved.
- A less important theorem that is helpful in the proof of other results is called a lemma (plural lemmas).

Introduction to proofs

- A proof is a valid argument that establishes the truth of a mathematical statement. A proof can use the hypotheses of the theorem, if any, axioms assumed to be true, and previously proven theorems. Using these ingredients and rules of inference, the final step of the proof establishes the truth of the statement being proved.
- A less important theorem that is helpful in the proof of other results is called a lemma (plural lemmas). Complicated proofs are usually easier to understand when they are proved using a series of lemmas, where each lemma is proved individually.

Introduction to proofs

- A proof is a valid argument that establishes the truth of a mathematical statement. A proof can use the hypotheses of the theorem, if any, axioms assumed to be true, and previously proven theorems. Using these ingredients and rules of inference, the final step of the proof establishes the truth of the statement being proved.
- A less important theorem that is helpful in the proof of other results is called a lemma (plural lemmas). Complicated proofs are usually easier to understand when they are proved using a series of lemmas, where each lemma is proved individually.
- A corollary is a theorem that can be established directly from a theorem that has been proved.

Introduction to proofs

- A proof is a valid argument that establishes the truth of a mathematical statement. A proof can use the hypotheses of the theorem, if any, axioms assumed to be true, and previously proven theorems. Using these ingredients and rules of inference, the final step of the proof establishes the truth of the statement being proved.
- A less important theorem that is helpful in the proof of other results is called a lemma (plural lemmas). Complicated proofs are usually easier to understand when they are proved using a series of lemmas, where each lemma is proved individually.
- A corollary is a theorem that can be established directly from a theorem that has been proved.
- A conjecture is a statement that is being proposed to be a true statement, usually on the basis of some partial evidence, a heuristic argument, or the intuition of an expert.

Introduction to proofs

- A proof is a valid argument that establishes the truth of a mathematical statement. A proof can use the hypotheses of the theorem, if any, axioms assumed to be true, and previously proven theorems. Using these ingredients and rules of inference, the final step of the proof establishes the truth of the statement being proved.
- A less important theorem that is helpful in the proof of other results is called a lemma (plural lemmas). Complicated proofs are usually easier to understand when they are proved using a series of lemmas, where each lemma is proved individually.
- A corollary is a theorem that can be established directly from a theorem that has been proved.
- A conjecture is a statement that is being proposed to be a true statement, usually on the basis of some partial evidence, a heuristic argument, or the intuition of an expert. When a proof of a conjecture is found, the conjecture becomes a theorem.

A direct proof of a conditional statement $p \rightarrow q$ is constructed when the first step is the assumption that p is true; subsequent steps are constructed using rules of inference, with the final step showing that q must also be true.

A direct proof of a conditional statement $p \rightarrow q$ is constructed when the first step is the assumption that p is true; subsequent steps are constructed using rules of inference, with the final step showing that q must also be true.

1. Give a direct proof of the theorem "If n is an odd integer, then n^2 is odd."

2. Give a direct proof that if m and n are both perfect squares, then nm is also a perfect square.

Proof by contraposition

Proof by contraposition make use of the fact that the conditional statement $p \rightarrow q$ is equivalent to its contrapositive, $\neg q \rightarrow \neg p$.

Proof by contraposition

Proof by contraposition make use of the fact that the conditional statement $p \rightarrow q$ is equivalent to its contrapositive, $\neg q \rightarrow \neg p$.

3. Prove that if n is an integer and $3n + 2$ is odd, then n is odd.

4. Prove that if $n = ab$, where a and b are positive integers, then $a \leq \sqrt{n}$ or $b \leq \sqrt{n}$.

6. Let $P(n)$ be "If a and b are positive integers with $a \geq b$, then $a^n \geq b^n$ ", where the domain consists of all nonnegative integers. Show that $P(0)$ is true. (Trivial Proof)

6. Let $P(n)$ be "If a and b are positive integers with $a \geq b$, then $a^n \geq b^n$ ", where the domain consists of all nonnegative integers. Show that $P(0)$ is true. (Trivial Proof)

Proof Strategies- Try the direct proof method, if it fails, then try with method of contraposition.

7. Prove that the sum of two rational numbers is rational.

6. Let $P(n)$ be "If a and b are positive integers with $a \geq b$, then $a^n \geq b^n$ ", where the domain consists of all nonnegative integers. Show that $P(0)$ is true. (Trivial Proof)

Proof Strategies- Try the direct proof method, if it fails, then try with method of contraposition.

7. Prove that the sum of two rational numbers is rational.

8. Prove that if n is an integer and n^2 is odd, then n is odd.

6. Let $P(n)$ be "If a and b are positive integers with $a \geq b$, then $a^n \geq b^n$ ", where the domain consists of all nonnegative integers. Show that $P(0)$ is true. (Trivial Proof)

Proof Strategies- Try the direct proof method, if it fails, then try with method of contraposition.

7. Prove that the sum of two rational numbers is rational.

8. Prove that if n is an integer and n^2 is odd, then n is odd.

Method of contradiction

10. Prove that $\sqrt{2}$ is irrational by giving a proof by contradiction.

9. Show that at least four of any 22 days must fall on the same day of the week.

9. Show that at least four of any 22 days must fall on the same day of the week.
11. Prove the theorem "If n is an integer, then n is odd if and only if n^2 is odd."

Proofs by equivalence

Sometimes a theorem states that several propositions are equivalent. Such a theorem states that propositions $p_1, p_2, p_3, \dots, p_n$ are equivalent. This can be written as $p_1 \leftrightarrow p_2 \leftrightarrow \dots \leftrightarrow p_n$
 $\equiv (p_1 \rightarrow p_2) \wedge (p_2 \rightarrow p_3) \wedge \dots (p_n \rightarrow p_1).$

Proofs by equivalence

Sometimes a theorem states that several propositions are equivalent. Such a theorem states that propositions $p_1, p_2, p_3, \dots, p_n$ are equivalent. This can be written as $p_1 \leftrightarrow p_2 \leftrightarrow \dots \leftrightarrow p_n$
 $\equiv (p_1 \rightarrow p_2) \wedge (p_2 \rightarrow p_3) \wedge \dots \wedge (p_n \rightarrow p_1).$

- Show that these statements about the integer n are equivalent:

p_1 : n is even.

p_2 : $n - 1$ is odd.

p_3 : n^2 is even.

Counter Examples

12. Show that the statement "Every positive integer is the sum of the squares of two integers" is false. (Give counter example)

Mistakes in Proofs

What is wrong with this famous supposed "proof" that $1 = 2$?

Mistakes in Proofs

What is wrong with this famous supposed "proof" that $1 = 2$?
"Proof": We use these steps, where a and b are two equal positive integers.

Mistakes in Proofs

What is wrong with this famous supposed "proof" that $1 = 2$?
"Proof": We use these steps, where a and b are two equal positive integers.

- Let $a = b$

Mistakes in Proofs

What is wrong with this famous supposed "proof" that $1 = 2$?
"Proof": We use these steps, where a and b are two equal positive integers.

- Let $a = b$
- $a^2 = ab$

Mistakes in Proofs

What is wrong with this famous supposed "proof" that $1 = 2$?
"Proof": We use these steps, where a and b are two equal positive integers.

- Let $a = b$
- $a^2 = ab$
- $a^2 - b^2 = ab - b^2$

Mistakes in Proofs

What is wrong with this famous supposed "proof" that $1 = 2$?
"Proof": We use these steps, where a and b are two equal positive integers.

- Let $a = b$
- $a^2 = ab$
- $a^2 - b^2 = ab - b^2$
- $(a - b)(a + b) = b(a - b)$

Mistakes in Proofs

What is wrong with this famous supposed "proof" that $1 = 2$?

"Proof": We use these steps, where a and b are two equal positive integers.

- Let $a = b$
- $a^2 = ab$
- $a^2 - b^2 = ab - b^2$
- $(a - b)(a + b) = b(a - b)$
- $a + b = b$

Mistakes in Proofs

What is wrong with this famous supposed "proof" that $1 = 2$?
"Proof": We use these steps, where a and b are two equal positive integers.

- Let $a = b$
- $a^2 = ab$
- $a^2 - b^2 = ab - b^2$
- $(a - b)(a + b) = b(a - b)$
- $a + b = b$
- $2b = b$

Mistakes in Proofs

What is wrong with this famous supposed "proof" that $1 = 2$?

"Proof": We use these steps, where a and b are two equal positive integers.

- Let $a = b$
- $a^2 = ab$
- $a^2 - b^2 = ab - b^2$
- $(a - b)(a + b) = b(a - b)$
- $a + b = b$
- $2b = b$
- $2 = 1$.

Mistakes in Proofs

What is wrong with this famous supposed "proof" that $1 = 2$?

"Proof": We use these steps, where a and b are two equal positive integers.

- Let $a = b$
- $a^2 = ab$
- $a^2 - b^2 = ab - b^2$
- $(a - b)(a + b) = b(a - b)$
- $a + b = b$
- $2b = b$
- $2 = 1$.

The error is that $a - b$ equals zero; division of both sides of an equation by the same quantity is valid as long as this quantity is not zero.

Circular reasoning

Is the following argument correct?

It supposedly shows that n is an even integer whenever n^2 is an even integer.

Circular reasoning

Is the following argument correct?

It supposedly shows that n is an even integer whenever n^2 is an even integer.

"Proof":

Suppose that n^2 is even. Then $n^2 = 2k$ for some integer k . Let $n = 2l$ for some integer l . This shows that n is even.

Exhaustive proof

Some theorems can be proved by examining a relatively small number of examples. Such proofs are called exhaustive proofs, or proofs by exhaustion.

Exhaustive proof

Some theorems can be proved by examining a relatively small number of examples. Such proofs are called exhaustive proofs, or proofs by exhaustion.

13. Prove that $(n + 1)^3 \geq 3n$ if n is a positive integer with $n \leq 4$.

Exhaustive proof

Some theorems can be proved by examining a relatively small number of examples. Such proofs are called exhaustive proofs, or proofs by exhaustion.

13. Prove that $(n + 1)^3 \geq 3n$ if n is a positive integer with $n \leq 4$.

Proof by Cases

14. Prove that if n is an integer, then $n^2 \geq n$.

Proof by Cases

14. Prove that if n is an integer, then $n^2 \geq n$.
15. Use a proof by cases to show that $|xy| = |x||y|$, where x and y are real numbers.

16. Formulate a conjecture about the final decimal digit of the square of an integer and prove your result.

Sometimes we can eliminate all but a few examples in a proof by cases.

Sometimes we can eliminate all but a few examples in a proof by cases.
17. Show that there are no solutions in integers x and y of $x^2 + 3y^2 = 8$.

WITHOUT LOSS OF GENERALITY

- In the proof in Example 15, we can dismiss the case where $x < 0$ and $y \geq 0$, because it is the same as case where $x \geq 0$ and $y < 0$, with the roles of x and y reversed.

WITHOUT LOSS OF GENERALITY

- In the proof in Example 15, we can dismiss the case where $x < 0$ and $y \geq 0$, because it is the same as case where $x \geq 0$ and $y < 0$, with the roles of x and y reversed.
- To shorten the proof, we could have proved cases together by assuming, without loss of generality, that $x \geq 0$ and $y < 0$.

Implicit in this statement is that we can complete the case with $x < 0$ and $y \geq 0$ using the same argument as we used for the case with $x \geq 0$ and $y < 0$, but with the obvious changes.

WITHOUT LOSS OF GENERALITY

- In the proof in Example 15, we can dismiss the case where $x < 0$ and $y \geq 0$, because it is the same as case where $x \geq 0$ and $y < 0$, with the roles of x and y reversed.
- To shorten the proof, we could have proved cases together by assuming, without loss of generality, that $x \geq 0$ and $y < 0$.

Implicit in this statement is that we can complete the case with $x < 0$ and $y \geq 0$ using the same argument as we used for the case with $x \geq 0$ and $y < 0$, but with the obvious changes.

We now illustrate a proof where without loss of generality is used effectively together with other proof techniques.

18. Show that if x and y are integers and both xy and $x + y$ are even, then both x and y are even.

18. Show that if x and y are integers and both xy and $x + y$ are even, then both x and y are even.

We will use proof by contraposition, the notion of without loss of generality, and proof by cases.

18. Show that if x and y are integers and both xy and $x + y$ are even, then both x and y are even.

We will use proof by contraposition, the notion of without loss of generality, and proof by cases.

- First, suppose that x and y are not both even. That is, assume that x is odd or that y is odd (or both).
- WLOG, we assume that x is odd, so that $x = 2m + 1$ for some integer k .

18. Show that if x and y are integers and both xy and $x + y$ are even, then both x and y are even.

We will use proof by contraposition, the notion of without loss of generality, and proof by cases.

- First, suppose that x and y are not both even. That is, assume that x is odd or that y is odd (or both).
- WLOG, we assume that x is odd, so that $x = 2m + 1$ for some integer k . To complete the proof, we need to show that xy is odd or $x + y$ is odd.
- Consider two cases: (i) y even, and (ii) y odd.

18. Show that if x and y are integers and both xy and $x + y$ are even, then both x and y are even.

We will use proof by contraposition, the notion of without loss of generality, and proof by cases.

- First, suppose that x and y are not both even. That is, assume that x is odd or that y is odd (or both).
- WLOG, we assume that x is odd, so that $x = 2m + 1$ for some integer k . To complete the proof, we need to show that xy is odd or $x + y$ is odd.
- Consider two cases: (i) y even, and (ii) y odd.
- In (i), $y = 2n$ for some integer n , so that

18. Show that if x and y are integers and both xy and $x + y$ are even, then both x and y are even.

We will use proof by contraposition, the notion of without loss of generality, and proof by cases.

- First, suppose that x and y are not both even. That is, assume that x is odd or that y is odd (or both).
- WLOG, we assume that x is odd, so that $x = 2m + 1$ for some integer k . To complete the proof, we need to show that xy is odd or $x + y$ is odd.
- Consider two cases: (i) y even, and (ii) y odd.
- In (i), $y = 2n$ for some integer n , so that $x + y = (2m + 1) + 2n = 2(m + n) + 1$ is odd.

18. Show that if x and y are integers and both xy and $x + y$ are even, then both x and y are even.

We will use proof by contraposition, the notion of without loss of generality, and proof by cases.

- First, suppose that x and y are not both even. That is, assume that x is odd or that y is odd (or both).
- WLOG, we assume that x is odd, so that $x = 2m + 1$ for some integer k . To complete the proof, we need to show that xy is odd or $x + y$ is odd.
- Consider two cases: (i) y even, and (ii) y odd.
- In (i), $y = 2n$ for some integer n , so that $x + y = (2m + 1) + 2n = 2(m + n) + 1$ is odd.
- In (ii), $y = 2n + 1$ for some integer n ,

18. Show that if x and y are integers and both xy and $x + y$ are even, then both x and y are even.

We will use proof by contraposition, the notion of without loss of generality, and proof by cases.

- First, suppose that x and y are not both even. That is, assume that x is odd or that y is odd (or both).
- WLOG, we assume that x is odd, so that $x = 2m + 1$ for some integer k . To complete the proof, we need to show that xy is odd or $x + y$ is odd.
- Consider two cases: (i) y even, and (ii) y odd.
- In (i), $y = 2n$ for some integer n , so that $x + y = (2m + 1) + 2n = 2(m + n) + 1$ is odd.
- In (ii), $y = 2n + 1$ for some integer n , so that $xy = (2m + 1)(2n + 1) = 4mn + 2m + 2n + 1 = 2(2mn + m + n) + 1$ is odd.

18. Show that if x and y are integers and both xy and $x + y$ are even, then both x and y are even.

We will use proof by contraposition, the notion of without loss of generality, and proof by cases.

- First, suppose that x and y are not both even. That is, assume that x is odd or that y is odd (or both).
- WLOG, we assume that x is odd, so that $x = 2m + 1$ for some integer k . To complete the proof, we need to show that xy is odd or $x + y$ is odd.
- Consider two cases: (i) y even, and (ii) y odd.
- In (i), $y = 2n$ for some integer n , so that $x + y = (2m + 1) + 2n = 2(m + n) + 1$ is odd.
- In (ii), $y = 2n + 1$ for some integer n , so that $xy = (2m + 1)(2n + 1) = 4mn + 2m + 2n + 1 = 2(2mn + m + n) + 1$ is odd. This completes the proof by contraposition.

19. What is wrong with this "proof?" If x is a real number, then x^2 is a positive real number.

Proof: " Let p_1 be " x is positive," let p_2 be " x is negative," and let q be " x^2 is positive." To show that $p_1 \rightarrow q$ is true, note that when x is positive, x^2 is positive because it is the product of two positive numbers, x and x . To show that $p_2 \rightarrow q$, note that when x is negative, x^2 is positive because it is the product of two negative numbers, x and x . This completes the proof.

A proof of a proposition of the form $\exists x P(x)$ is called an existence proof.

A proof of a proposition of the form $\exists xP(x)$ is called an existence proof. There are several ways to prove a theorem of this type. Sometimes an existence proof of $\exists xP(x)$ can be given by finding an element a , called a witness, such that $P(a)$ is true. This type of existence proof is called constructive.

A proof of a proposition of the form $\exists xP(x)$ is called an existence proof. There are several ways to prove a theorem of this type. Sometimes an existence proof of $\exists xP(x)$ can be given by finding an element a , called a witness, such that $P(a)$ is true. This type of existence proof is called constructive. It is also possible to give an existence proof that is nonconstructive; that is, we do not find an element a such that $P(a)$ is true, but rather prove that $\exists xP(x)$ is true in some other way.

A Constructive Existence Proof

20. Show that there is a positive integer that can be written as the sum of cubes of positive integers in two different ways.

A Constructive Existence Proof

20. Show that there is a positive integer that can be written as the sum of cubes of positive integers in two different ways.

$$1729 = 12^3 + 1 = 10^3 + 9^3.$$

A Nonconstructive Existence Proof

21. Show that there exist irrational numbers x and y such that x^y is rational.

A Nonconstructive Existence Proof

21. Show that there exist irrational numbers x and y such that x^y is rational.

We know that $\sqrt{2}$ is irrational.

A Nonconstructive Existence Proof

21. Show that there exist irrational numbers x and y such that x^y is rational.

We know that $\sqrt{2}$ is irrational. Consider the number $\sqrt{2}^{\sqrt{2}}$.

A Nonconstructive Existence Proof

21. Show that there exist irrational numbers x and y such that x^y is rational.

We know that $\sqrt{2}$ is irrational. Consider the number $\sqrt{2}^{\sqrt{2}}$. If it is rational, we have two irrational numbers x and y with x^y rational, namely, $x = \sqrt{2}$ and $y = \sqrt{2}$.

A Nonconstructive Existence Proof

21. Show that there exist irrational numbers x and y such that x^y is rational.

We know that $\sqrt{2}$ is irrational. Consider the number $\sqrt{2}^{\sqrt{2}}$. If it is rational, we have two irrational numbers x and y with x^y rational, namely, $x = \sqrt{2}$ and $y = \sqrt{2}$. On the other hand if $\sqrt{2}^{\sqrt{2}}$ is irrational, then we can let $x = \sqrt{2}^{\sqrt{2}}$ and $y = \sqrt{2}$.

22. Show that if a and b are real numbers and $a \neq 0$, then there is a unique real number r such that $ar + b = 0$.

Tiling

A checkerboard is a rectangle divided into squares of the same size by horizontal and vertical lines.

Tiling

A checkerboard is a rectangle divided into squares of the same size by horizontal and vertical lines. The game of checkers is played on a board with 8 rows and 8 columns; this board is called the standard checkerboard.

Tiling

A checkerboard is a rectangle divided into squares of the same size by horizontal and vertical lines. The game of checkers is played on a board with 8 rows and 8 columns; this board is called the standard checkerboard. In this section we use the term board to refer to a checkerboard of any rectangular size as well as parts of checkerboards obtained by removing one or more squares.

Tiling

A checkerboard is a rectangle divided into squares of the same size by horizontal and vertical lines. The game of checkers is played on a board with 8 rows and 8 columns; this board is called the standard checkerboard. In this section we use the term board to refer to a checkerboard of any rectangular size as well as parts of checkerboards obtained by removing one or more squares. A domino is a rectangular piece that is one square by two squares.

Tiling

A checkerboard is a rectangle divided into squares of the same size by horizontal and vertical lines. The game of checkers is played on a board with 8 rows and 8 columns; this board is called the standard checkerboard. In this section we use the term board to refer to a checkerboard of any rectangular size as well as parts of checkerboards obtained by removing one or more squares. A domino is a rectangular piece that is one square by two squares. We say that a board is tiled by dominoes when all its squares are covered with no overlapping dominoes and no dominoes overhanging the board.

1. Can we tile the standard checkerboard using dominoes?

Solution: We can find manyways to tile the standard checkerboard using dominoes.

1. Can we tile the standard checkerboard using dominoes?

Solution: We can find manyways to tile the standard checkerboard using dominoes. For example, we can tile it by placing 32 dominoes horizontally.

1. Can we tile the standard checkerboard using dominoes?

Solution: We can find manyways to tile the standard checkerboard using dominoes. For example, we can tile it by placing 32 dominoes horizontally.

2. Can we tile a board obtained by removing one of the four corner squares of a standard checkerboard?

1. Can we tile the standard checkerboard using dominoes?

Solution: We can find manyways to tile the standard checkerboard using dominoes. For example, we can tile it by placing 32 dominoes horizontally.

2. Can we tile a board obtained by removing one of the four corner squares of a standard checkerboard?

Solution: To answer this question, note that a standard checkerboard has 64 squares, so removing a square produces a board with 63 squares.

1. Can we tile the standard checkerboard using dominoes?

Solution: We can find manyways to tile the standard checkerboard using dominoes. For example, we can tile it by placing 32 dominoes horizontally.

2. Can we tile a board obtained by removing one of the four corner squares of a standard checkerboard?

Solution: To answer this question, note that a standard checkerboard has 64 squares, so removing a square produces a board with 63 squares. Now suppose that we could tile a board obtained from the standard checkerboard by removing a corner square.

1. Can we tile the standard checkerboard using dominoes?

Solution: We can find manyways to tile the standard checkerboard using dominoes. For example, we can tile it by placing 32 dominoes horizontally.

2. Can we tile a board obtained by removing one of the four corner squares of a standard checkerboard?

Solution: To answer this question, note that a standard checkerboard has 64 squares, so removing a square produces a board with 63 squares. Now suppose that we could tile a board obtained from the standard checkerboard by removing a corner square. The board has an even number of squares because each domino covers two squares and no two dominoes overlap and no dominoes overhang the board.

1. Can we tile the standard checkerboard using dominoes?

Solution: We can find manyways to tile the standard checkerboard using dominoes. For example, we can tile it by placing 32 dominoes horizontally.

2. Can we tile a board obtained by removing one of the four corner squares of a standard checkerboard?

Solution: To answer this question, note that a standard checkerboard has 64 squares, so removing a square produces a board with 63 squares. Now suppose that we could tile a board obtained from the standard checkerboard by removing a corner square. The board has an even number of squares because each domino covers two squares and no two dominoes overlap and no dominoes overhang the board. Consequently, we can prove by contradiction that a standard checkerboard with one square removed cannot be tiled using dominoes because such a board has an odd number of squares.

3. Can we tile the board obtained by deleting the upper left and lower right corner squares of a standard checkerboard?

3. Can we tile the board obtained by deleting the upper left and lower right corner squares of a standard checkerboard?
Method of Problem 2 fails here.

3. Can we tile the board obtained by deleting the upper left and lower right corner squares of a standard checkerboard?

Method of Problem 2 fails here.

We might try to prove that no tiling exists by showing that we reach a dead end however we successively place dominoes on the board.

3. Can we tile the board obtained by deleting the upper left and lower right corner squares of a standard checkerboard?

Method of Problem 2 fails here.

We might try to prove that no tiling exists by showing that we reach a dead end however we successively place dominoes on the board. To construct such a proof, we would have to consider all possible cases that arise as we run through all possible choices of successively placing dominoes.

3. Can we tile the board obtained by deleting the upper left and lower right corner squares of a standard checkerboard?

Method of Problem 2 fails here.

We might try to prove that no tiling exists by showing that we reach a dead end however we successively place dominoes on the board. To construct such a proof, we would have to consider all possible cases that arise as we run through all possible choices of successively placing dominoes. For example, we have two choices for covering the square in the second column of the first row, next to the removed top left corner. We could cover it with a horizontally placed tile or a vertically placed tile. Each of these two choices leads to further choices, and so on.

We color the squares of this checkerboard using alternating white and black squares.

3. Can we tile the board obtained by deleting the upper left and lower right corner squares of a standard checkerboard?

Method of Problem 2 fails here.

We might try to prove that no tiling exists by showing that we reach a dead end however we successively place dominoes on the board. To construct such a proof, we would have to consider all possible cases that arise as we run through all possible choices of successively placing dominoes. For example, we have two choices for covering the square in the second column of the first row, next to the removed top left corner. We could cover it with a horizontally placed tile or a vertically placed tile. Each of these two choices leads to further choices, and so on.

We color the squares of this checkerboard using alternating white and black squares. Observe that a domino in a tiling of such a board covers one white square and one black square.

3. Can we tile the board obtained by deleting the upper left and lower right corner squares of a standard checkerboard?

Method of Problem 2 fails here.

We might try to prove that no tiling exists by showing that we reach a dead end however we successively place dominoes on the board. To construct such a proof, we would have to consider all possible cases that arise as we run through all possible choices of successively placing dominoes. For example, we have two choices for covering the square in the second column of the first row, next to the removed top left corner. We could cover it with a horizontally placed tile or a vertically placed tile. Each of these two choices leads to further choices, and so on.

We color the squares of this checkerboard using alternating white and black squares. Observe that a domino in a tiling of such a board covers one white square and one black square. Next, note that this board has unequal numbers of white square and black squares.

Suppose we can use dominoes to tile a standard checkerboard with opposite corners removed.

Suppose we can use dominoes to tile a standard checkerboard with opposite corners removed. Note that the standard checkerboard with opposite corners removed contains $64 - 2 = 62$ squares.

Suppose we can use dominoes to tile a standard checkerboard with opposite corners removed. Note that the standard checkerboard with opposite corners removed contains $64 - 2 = 62$ squares. The tiling would use $62/2 = 31$ dominoes.

Suppose we can use dominoes to tile a standard checkerboard with opposite corners removed. Note that the standard checkerboard with opposite corners removed contains $64 - 2 = 62$ squares. The tiling would use $62/2 = 31$ dominoes. Note that each domino in this tiling covers one white and one black square.

Suppose we can use dominoes to tile a standard checkerboard with opposite corners removed. Note that the standard checkerboard with opposite corners removed contains $64 - 2 = 62$ squares. The tiling would use $62/2 = 31$ dominoes. Note that each domino in this tiling covers one white and one black square. Consequently, the tiling covers 31 white squares and 31 black squares.

Suppose we can use dominoes to tile a standard checkerboard with opposite corners removed. Note that the standard checkerboard with opposite corners removed contains $64 - 2 = 62$ squares. The tiling would use $62/2 = 31$ dominoes. Note that each domino in this tiling covers one white and one black square. Consequently, the tiling covers 31 white squares and 31 black squares. However, when we remove two opposite corner squares, either 32 of the remaining squares are white and 30 are black or else 30 are white and 32 are black.

Suppose we can use dominoes to tile a standard checkerboard with opposite corners removed. Note that the standard checkerboard with opposite corners removed contains $64 - 2 = 62$ squares. The tiling would use $62/2 = 31$ dominoes. Note that each domino in this tiling covers one white and one black square. Consequently, the tiling covers 31 white squares and 31 black squares. However, when we remove two opposite corner squares, either 32 of the remaining squares are white and 30 are black or else 30 are white and 32 are black. This contradicts the assumption that we can use dominoes to cover a standard checkerboard with opposite corners removed, completing the proof.