Introduction to Logic

Department of Science and Mathematics

IIIT Guwahati.

August 13, 2020

Consider the statement "x is greater than 3".

Consider the statement "x is greater than 3".

- The first part, the variable *x*, is the subject of the statement.
- The second part the predicate, "is greater than 3" refers to a property that the subject of the statement can have.

Consider the statement "x is greater than 3".

- The first part, the variable *x*, is the subject of the statement.
- The second part the predicate, "is greater than 3" refers to a property that the subject of the statement can have.

Denote the statement "x is greater than 3" by P(x), where P denotes the predicate "is greater than 3" and x is the variable.

Consider the statement "x is greater than 3".

- The first part, the variable *x*, is the subject of the statement.
- The second part the predicate, "is greater than 3" refers to a property that the subject of the statement can have.

Denote the statement "x is greater than 3" by P(x), where P denotes the predicate "is greater than 3" and x is the variable.

- The statement P(x) is also said to be the value of the propositional function P at x.
- Once a value has been assigned to the variable x, the statement P(x) becomes a proposition and has a truth value.

Consider the statement "x is greater than 3".

- The first part, the variable x, is the subject of the statement.
- The second part the predicate, "is greater than 3" refers to a property that the subject of the statement can have.

Denote the statement "x is greater than 3" by P(x), where P denotes the predicate "is greater than 3" and x is the variable.

- The statement P(x) is also said to be the value of the propositional function P at x.
- Once a value has been assigned to the variable x, the statement P(x) becomes a proposition and has a truth value.

Example

Let P(x) denote the statement x > 3. What are the truth values of P(4) and P(2)?

Example

Let A(x) denote the statement : Computer x is under attack by an intruder. Suppose that of the computers on campus, only CS2 and MATH1 are currently under attack by intruders. What are truth values of A(CS1), A(CS2), and A(MATH1)?

Example

Let A(x) denote the statement : Computer x is under attack by an intruder. Suppose that of the computers on campus, only CS2 and MATH1 are currently under attack by intruders. What are truth values of A(CS1), A(CS2), and A(MATH1)?

Example

Let Q(x, y) denote the statement : x = y + 3. What are the truth values of the propositions Q(1, 2) and Q(3, 0)?

Quantifiers

Quantification expresses the extent to which a predicate is true over a range of elements. We will focus on two types of quantification here: universal quantification, which tells us that a predicate is true for every element under consideration (domain), and existential quantification, which tells us that there is one or more element under consideration for which the predicate is true.

Quantifiers

Quantification expresses the extent to which a predicate is true over a range of elements. We will focus on two types of quantification here: universal quantification, which tells us that a predicate is true for every element under consideration (domain), and existential quantification, which tells us that there is one or more element under consideration for which the predicate is true.

THE UNIVERSAL QUANTIFIER: The meaning of the universal quantification of P(x) changes when we change the domain. The domain must always be specified when a universal quantifier is used; without it, the universal quantification of a statement is not defined.

The universal quantification of P(x) is the statement P(x) for all values of x in the domain. The notation $\forall x \ P(x)$ denotes the universal quantification of P(x). Here $\forall x$ is called the universal quantifier. We read $\forall x \ P(x)$ as "for all x, P(x)" or "for every x, P(x)." An element for which P(x) is false is called a counterexample of $\forall x \ P(x)$.

The universal quantification of P(x) is the statement P(x) for all values of x in the domain. The notation $\forall x \ P(x)$ denotes the universal quantification of P(x). Here $\forall x$ is called the universal quantifier. We read $\forall x \ P(x)$ as "for all x, P(x)" or "for every x, P(x)." An element for which P(x) is false is called a counterexample of $\forall x \ P(x)$.

• Let P(x) be the statement "x + 1 > x." What is the truth value of the quantification $\forall x \ P(x)$, where the domain consists of all real numbers?

The universal quantification of P(x) is the statement P(x) for all values of x in the domain. The notation $\forall x \ P(x)$ denotes the universal quantification of P(x). Here $\forall x$ is called the universal quantifier. We read $\forall x \ P(x)$ as "for all x, P(x)" or "for every x, P(x)." An element for which P(x) is false is called a counterexample of $\forall x \ P(x)$.

- Let P(x) be the statement "x + 1 > x." What is the truth value of the quantification $\forall x \ P(x)$, where the domain consists of all real numbers?
- Let Q(x) be the statement "x < 2." What is the truth value of the quantification $\forall x \ Q(x)$, where the domain consists of all real numbers?

 What is the truth value of ∀x P(x), where P(x) is the statement x² < 10 and the domain consists of the positive integers not exceeding 4?

- What is the truth value of ∀x P(x), where P(x) is the statement x² < 10 and the domain consists of the positive integers not exceeding 4?
- What is the truth value of $\forall x(x^2 \ge x)$ if the domain consists of all real numbers?

- What is the truth value of ∀x P(x), where P(x) is the statement x² < 10 and the domain consists of the positive integers not exceeding 4?
- What is the truth value of $\forall x(x^2 \ge x)$ if the domain consists of all real numbers?
- What is the truth value of this statement if the domain consists of all integers?

- What is the truth value of ∀x P(x), where P(x) is the statement x² < 10 and the domain consists of the positive integers not exceeding 4?
- What is the truth value of $\forall x(x^2 \ge x)$ if the domain consists of all real numbers?
- What is the truth value of this statement if the domain consists of all integers?

The existential quantification of P(x) is the proposition "There exists an element x in the domain such that P(x)." We use the notation $\exists x \ P(x)$ for the existential quantification of P(x). Here \exists is called the existential quantifier.

The existential quantification of P(x) is the proposition "There exists an element x in the domain such that P(x)." We use the notation $\exists x \ P(x)$ for the existential quantification of P(x). Here \exists is called the existential quantifier.

A domain must always be specified when a statement $\exists x P(x)$ is used. We can also express existential quantification in many other ways, such as by using the words "for some, for at least one," or "there is."

• Let P(x) denote the statement x > 3. What is the truth value of the quantification $\exists x \ P(x)$, where the domain consists of all real numbers?

- Let P(x) denote the statement x > 3. What is the truth value of the quantification $\exists x \ P(x)$, where the domain consists of all real numbers?
- Let Q(x) denote the statement x = x + 1. What is the truth value of the quantification $\exists x \ Q(x)$, where the domain consists of all real numbers?

- Let P(x) denote the statement x > 3. What is the truth value of the quantification $\exists x \ P(x)$, where the domain consists of all real numbers?
- Let Q(x) denote the statement x = x + 1. What is the truth value of the quantification $\exists x \ Q(x)$, where the domain consists of all real numbers?
- What is the truth value of $\exists x \ P(x)$, where P(x) is the statement $x^2 > 10$ and the universe of discourse consists of the positive integers not exceeding 4?

- Let P(x) denote the statement x > 3. What is the truth value of the quantification $\exists x \ P(x)$, where the domain consists of all real numbers?
- Let Q(x) denote the statement x = x + 1. What is the truth value of the quantification $\exists x \ Q(x)$, where the domain consists of all real numbers?
- What is the truth value of $\exists x \ P(x)$, where P(x) is the statement $x^2 > 10$ and the universe of discourse consists of the positive integers not exceeding 4?

The quantifiers \forall and \exists have higher precedence than all logical operators from propositional calculus.