

Introduction to Logic

Department of Science and Mathematics

IIIT Guwahati.

August 17, 2020

Definition

Statements involving predicates and quantifiers are logically equivalent if and only if they have the same truth value no matter which predicates are substituted into these statements and which domain of discourse is used for the variables in these propositional functions. We use the notation $S \equiv T$ to indicate that two statements S and T involving predicates and quantifiers are logically equivalent.

Definition

Statements involving predicates and quantifiers are logically equivalent if and only if they have the same truth value no matter which predicates are substituted into these statements and which domain of discourse is used for the variables in these propositional functions. We use the notation $S \equiv T$ to indicate that two statements S and T involving predicates and quantifiers are logically equivalent.

1. Show that $\forall x(P(x) \wedge Q(x))$ and $\forall xP(x) \wedge \forall xQ(x)$ are logically equivalent (where the same domain is used throughout).

Definition

Statements involving predicates and quantifiers are logically equivalent if and only if they have the same truth value no matter which predicates are substituted into these statements and which domain of discourse is used for the variables in these propositional functions. We use the notation $S \equiv T$ to indicate that two statements S and T involving predicates and quantifiers are logically equivalent.

1. Show that $\forall x(P(x) \wedge Q(x))$ and $\forall xP(x) \wedge \forall xQ(x)$ are logically equivalent (where the same domain is used throughout).

Solution: Suppose we have particular predicates P and Q , with a common domain. We can show that $\forall x(P(x) \wedge Q(x))$ and $\forall xP(x) \wedge \forall xQ(x)$ are logically equivalent by proving $\forall x(P(x) \wedge Q(x))$ iff $\forall xP(x) \wedge \forall xQ(x)$.

Definition

Statements involving predicates and quantifiers are logically equivalent if and only if they have the same truth value no matter which predicates are substituted into these statements and which domain of discourse is used for the variables in these propositional functions. We use the notation $S \equiv T$ to indicate that two statements S and T involving predicates and quantifiers are logically equivalent.

1. Show that $\forall x(P(x) \wedge Q(x))$ and $\forall xP(x) \wedge \forall xQ(x)$ are logically equivalent (where the same domain is used throughout).

Solution: Suppose we have particular predicates P and Q , with a common domain. We can show that $\forall x(P(x) \wedge Q(x))$ and $\forall xP(x) \wedge \forall xQ(x)$ are logically equivalent by proving $\forall x(P(x) \wedge Q(x))$ iff $\forall xP(x) \wedge \forall xQ(x)$.

First, we show that if $\forall x(P(x) \wedge Q(x))$ is true, then $\forall xP(x) \wedge \forall xQ(x)$ is true. Second, we show that if $\forall xP(x) \wedge \forall xQ(x)$ is true, then $\forall x(P(x) \wedge Q(x))$ is true.

Logical equivalence of statements involving predicates and quantifiers

- Let $\forall x(P(x) \wedge Q(x))$ is true.

Logical equivalence of statements involving predicates and quantifiers

- Let $\forall x(P(x) \wedge Q(x))$ is true.
- i.e If a is any element (arbitrary element) in domain, then $P(a) \wedge Q(a)$ is true.

Logical equivalence of statements involving predicates and quantifiers

- Let $\forall x(P(x) \wedge Q(x))$ is true.
- i.e If a is any element (arbitrary element) in domain, then $P(a) \wedge Q(a)$ is true.
- $P(a)$ is true and $Q(a)$ is true

Logical equivalence of statements involving predicates and quantifiers

- Let $\forall x(P(x) \wedge Q(x))$ is true.
- i.e If a is any element (arbitrary element) in domain, then $P(a) \wedge Q(a)$ is true.
- $P(a)$ is true and $Q(a)$ is true
- $\forall xP(x)$ is true and $\forall xQ(x)$ is true.

Logical equivalence of statements involving predicates and quantifiers

- Let $\forall x(P(x) \wedge Q(x))$ is true.
- i.e If a is any element (arbitrary element) in domain, then $P(a) \wedge Q(a)$ is true.
- $P(a)$ is true and $Q(a)$ is true
- $\forall xP(x)$ is true and $\forall xQ(x)$ is true.
- Thus, $\forall xP(x) \wedge \forall xQ(x)$ is true.

Therefore

- We can distribute a universal quantifier over a conjunction.

Therefore

- We can distribute a universal quantifier over a conjunction.
- Can we distribute a universal quantifier over a disjunction?

Therefore

- We can distribute a universal quantifier over a conjunction.
- Can we distribute a universal quantifier over a disjunction?
- Can we distribute an existential quantifier over a conjunction?

Therefore

- We can distribute a universal quantifier over a conjunction.
- Can we distribute a universal quantifier over a disjunction?
- Can we distribute an existential quantifier over a conjunction?
- Can we distribute an existential quantifier over a disjunction?

Therefore

- We can distribute a universal quantifier over a conjunction.
- Can we distribute a universal quantifier over a disjunction?
- Can we distribute an existential quantifier over a conjunction?
- Can we distribute an existential quantifier over a disjunction?

Negating Quantified Expressions

Consider the negation of the statement

Negating Quantified Expressions

Consider the negation of the statement

- Every student in your class has taken a course in calculus.

Negating Quantified Expressions

Consider the negation of the statement

- Every student in your class has taken a course in calculus.
Let $P(x)$ be the statement "x has taken a course in calculus" and the domain consists of the students in your class.

Negating Quantified Expressions

Consider the negation of the statement

- Every student in your class has taken a course in calculus.
Let $P(x)$ be the statement "x has taken a course in calculus" and the domain consists of the students in your class.
i.e $\forall x P(x)$

Negating Quantified Expressions

Consider the negation of the statement

- Every student in your class has taken a course in calculus.
Let $P(x)$ be the statement "x has taken a course in calculus" and the domain consists of the students in your class.
i.e $\forall x P(x)$
- Negation: It is not the case that every student in your class has taken a course in calculus.

Negating Quantified Expressions

Consider the negation of the statement

- Every student in your class has taken a course in calculus.
Let $P(x)$ be the statement "x has taken a course in calculus" and the domain consists of the students in your class.
i.e $\forall x P(x)$
- Negation: It is not the case that every student in your class has taken a course in calculus.
i.e There is a student in your class who has not taken a course in calculus.

Negating Quantified Expressions

Consider the negation of the statement

- Every student in your class has taken a course in calculus.
Let $P(x)$ be the statement "x has taken a course in calculus" and the domain consists of the students in your class.
i.e $\forall x P(x)$
- Negation: It is not the case that every student in your class has taken a course in calculus.
i.e There is a student in your class who has not taken a course in calculus.
i.e $\exists \neg P(x)$

Negating Quantified Expressions

Consider the negation of the statement

- Every student in your class has taken a course in calculus.
Let $P(x)$ be the statement "x has taken a course in calculus" and the domain consists of the students in your class.
i.e $\forall x P(x)$
- Negation: It is not the case that every student in your class has taken a course in calculus.
i.e There is a student in your class who has not taken a course in calculus.
i.e $\exists \neg P(x)$
- Is $\neg(\forall x P(x)) \equiv \exists \neg P(x)$?

Prove that $\neg(\forall x P(x)) \equiv \exists \neg P(x)$

Prove that $\neg(\forall x P(x)) \equiv \exists \neg P(x)$

- $\neg(\forall x P(x))$ is true if and only if $\forall x P(x)$ is false.

Prove that $\neg(\forall x P(x)) \equiv \exists \neg P(x)$

- $\neg(\forall x P(x))$ is true if and only if $\forall x P(x)$ is false.
- $\forall x P(x)$ is false if and only if there is an element x in the domain for which $P(x)$ is false.

Prove that $\neg(\forall x P(x)) \equiv \exists \neg P(x)$

- $\neg(\forall x P(x))$ is true if and only if $\forall x P(x)$ is false.
- $\forall x P(x)$ is false if and only if there is an element x in the domain for which $P(x)$ is false.
- This holds if and only if there is an element x in the domain for which $\neg P(x)$ is true.

Prove that $\neg(\forall x P(x)) \equiv \exists \neg P(x)$

- $\neg(\forall x P(x))$ is true if and only if $\forall x P(x)$ is false.
- $\forall x P(x)$ is false if and only if there is an element x in the domain for which $P(x)$ is false.
- This holds if and only if there is an element x in the domain for which $\neg P(x)$ is true.
- Finally, there is an element x in the domain for which $\neg P(x)$ is true if and only if $\exists x \neg P(x)$ is true.

Consider the statement:

- There is a student in this class who has taken a course in calculus.

Consider the statement:

- There is a student in this class who has taken a course in calculus.
Let $Q(x)$ be the statement "x has taken a course in calculus."

Consider the statement:

- There is a student in this class who has taken a course in calculus.
Let $Q(x)$ be the statement "x has taken a course in calculus."
i.e $\exists x Q(x)$

Consider the statement:

- There is a student in this class who has taken a course in calculus.
Let $Q(x)$ be the statement "x has taken a course in calculus."
i.e $\exists x Q(x)$
- Negation: It is not the case that there is a student in this class who has taken a course in calculus.

Consider the statement:

- There is a student in this class who has taken a course in calculus.
Let $Q(x)$ be the statement "x has taken a course in calculus."
i.e $\exists x Q(x)$
- Negation: It is not the case that there is a student in this class who has taken a course in calculus.
i.e "Every student in this class has not taken calculus"

Consider the statement:

- There is a student in this class who has taken a course in calculus.
Let $Q(x)$ be the statement "x has taken a course in calculus."
i.e $\exists x Q(x)$
- Negation: It is not the case that there is a student in this class who has taken a course in calculus.
i.e "Every student in this class has not taken calculus"
i.e $\forall x \neg Q(x)$.

Consider the statement:

- There is a student in this class who has taken a course in calculus.
Let $Q(x)$ be the statement "x has taken a course in calculus."
i.e $\exists x Q(x)$
- Negation: It is not the case that there is a student in this class who has taken a course in calculus.
i.e "Every student in this class has not taken calculus"
i.e $\forall x \neg Q(x)$.
- Is $\neg \exists Q(x) \equiv \forall x \neg Q(x)$?

Consider the statement:

- There is a student in this class who has taken a course in calculus.
Let $Q(x)$ be the statement "x has taken a course in calculus."
i.e $\exists x Q(x)$
- Negation: It is not the case that there is a student in this class who has taken a course in calculus.
i.e "Every student in this class has not taken calculus"
i.e $\forall x \neg Q(x)$.
- Is $\neg \exists Q(x) \equiv \forall x \neg Q(x)$?

Prove that $\neg \exists Q(x) \equiv \forall x \neg Q(x)$.

Prove that $\neg \exists Q(x) \equiv \forall x \neg Q(x)$.

- $\neg \exists x Q(x)$ is true if and only if $\exists x Q(x)$ is false.

Prove that $\neg \exists x Q(x) \equiv \forall x \neg Q(x)$.

- $\neg \exists x Q(x)$ is true if and only if $\exists x Q(x)$ is false.
- This is true if and only if no x exists in the domain for which $Q(x)$ is true.

Prove that $\neg \exists x Q(x) \equiv \forall x \neg Q(x)$.

- $\neg \exists x Q(x)$ is true if and only if $\exists x Q(x)$ is false.
- This is true if and only if no x exists in the domain for which $Q(x)$ is true.
- This is true if and only if $Q(x)$ is false for every x in the domain.

Prove that $\neg \exists x Q(x) \equiv \forall x \neg Q(x)$.

- $\neg \exists x Q(x)$ is true if and only if $\exists x Q(x)$ is false.
- This is true if and only if no x exists in the domain for which $Q(x)$ is true.
- This is true if and only if $Q(x)$ is false for every x in the domain.
- $Q(x)$ is false for every x in the domain if and only if $\neg Q(x)$ is true for all x in the domain

Prove that $\neg \exists x Q(x) \equiv \forall x \neg Q(x)$.

- $\neg \exists x Q(x)$ is true if and only if $\exists x Q(x)$ is false.
- This is true if and only if no x exists in the domain for which $Q(x)$ is true.
- This is true if and only if $Q(x)$ is false for every x in the domain.
- $Q(x)$ is false for every x in the domain if and only if $\neg Q(x)$ is true for all x in the domain
- This holds if and only if $\forall x \neg Q(x)$ is true.

Prove that $\neg \exists x Q(x) \equiv \forall x \neg Q(x)$.

- $\neg \exists x Q(x)$ is true if and only if $\exists x Q(x)$ is false.
- This is true if and only if no x exists in the domain for which $Q(x)$ is true.
- This is true if and only if $Q(x)$ is false for every x in the domain.
- $Q(x)$ is false for every x in the domain if and only if $\neg Q(x)$ is true for all x in the domain
- This holds if and only if $\forall x \neg Q(x)$ is true.

We conclude that $\neg \exists x Q(x)$ and $\forall x \neg Q(x)$ are logically equivalent.

Prove that $\neg \exists x Q(x) \equiv \forall x \neg Q(x)$.

- $\neg \exists x Q(x)$ is true if and only if $\exists x Q(x)$ is false.
- This is true if and only if no x exists in the domain for which $Q(x)$ is true.
- This is true if and only if $Q(x)$ is false for every x in the domain.
- $Q(x)$ is false for every x in the domain if and only if $\neg Q(x)$ is true for all x in the domain
- This holds if and only if $\forall x \neg Q(x)$ is true.

We conclude that $\neg \exists x Q(x)$ and $\forall x \neg Q(x)$ are logically equivalent.

- De Morgan's laws for quantifiers

$$\neg(\forall x P(x)) \equiv \exists x \neg P(x)$$

$$\neg(\exists x P(x)) \equiv \forall x \neg P(x)$$

When the domain has n elements x_1, x_2, \dots, x_n , it follows that $\neg \forall x P(x)$ is the same as $\neg(P(x_1) \wedge P(x_2) \wedge \dots \wedge P(x_n))$, which is equivalent to $\neg P(x_1) \vee \neg P(x_2) \vee \dots \vee \neg P(x_n)$ by De Morgan's laws, and this is the same as $\exists x \neg P(x)$. Similarly, $\neg \exists x P(x)$ is the same as $\neg(P(x_1) \vee P(x_2) \vee \dots \vee P(x_n))$, which by De Morgans laws is equivalent to $\neg P(x_1) \wedge \neg P(x_2) \wedge \dots \wedge \neg P(x_n)$, and this is the same as $\forall x \neg P(x)$.

Find negations of the statements

- There is an honest politician.

Find negations of the statements

- There is an honest politician.
Let $H(x)$ denote "x is honest."

Find negations of the statements

- There is an honest politician.

Let $H(x)$ denote "x is honest."

Then the statement "There is an honest politician" is represented

Find negations of the statements

- There is an honest politician.

Let $H(x)$ denote "x is honest."

Then the statement "There is an honest politician" is represented by $\exists x H(x)$, where the domain consists of all politicians.

Find negations of the statements

- There is an honest politician.

Let $H(x)$ denote "x is honest."

Then the statement "There is an honest politician" is represented by $\exists x H(x)$, where the domain consists of all politicians.

- The negation is $\neg \exists x H(x) \equiv \forall x \neg H(x)$.

Find negations of the statements

- There is an honest politician.

Let $H(x)$ denote "x is honest."

Then the statement "There is an honest politician" is represented by $\exists x H(x)$, where the domain consists of all politicians.

- The negation is $\neg \exists x H(x) \equiv \forall x \neg H(x)$.

This negation can be expressed as "Every politician is dishonest."

Find negations of the statements

- There is an honest politician.

Let $H(x)$ denote "x is honest."

Then the statement "There is an honest politician" is represented by $\exists x H(x)$, where the domain consists of all politicians.

- The negation is $\neg \exists x H(x) \equiv \forall x \neg H(x)$.

This negation can be expressed as "Every politician is dishonest."

Note: In English, the statement "All politicians are not honest" is ambiguous. In common usage, this statement often means "Not all politicians are honest." Consequently, we do not use this statement to express this negation.

- All Indians drink tea.

- All Indians drink tea.

Let $C(x)$ denote "x drinks tea." Then the given statement is represented by $\forall x C(x)$, where the domain consists of all Indians.

- All Indians drink tea.

Let $C(x)$ denote "x drinks tea." Then the given statement is represented by $\forall x C(x)$, where the domain consists of all Indians.

- Negation is $\neg \forall x C(x) \equiv \exists x \neg C(x)$.

- All Indians drink tea.

Let $C(x)$ denote "x drinks tea." Then the given statement is represented by $\forall x C(x)$, where the domain consists of all Indians.

- Negation is $\neg \forall x C(x) \equiv \exists x \neg C(x)$.

This negation can be expressed in several different ways, including "Some Indian does not drink tea" and "There is an Indian who does not drink tea."

- What are the negations of the statements

$$\forall x (x^2 > x)$$

$$\exists x (x^2 = 2)?$$

- All Indians drink tea.

Let $C(x)$ denote "x drinks tea." Then the given statement is represented by $\forall x C(x)$, where the domain consists of all Indians.

- Negation is $\neg \forall x C(x) \equiv \exists x \neg C(x)$.

This negation can be expressed in several different ways, including "Some Indian does not drink tea" and "There is an Indian who does not drink tea."

- What are the negations of the statements

$$\forall x (x^2 > x)$$

$$\exists x (x^2 = 2)? \text{ (Truth values will depend on the domain.)}$$

- Show that $\neg\forall x(P(x) \rightarrow Q(x))$ and $\exists x(P(x) \wedge \neg Q(x))$ are logically equivalent.

- Show that $\neg\forall x(P(x) \rightarrow Q(x))$ and $\exists x(P(x) \wedge \neg Q(x))$ are logically equivalent. (Hint: $p \rightarrow q \equiv \neg p \vee q$)

Translating from English into Logical Expressions

1. Express the statement "Every student in this class has studied calculus" using predicates and quantifiers.

Translating from English into Logical Expressions

1. Express the statement "Every student in this class has studied calculus" using predicates and quantifiers.
 - "For every student in this class, that student has studied calculus."

Translating from English into Logical Expressions

1. Express the statement "Every student in this class has studied calculus" using predicates and quantifiers.
 - "For every student in this class, that student has studied calculus."
 - Introduce a variable x so that our statement becomes "For every student x in this class, x has studied calculus".

Translating from English into Logical Expressions

1. Express the statement "Every student in this class has studied calculus" using predicates and quantifiers.

- "For every student in this class, that student has studied calculus."
- Introduce a variable x so that our statement becomes "For every student x in this class, x has studied calculus".
- Introduce $C(x)$: " x has studied calculus."

Translating from English into Logical Expressions

1. Express the statement "Every student in this class has studied calculus" using predicates and quantifiers.

- "For every student in this class, that student has studied calculus."
- Introduce a variable x so that our statement becomes "For every student x in this class, x has studied calculus".
- Introduce $C(x)$: " x has studied calculus."

If the domain for x consists of the students in the class, we can translate our statement as

Translating from English into Logical Expressions

1. Express the statement "Every student in this class has studied calculus" using predicates and quantifiers.

- "For every student in this class, that student has studied calculus."
- Introduce a variable x so that our statement becomes "For every student x in this class, x has studied calculus".
- Introduce $C(x)$: " x has studied calculus."

If the domain for x consists of the students in the class, we can translate our statement as $\forall x C(x)$.

Translating from English into Logical Expressions

1. Express the statement "Every student in this class has studied calculus" using predicates and quantifiers.

- "For every student in this class, that student has studied calculus."
- Introduce a variable x so that our statement becomes "For every student x in this class, x has studied calculus".
- Introduce $C(x)$: " x has studied calculus."

If the domain for x consists of the students in the class, we can translate our statement as $\forall x C(x)$.

If we change the domain to consist of all people, we will need to express our statement as

Translating from English into Logical Expressions

1. Express the statement "Every student in this class has studied calculus" using predicates and quantifiers.

- "For every student in this class, that student has studied calculus."
- Introduce a variable x so that our statement becomes "For every student x in this class, x has studied calculus".
- Introduce $C(x)$: " x has studied calculus."

If the domain for x consists of the students in the class, we can translate our statement as $\forall x C(x)$.

If we change the domain to consist of all people, we will need to express our statement as

- "For every person x , if person x is a student in this class then x has studied calculus."

Translating from English into Logical Expressions

1. Express the statement "Every student in this class has studied calculus" using predicates and quantifiers.

- "For every student in this class, that student has studied calculus."
- Introduce a variable x so that our statement becomes "For every student x in this class, x has studied calculus".
- Introduce $C(x)$: " x has studied calculus."

If the domain for x consists of the students in the class, we can translate our statement as $\forall x C(x)$.

If we change the domain to consist of all people, we will need to express our statement as

- "For every person x , if person x is a student in this class then x has studied calculus."

If $S(x)$ represents the statement that person x is in this class, we see that our statement can be expressed as

Translating from English into Logical Expressions

1. Express the statement "Every student in this class has studied calculus" using predicates and quantifiers.

- "For every student in this class, that student has studied calculus."
- Introduce a variable x so that our statement becomes "For every student x in this class, x has studied calculus".
- Introduce $C(x)$: " x has studied calculus."

If the domain for x consists of the students in the class, we can translate our statement as $\forall x C(x)$.

If we change the domain to consist of all people, we will need to express our statement as

- "For every person x , if person x is a student in this class then x has studied calculus."

If $S(x)$ represents the statement that person x is in this class, we see that our statement can be expressed as $\forall x (S(x) \rightarrow C(x))$.

Translating from English into Logical Expressions

1. Express the statement "Every student in this class has studied calculus" using predicates and quantifiers.

- "For every student in this class, that student has studied calculus."
- Introduce a variable x so that our statement becomes "For every student x in this class, x has studied calculus".
- Introduce $C(x)$: " x has studied calculus."

If the domain for x consists of the students in the class, we can translate our statement as $\forall x C(x)$.

If we change the domain to consist of all people, we will need to express our statement as

- "For every person x , if person x is a student in this class then x has studied calculus."

If $S(x)$ represents the statement that person x is in this class, we see that our statement can be expressed as $\forall x (S(x) \rightarrow C(x))$.

What does $\forall x (S(x) \wedge C(x))$ represent?