

□ Question - 1 :

• problem : 8 coins are thrown simultaneously. Find the chance of obtaining — (i) at least 6 heads, (ii) no heads, and (iii) all heads.

• Manual Calculation : Here,  $n = 8$ ,  $p = \frac{1}{2}$ ,  $q = 1 - p = \frac{1}{2}$ .

$$\begin{aligned}
 \text{(i)} \quad P(X \geq 6) &= P(X=6) + P(X=7) + P(X=8) \\
 &= {}^8C_6 \left(\frac{1}{2}\right)^6 \left(\frac{1}{2}\right)^2 + {}^8C_7 \left(\frac{1}{2}\right)^7 \left(\frac{1}{2}\right)^1 + {}^8C_8 \left(\frac{1}{2}\right)^8 \left(\frac{1}{2}\right)^0 \\
 &= 2^8 \times \frac{1}{2^8} + 8 \times \frac{1}{2^8} + 1 \times \frac{1}{2^8} \\
 &= \frac{37}{256} = 0.1445
 \end{aligned}$$

$$\begin{aligned}
 \text{(ii)} \quad P(X=0) &= {}^8C_0 \left(\frac{1}{2}\right)^8 \left(\frac{1}{2}\right)^0 \\
 &= \frac{1}{2^8} \\
 &= \frac{1}{256} = 0.0039
 \end{aligned}$$

$$\begin{aligned}
 \text{(iii)} \quad P(X=8) &= {}^8C_8 \left(\frac{1}{2}\right)^8 \left(\frac{1}{2}\right)^0 \\
 &= \frac{1}{2^8} \\
 &= 0.0039
 \end{aligned}$$

• Code in R-console :

```
> 1 - pbinom(5, 8, 0.5)
```

```
// P(X ≥ 6)
```

```
[1] 0.1445312
```

```
> dbinom(0, 8, 0.5)
```

```
// P(X = 0)
```

```
[1] 0.00390625
```

```
> dbinom(8, 8, 0.5)
```

```
// P(X = 8)
```

```
[1] 0.00390625
```

□ Question - 2 :

- problem : A multi-choice test questions consists of 8 questions with 3 answers to each question (of which only one is correct). A student answers each question by rolling a balanced dice and checking the first answer if he gets 1 or 2, the second answer if he gets 3 or 4 and the three answers if he gets 5 or 6. To get a distinction, the student must secure at least 75% correct answers. If there is no negative marking, what is the probability that the student secures a distinction?

- Manual Calculation : Here,  $p = \frac{1}{3}$ ,  $q = 1 - p = \frac{2}{3}$ ,  $n = 8$

Acc. to quest<sup>n</sup>, in order to get at least 75%, the student must answer at least 6 or more questions.

$$\begin{aligned}
 \therefore P(X \geq 6) &= P(X=6) + P(X=7) + P(X=8) \\
 &= {}^8C_6 \left(\frac{1}{3}\right)^6 \left(\frac{2}{3}\right)^2 + {}^8C_7 \left(\frac{1}{3}\right)^7 \left(\frac{2}{3}\right)^1 \\
 &\quad + {}^8C_8 \left(\frac{1}{3}\right)^8 \left(\frac{2}{3}\right)^0 \\
 &= 28 \times \frac{1}{729} \times \frac{4}{9} + 8 \times \frac{1}{2187} \times \frac{2}{3} \\
 &\quad + 1 \times \frac{1}{6561} \\
 &= 0.01707 + 0.002438 + 0.00015 \\
 &= 0.01965
 \end{aligned}$$

- Code in R-console :

> 1 - pbinom(5, 8,  $\frac{1}{3}$ )

//  $P(X \geq 6)$

[1] 0.01966164

### □ Question-3 :

# 19BCE2105

- problem : In a town 10 accidents took place in a span of 50 days. Assuming that the number of accidents per day follows the Poisson distribution, find the probability that there will be 3 or more accidents in a day.

- Manual calculation : Here,  $\lambda = \frac{10}{50} = \frac{1}{5}$

$$\begin{aligned}
 P(X \geq 3) &= 1 - [P(X=0) + P(X=1) + P(X=2)] \\
 &= 1 - \left[ e^{-\frac{1}{5}} \left( \frac{(\frac{1}{5})^0}{0!} + \frac{(\frac{1}{5})^1}{1!} + \frac{(\frac{1}{5})^2}{2!} \right) \right] \\
 &= 1 - [2.7183]^{-\frac{1}{5}} \left( 1 + \frac{1}{5} + \frac{1}{50} \right) \\
 &= 1 - (2.7183)^{-\frac{1}{5}} \times \frac{61}{50} \\
 &= 1 - 0.81873 \times \frac{61}{50} \\
 &= 0.0011494
 \end{aligned}$$

- Code in R console :

```

> 1 - ppois(2, lambda = 1/5) // P(X ≥ 3)
[1] 0.001148481

```

### □ Question-4 :


- problem : 1000 light bulbs with a mean life of 120 days are installed in a new factory; their length of life is normally distributed with standard deviation 20 days. (i) how many bulbs will expire in less than 90 days? (ii) if it is decided to replace all the bulbs together, what intervals should be allowed between replacements if not more than 10% should expire before replacement?

• Manual calculation:

Here,  $\mu = 120$ ,  $\sigma = 20$

(i)  $x = 90$

$$\therefore Z = \frac{x - \mu}{\sigma} = \frac{90 - 120}{20} = -1.5$$



$\therefore$  No. of bulbs that will expire in less than 90 days

$$= A_2 \times 1000$$

$$= 0.06681 \times 1000$$

$$= 66.8 \text{ bulbs}$$

$$\approx 67 \text{ bulbs}$$

(ii)  $P(x \leq x_1) = 0.1$

$$P\left(z \leq \frac{x_1 - 120}{20}\right) = 0.1$$

or,  $\frac{x_1 - 120}{20} = -1.3$  [from table]

$$\therefore x_1 = (-1.3 \times 20) + 120$$

$$= (120 - 26) \text{ days}$$

$$= 94 \text{ days}$$



Q Code in R-console :-

```
> pnorm(90, mean = 120, sd = 20) * 1000
```

[1] 66.8072

— END —