

Certified Local Weil Positivity in the Even Pole-Moment-Null Sector through Radius $3/5$

Sharan Thota

Abstract

We study the compactly supported Weil quadratic form associated with the Riemann zeta function after restriction to even functions whose $\cosh(x/2)$ -moment vanishes. For a support radius a , its Fourier multiplier consists of the archimedean digamma term and the prime-power atoms with $\log n < 2a$. We prove, with a reproducible interval-arithmetic certificate, that the resulting closed form satisfies

$$T_a[f] \geq \frac{3}{100000} \|f\|_2^2 \quad (0 < a \leq 3/5).$$

At the endpoint, the arithmetic part contains exactly the atoms $n = 2, 3$. The proof is an infinite-dimensional estimate, not a finite-section Ritz calculation. We split the form into a compact adverse operator and a positive favorable form, retain a trace-class part of the favorable form, and control the discarded directions by a matrix-valued Schur estimate. A rank-44 interval LDL^* certificate then gives the stated rational gap. We also prove a general completeness theorem for this positive-rescue scheme, derive an exact positive-background Poisson-frame representation, and record the corresponding normalized shorted-graph Loewner inequality. The result is local in the support radius; the all-radius positivity criterion, which is equivalent to the Riemann hypothesis, is not asserted.

2020 Mathematics Subject Classification. 65G40, 47A07, 47B65, 11M26.

Keywords. Validated numerics, interval arithmetic, localized quadratic form, Schur complement, Poisson frame, Weil's criterion, Riemann zeta function.

1 Introduction

Let Λ denote the von Mangoldt function and $\psi = \Gamma'/\Gamma$ the digamma function. For $a > 0$, the compact-support explicit formula leads to the real multiplier

$$b_a(t) = \operatorname{Re} \psi\left(\frac{1}{4} + \frac{it}{2}\right) - \log \pi - 2 \sum_{1 < n < e^{2a}} \frac{\Lambda(n)}{\sqrt{n}} \cos(t \log n). \quad (1)$$

The finite sum changes only when $2a$ crosses the logarithm of a prime power. This makes fixed-support positivity a natural local problem: it retains exact arithmetic coefficients while involving only finitely many prime-power translations.

The purpose of this paper is to establish a strict local theorem after the first two arithmetic atoms have entered. The principal result is the following. All notation is defined in Section 2.

Theorem 1.1 (Prime-inclusive local coercivity). *For every $0 < a \leq 3/5$, the closed form T_a on the complex-even, $\cosh(x/2)$ -moment-null space satisfies*

$$T_a[f] \geq \frac{3}{100000} \|f\|_2^2, \quad f \in \mathcal{Q}_a^0. \quad (2)$$

At $a = 3/5$, the prime-power sum in (1) contains exactly $n = 2$ and $n = 3$.

The proof has three parts. First, a projected cosine-frame representation turns multiplication by the negative part of the symbol into a positive trace-class operator. Second, a favorable, sign-certified portion of the positive part is retained without discarding its orientation relative to the adverse operator. A matrix Schur estimate controls the infinite orthogonal complement by trace tails. Third, outward-rounded Arb arithmetic verifies the resulting finite matrix inequalities and every scalar sign decision. Exact zero-extension then transfers the endpoint bound to all smaller supports, including across the thresholds $(\log 2)/2$ and $(\log 3)/2$.

The finite-dimensional space in the certificate is used only to bound the complement of a trace-class operator. No smallest eigenvalue of a compression of T_a is used as a lower bound for the continuum form; such a use would have the wrong variational direction.

The connection with the Riemann hypothesis is contextual rather than a claim of this paper. Weil’s positivity criterion, in its localized Hermitian-form version, identifies the Riemann hypothesis with positivity for every support radius. Theorem 1.1 supplies one explicit initial interval of radii and therefore does not settle the all-radius problem; the precise statement is recorded in Section 9.

The localization of Weil’s quadratic form was initiated by Yoshida, who proved positivity for sufficiently small support and reduced the problem to finite calculations on suitable localized spaces. Bombieri subsequently developed the associated variational problem and recorded the classical calculation at the prime-free radius $a = (\log 2)/2$ [3, 4, 5]. Connes and Consani gave an operator-theoretic explanation of the archimedean inequality at that scale and later reported numerical semi-local evidence across successive arithmetic thresholds [6, 7]. Connes and van Suijlekom studied the quadratic forms attached to finite truncations and proved an extension of the Carathéodory–Fejér theorem for Toeplitz matrices supplying the required self-adjointness [8]. Building on that framework, Connes, Consani and Moscovici construct self-adjoint rank-one perturbations of the scaling spectral triple on an interval whose spectra approach the low-lying zeros; their convergence is reported numerically, and a rigorous proof of it would give the Riemann hypothesis [9]. Suzuki constructed the localized Friedrichs operator and proved continuity of its lowest eigenvalue, but his unconditional positivity theorem remains qualitative for sufficiently small a [10, 11]. A recent finite-Galerkin study supplies an exact dictionary between real even coefficient vectors and band-limited Guinand–Weil test functions, together with a certification rule for finite-cutoff positivity, but not a lower bound for the exact infinite-dimensional pole-moment-null continuum form [12].

To the author’s knowledge, Theorem 1.1 is the first explicit rigorous continuum lower bound in the even pole-moment-null sector at a support radius beyond the $n = 3$ entry threshold. This comparison is restricted to that sector; it does not concern Suzuki’s unrestricted localized operator or the all-radius problem.

The paper is organized as follows. Section 2 fixes the Fourier normalization and the form domain. Section 3 gives the exact projected cosine kernel. Section 4 proves the abstract positive-rescue estimates. Section 5 gives the complete theorem-bearing certificate at $a = 3/5$. Section 6 proves inward nesting, Section 7 gives the exact grouped Poisson frame, and Section 8 records the shorted-form consequence. Section 9 records the all-radius criterion that delimits the scope of the result. Reproduction data appear in Appendix A.

2 The local Weil form

Write $I_a = (-a, a)$, and identify every function on I_a with its zero extension to $\mathbb{R}$. We use

$$\widehat{f}(t) = \int_{-a}^a f(x) e^{-itx} dx, \quad \|f\|_2^2 = \frac{1}{2\pi} \int_{\mathbb{R}} |\widehat{f}(t)|^2 dt. \quad (3)$$

Let

$$\mathcal{V}_a^0 = \left\{ f \in L^2(I_a; \mathbb{C}) : f(-x) = f(x) \text{ a.e., } \int_{-a}^a f(x) \cosh(x/2) dx = 0 \right\}. \quad (4)$$

For an even function the odd $\sinh(x/2)$ -moment vanishes, and hence

$$\widehat{f}(i/2) = \widehat{f}(-i/2) = 0 \iff \int_{-a}^a f(x) \cosh(x/2) dx = 0. \quad (5)$$

Thus the single condition in (4) removes both pole terms from the even restriction of the Weil form.

Put

$$g_\infty(t) = \operatorname{Re} \psi\left(\frac{1}{4} + \frac{it}{2}\right) - \log \pi. \quad (6)$$

Since $g_\infty(t) = \log(2 + |t|) + O(1)$, while the arithmetic part of (1) is bounded for fixed a , the natural closed form domain is

$$\mathcal{Q}_a^0 = \left\{ f \in \mathcal{V}_a^0 : \int_{\mathbb{R}} \log(2 + |t|) |\widehat{f}(t)|^2 dt < \infty \right\}. \quad (7)$$

On this domain define

$$T_a[f, g] = \frac{1}{2\pi} \int_{\mathbb{R}} b_a(t) \widehat{f}(t) \overline{\widehat{g}(t)} dt, \quad T_a[f] = T_a[f, f]. \quad (8)$$

The form is densely defined, closed, and bounded below. Including an atom with $\log n = 2a$ does not alter the compressed form: the corresponding translation meets the support only on a null set.

3 Projected cosine frames

Let $m_a(x) = \cosh(x/2)$, and let P_a^0 be the orthogonal projection from the even subspace of $L^2(I_a)$ onto $m_a^\perp$. For $t \geq 0$ put

$$v_{a,t} = P_a^0 \cos(tx). \quad (9)$$

For $f \in \mathcal{V}_a^0$, evenness and the moment constraint give

$$\widehat{f}(t) = \langle f, v_{a,t} \rangle. \quad (10)$$

Consequently, if $w \geq 0$ is integrable and compactly supported, then

$$K_{a,w} = \frac{1}{\pi} \int_0^\infty w(t) |v_{a,t}\rangle \langle v_{a,t}| dt \quad (11)$$

is positive and trace class, with

$$\langle K_{a,w} f, f \rangle = \frac{1}{\pi} \int_0^\infty w(t) |\widehat{f}(t)|^2 dt, \quad \operatorname{Tr} K_{a,w} = \frac{1}{\pi} \int_0^\infty w(t) \|v_{a,t}\|^2 dt. \quad (12)$$

The kernel needed by the certificate is elementary. With $\text{sinc } z = \sin z/z$, continuously extended at zero,

$$\|m_a\|_2^2 = a + \sinh a, \quad (13)$$

$$\langle m_a, \cos(tx) \rangle = 2 \frac{\frac{1}{2} \sinh(a/2) \cos(at) + t \cosh(a/2) \sin(at)}{t^2 + 1/4}, \quad (14)$$

$$\begin{aligned} \langle v_{a,s}, v_{a,t} \rangle &= a \{ \text{sinc}(a(s-t)) + \text{sinc}(a(s+t)) \} \\ &\quad - \frac{\langle m_a, \cos(sx) \rangle \langle m_a, \cos(tx) \rangle}{a + \sinh a}. \end{aligned} \quad (15)$$

These identities make both the Gram matrix and all operator compressions amenable to rigorous interval integration.

4 Positive rescue and a matrix Schur estimate

4.1 The canonical split

Let b be any real multiplier satisfying $b(t) \rightarrow +\infty$, fix $\mu > 0$, and set

$$w = (\mu - b)_+, \quad d = (b - \mu)_+. \quad (16)$$

On a projected cosine space as above, let

$$A = K_w, \quad D[f] = \frac{1}{\pi} \int_0^\infty d(t) |\widehat{f}(t)|^2 dt. \quad (17)$$

Then A is positive trace class, D is a closed positive form, and the multiplier form has the exact decomposition

$$T = \mu I - A + D. \quad (18)$$

If C is positive trace class and $0 \preceq C \preceq D$ in form sense, then

$$T \succeq \mu I - (A - C). \quad (19)$$

4.2 The matrix residual

We first isolate the operator-theoretic estimate used in the certificate.

Lemma 4.1 (Schur residual). *Let $S = S^*$ be bounded, let P be an orthogonal projection, and put $Q = I - P$. Write*

$$D_P = PSP, \quad X = QSP, \quad E = QSQ.$$

*Suppose that $E \preceq \delta Q$ and $X^*X \preceq R_P$, where $R_P \succeq 0$ acts on $P\mathcal{H}$. If $\lambda > \delta$ and*

$$\lambda P - D_P - \frac{R_P}{\lambda - \delta} \succeq 0, \quad (20)$$

then $S \preceq \lambda I$.

Proof. Relative to $P\mathcal{H} \oplus Q\mathcal{H}$,

$$\lambda I - S = \begin{pmatrix} \lambda P - D_P & -X^* \\ -X & \lambda Q - E \end{pmatrix}.$$

The lower-right block is at least $(\lambda - \delta)Q$. Its inverse is therefore at most $(\lambda - \delta)^{-1}Q$, and its Schur complement is bounded below by the left side of (20). The Schur complement criterion proves the assertion. $\square$

Corollary 4.4 (Finite-projector completeness). *Under the final hypothesis of Theorem 4.3, assume moreover that A and the selected C_R are trace class, and choose R so that $\lambda_R < \mu$. If $P_N \uparrow I$ are finite-rank orthogonal projections, then the trace tails in Theorem 4.2 tend to zero and its finite LMI certifies a bound strictly below μ for all sufficiently large N .*

Proof. For the trace-class operators A and C_R , $\text{Tr}((I - P_N)A(I - P_N)) \rightarrow 0$ and likewise for C_R . Since $A - C_R$ is compact, $P_N(A - C_R)P_N \rightarrow A - C_R$ in norm. The residual in (22) therefore tends to zero, and the finite Schur test converges to the strict inequality $\lambda_R < \mu$. $\square$

5 The certified endpoint $a = 3/5$

5.1 The exact arithmetic cell

Set $a_0 = 3/5$. Rigorous interval evaluation gives

$$\log 3 < \frac{6}{5} < \log 4. \quad (27)$$

Thus the endpoint symbol is

$$b_{2,3}(t) = g_\infty(t) - \sqrt{2} \log 2 \cos(t \log 2) - \frac{2 \log 3}{\sqrt{3}} \cos(t \log 3). \quad (28)$$

Put $d = 1 - b_{2,3}$. In the notation of (18),

$$T_{a_0} = I - A + C_{\text{full}}, \quad A = K_{a_0, d_+}, \quad C_{\text{full}} = K_{a_0, (-d)_+}. \quad (29)$$

The retained rescue $C \preceq C_{\text{full}}$ is obtained by restricting $(-d)_+$ to the following union J of exact rational intervals:

$$\begin{aligned} J = & [60382193/20000000, 380082633/100000000] \\ & \cup [39235217/3125000, 1573243267/100000000] \\ & \cup [1908657941/100000000, 228037531/100000000] \\ & \cup [23057517/1000000, 1340114843/50000000] \\ & \cup [184766397/6250000, 3362720043/100000000] \\ & \cup [717875089/20000000, 4443581909/100000000]. \end{aligned} \quad (30)$$

Interval subdivision proves $d < 0$ throughout each component of J , so

$$C = K_{a_0, (-d)\mathbf{1}_J} \quad \text{satisfies} \quad 0 \preceq C \preceq C_{\text{full}}. \quad (31)$$

5.2 Complete sign and tail control

The derivative of the archimedean term has the positive expansion

$$g'_\infty(t) = \frac{t}{2} \sum_{k \geq 0} \frac{k + 1/4}{((k + 1/4)^2 + (t/2)^2)^2} > 0 \quad (t > 0). \quad (32)$$

Using

$$\sup_{y \geq 0} \frac{yu}{(u^2 + y^2)^2} = \frac{9}{16\sqrt{3}} \frac{1}{u^2}, \quad \sum_{k \geq 0} \frac{1}{(k + 1/4)^2} = \pi^2 + 8G,$$

All subtractions in (40) are one-sided: the full trace is rounded upward and the captured trace downward. The root charge (34) is added only in the adverse direction.

For

$$\eta = \frac{1}{16}, \quad \lambda = \frac{99997}{100000}, \quad \bar{\tau}_A = \frac{1}{200000}, \quad \bar{\tau}_C = \frac{1}{500000}, \quad (41)$$

define the Gram-coordinate matrix

$$\begin{aligned} \mathcal{M} = & \lambda G - (B_{A,0} + \varepsilon_{\text{root}} G - B_C) \\ & - \frac{(1 + \eta)\bar{\tau}_A(B_{A,0} + \varepsilon_{\text{root}} G) + (1 + \eta^{-1})\bar{\tau}_C B_C}{\lambda - \bar{\tau}_A}. \end{aligned} \quad (42)$$

An outward-rounded interval LDL^* factorization gives 44 strictly positive pivots, the smallest with lower endpoint greater than

$$0.00979334885665161. \quad (43)$$

Proposition 5.1 (Endpoint certificate). *On $\mathcal{Q}_{3/5}^0$,*

$$T_{3/5} \succeq \frac{3}{100000} I. \quad (44)$$

Proof. The sign proof gives $A \preceq A_0 + R_{\text{root}}$, while (31) gives $C \preceq C_{\text{full}}$. Thus

$$T_{3/5} \succeq I - (A - C).$$

Equations (40)–(43) verify the hypothesis of Theorem 4.2, so

$$A - C \preceq \frac{99997}{100000} I.$$

Subtracting this inequality from the identity proves (44). $\square$

6 Exact inward nesting

The passage from the endpoint to all smaller supports is exact and does not use numerical continuity in a .

Lemma 6.1 (Zero-extension identity). *Let $0 < a \leq a_0$, and let $f \in \mathcal{Q}_a^0$. If f^{ext} denotes its zero extension from $(-a, a)$ to $(-a_0, a_0)$, then*

$$\|f^{\text{ext}}\|_2 = \|f\|_2, \quad T_{a_0}[f^{\text{ext}}] = T_a[f]. \quad (45)$$

Proof. Zero extension preserves evenness, the $\cosh(x/2)$ -moment, the norm, and the Fourier transform. It remains to compare the arithmetic sums. If an atom n is active at a_0 but inactive at a , then $\log n \geq 2a$. Its contribution to the form is a constant multiple of the correlation of f with its translate by $\log n$. The two supports are disjoint, except possibly at a null endpoint when $\log n = 2a$, so this correlation is zero. All other terms agree. $\square$

Proof of Theorem 1.1. Apply Proposition 5.1 to f^{ext} with $a_0 = 3/5$, and use Lemma 6.1. $\square$

the ledger. The 60-digit interval balls and the ledger hash provide the data required for independent replay.

After cloning the repository and changing to its top-level directory, the certificate is reproduced on a POSIX shell by

```
python -m pip install -r requirements.txt
mkdir -p build
python certificate/certify.py \
  --workers 8 --precision-bits 256 --tolerance 1e-14 \
  --eval-limit 500000 --json build/fresh-ledger.json
python certificate/validate.py \
  --ledger build/fresh-ledger.json \
  --output build/fresh-schur-lmi.json --precision 256
```

Parallel worker scheduling affects elapsed time but not the canonical proof payload.

B Exact data flow for the continuum bound

For reference, the logical dependence of the computer-assisted part is

$$\begin{array}{c}
 \text{interval proof of (27), signs, and tail} \\
 \Downarrow \\
 A \preceq A_0 + \varepsilon_{\text{root}} I, \quad 0 \preceq C \preceq C_{\text{full}} \\
 \Downarrow \\
 \text{interval Gram/compression integrals and one-sided trace tails} \\
 \Downarrow \\
 \mathcal{M} \succ 0 \text{ by interval } LDL^* \\
 \Downarrow \\
 A - C \preceq (99997/100000)I \\
 \Downarrow \\
 T_{3/5} \succeq (3/100000)I \\
 \Downarrow \\
 T_a \succeq (3/100000)I \quad (0 < a \leq 3/5).
 \end{array}$$

Every arrow except the three interval-verification steps is proved symbolically in the body of the paper.

References

- [1] A. Weil, Sur les “formules explicites” de la théorie des nombres premiers, in *Comm. Sém. Math. Univ. Lund, Tome Supplémentaire*, 1952, pp. 252–265. <https://cds.cern.ch/record/471308>.
- [2] A. Weil, Sur les formules explicites de théorie des nombres, *Math. USSR-Izv.* **6** (1972), 1–17. <https://doi.org/10.1070/IM1972v006n01ABEH001866>.
- [3] H. Yoshida, On Hermitian forms attached to zeta functions, in *Zeta Functions in Geometry*, Adv. Stud. Pure Math. **21**, Kinokuniya, Tokyo, 1992, pp. 281–325. <https://doi.org/10.2969/aspm/02110281>.

