

PERMADOC CONCENTRATE

FOR USE ON RESIDENTIAL LAWNS, GARDENS, PATIOS AND HOME PICNIC AREAS

Controls Listed Pests for up to 4 Weeks • Water Based Formula • Kills Bugs on Your Lawns • Trees and Shrubs • Roses and Flowers Aids in Reducing Annovance from Mosquitoes and Other Flying Insects. • Kills Gnats, Fleas, Deer Ticks, Brown Dog Ticks and other Crawling Insects.

Permethrin [(3-phenoxyphenyl)methyl(±)-cis, trans-3-(2,2-dichloroetheynl)-2,2-dimethylcyclopropanecarboxylate] 2.50% Other Ingredients: 97.50% * Cis, trans ratio: Min. 35%(±) cis and max (±) trans.

KEEP OUT OF REACH OF CHILDREN CAUTION

PRECAUTIONARY STATEMENTS: HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION: May be harmful if swallowed or absorbed through skin. Avoid breathing spray mist. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling.

FIRST AID

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or

IF ON SKIN OR CLOTHING: Take on contamination doming. Taking skill minimization, the plant of the property of

ENVIRONMENTAL HAZARDS: This pesticide is extremely toxic to fish. Do not apply directly to water. Drifts and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters. This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively visiting the

EPA REG. NO. 10088-110-11547 • EPA EST. 11547-WI-1

NET CONTENTS: 20 GALLONS

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climactic modification and being grown in interior plantscapes, ornamental gardens or parks, or on golf courses or lawns

SHAKE WELL BEFORE USE.

This product helps control many annoying and troublesome outdoor insects. You can help protect your property by controlling the insects listed on this label. Do not spray animals. People and pets should not be allowed in treated areas until

This product can be applied using a hose-end sprayer, pump up pressure sprayer or by watering can. Carefully dilute the recommended amount of concentrate and water according to the directions found under the desired insect to control. Use a clean sprayer. Apply insecticide when insects first appear and repeat as necessary but not more frequently than every 3 days. Flush sprayer or watering can with clean water after each use.

Use on Lawns: Bent, Bermuda, Bluegrass, Dichondra, Fescue, Irish Moss, Merlon and St. Augustine.

Pests: Ants, Armyworms, Brown dog ticks, Chinch bugs, Crickets, Cutworms, Deer ticks, Fleas, Grasshoppers, Japanese beetles grubs, Mole crickets and Sod

Thoroughly wet down grass a few hours before applying. Home lawns should be no taller than 3 inches at time of application. Slightly more water may be used as long as the amount of product per area is as listed below. For heavy infestations, repeat application after 2 weeks. For ants, crickets and grasshoppers use 2-fl. oz. in 3 gallons of water to cover 1,000 sq. ft. For other listed insects, use 6 fl.oz. in 10 gallons of water to cover 1,000 sq. ft.

Use on Trees and Shrubs: Arizona Cypress, Azalea, Birch, Non-Bearing Citrus, Conifers, Elm, English Ivy, Euonymus, Weeping Fig, Fir, Honeysuckle, Lilac, Mock-Orange, Oak, Palm, Philodendron, Pine, Poinsettia, Tulip Poplar,

<u>Pests:</u> Aphids, Bagworms, Cicadas, Exposed thrips, Fall Cankerworms, Inchworms, Leafminers, Mealybugs, Spider mites, Whiteflies. Fall webworms on Cherry and Honeysuckle, Budworms on Conifer, Elm leaf beetles, Elm spanworms and Fall webworms on Elms. Tussock moths on Douglas and Grand Firs, Citrus black flies on Non-Bearing Citrus, Oakworms on Oaks, Pine beetles, Pine moths, Pine needleminers, and Needle scales on Pines.

Use 1 fl. oz. in 1 gallon of water. Apply as a spray, wetting leaves and branches thoroughly. Try to penetrate dense foliage. Spray in the late afternoon or

evening, when the temperature ranges from 50 to 75°F and when there is little or no wind. Spray at the first sign of insects. Repeat as necessary; use intervals of 4-8

Use on Roses and Flowers: Ageratum, Aster, Azalea, Begonia, Coleus, Common Ninebark and Snowberry, Exacum, Gladiolus, Gold Bells, Hypoestes, Ivy, Lilac, Marigold, Mock-Orange, Nannyberry, Orchid, Pansy, Pea Shrub, Petunia, Poinsettia, Rhododendron, Rose, Snapdragon, Zinnia.

Pests: Aphids, Bagworms, Cicadas, Exposed thrips, Fall Cankerworm, Fall webworms, Inchworms, Leafminers, Leafrollers, Mealybugs, Spider mites, and

Use 1 fl. oz. in 1 gallon of water. Apply as a spray, wetting the plants thoroughly. Try to hit the underside of leaves and penetrate dense foliage. Spray in the later afternoon or evening, when the temperature ranges from 50 to 75°F and when there is little or no wind. Spray at the first sign of insects. Repeat as necessary; use

Barrier Spray For Nuisance Pests: Such as Ants, Clover mites, Cluster flies, Crickets, Earwigs, Fleas, Flying moths, Gnats, House flies, Millipedes, Mosquitoes, Sowbugs (pillbugs), Spiders, Tent caterpillars, Ticks (Brown dog, Deer, Lone star).

GENERAL DIRECTIONS:

Use 6-2/3 fl.oz. in 1 quart of water, treating 400 linear feet with a 6-inch spray band.

Outside Surfaces of Buildings:
Pests: Ants, Clover mites, Crickets, Earwigs, Millipedes, Sowbugs (pillbugs). How to Use: To help prevent infestation of buildings, treat the building foundation to a height of 2 to 3 feet, where pests are active and may find entrance. Also, apply as a residual spray to outside surfaces of buildings including porches screens

window frames, eaves, patios, picnic areas, garages, refuse piles, and other areas where these pests congregate or have been seen. Repeat application may be needed, if re-infestation occurs.

Other Outside Areas:

Pests: Ants, Crickets, Earwigs, Fleas, Spiders, Ticks (Brown dog, Deer, Lone star). How to Use: For treatment of localized infestations of these insects in areas where there are weeds, shrubs or bushy non-crop areas, spray infested areas thoroughly For ants, thoroughly wet hills and runways. Repeat application may be needed, if re-

Pests: Cluster flies, Flying moths, Gnats, House flies, Mosquitoes.

How to Use: For use only as an aid in reducing annoyance from these insects. Spray outside surfaces of screens, doors, window fames or wherever these insect may enter the room. Also treat surfaces around light fixtures on porches, under patio

furniture, in garages and other places where these insects alight or congregate. Repeat application may be needed, if re-infestation occurs.

How to Use: Application should be made when caterpillars are young and tents are first noticed. For best results, apply in the late afternoon or evening, when caterpillars have returned to their tents. Wet tents with spray on all sides. A few caterpillars may be seen escaping from the tent, however, they should die within

Product with Hose End Sprayer Attached: Make sure knob is twisted fully clockwise to OFF position. Shake well before attaching to hose. Turn Product Control Button clockwise until its point aligns with triangle, then push all the way in. Avoid squeezing bottle. Turn water on at faucet, aim nozzle toward lawn and turn knob counter clockwise to begin spraying. When finished push Product Control Button to OFF position from the opposite side. Return knob and faucet to OFF position. Discharge water pressure by turning knob ON and OFF again. Turn Product Control Button counter clockwise away from triangle to prevent accidental discharge. Store out of reach of children.

STORAGE AND DISPOSAL

Do Not Contaminate Water, Food, or Feed By Storage or Disposal. Pesticide Storage: Store in a dry place no lower in temperature than 50°F or higher than 120°F

Pesticide Disposal: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container Disposal: Nonrefillable container. Do not reuse or refill container Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

0845•032410•

Sold By: Share Corporation • P.O. Box 245013 • Milwaukee, WI 53224 • (800) 776-7192 • www.sharecorp.com