



GENERAL CHEMISTRY

LAB COMPONENT CHE101L

GUIDED INQUIRY EXPERIMENTS

CONTENT: LAB 3

DISSOLUTION REACTIONS: HEATS OF DISSOCIATION

NAME

SECTION

STUDENT ID.....

DATE

TIME.....

NAME OF THE INSTRUCTOR


SIGNATURE & DATE

REPORT SUBMISSION DATE (ASSIGNED BY INSTRUCTOR)

EXPERIMENT 3

DISSOLUTION REACTIONS: HEATS OF DISSOCIATION

Heats (exothermic or endothermic) are associated with chemical reactions. Quantity of heat evolved or absorbed is directly proportional to the amount reacted. Consider the reaction:



Heat could be generated or absorbed in this reaction. When heat is generated/released from a chemical reaction it is called exothermic reaction (you can feel it by touching the reaction container (warmer) and when heat is absorbed the reaction is called endothermic (colder). When reactions occur in a reaction vessel (e.g., Beaker) in aqueous condition, formation and dissociation of chemical bonds occur simultaneously. Bond formation and dissociation involves heat energy of the system which is expressed by the term Q which is called enthalpy.

PROBLEM STATEMENT: *Is heat energy related to chemical reactions, how?*

This experiment is subdivided into two parts:

- I. QUALITATIVE & II. QUANTITATIVE

PART I. QUALITATIVE

DATA COLLECTION:

Place about 30 mL of distilled water into a 50 mL beaker. Suspend a thermometer (having 0.1°C division mark) into the beaker using thermometer clamp and ring stand. Please make sure that the thermometer is not touching the bottom of the beaker, as any movement of the beaker could break the thermometer. Record the temperature of water in the beaker in every 30 seconds for 240 seconds.




FIGURE 1: Experimental setup for dissolution reaction

Place a moderate amount (which would be 1 to 3 cm³) of supplied anhydrous magnesium sulfate (MgSO₄) to the beaker. Mix vigorously with the glass rod for 5 minutes. Record your observations. **(2 points)**

Repeat this procedure with each of the following compounds: **(2 points)**

- a. Sodium Nitrate, NaNO₃
- b. Sodium Chloride, NaCl
- c. Hydrated Calcium Chloride, CaCl₂·2H₂O
- d. Ammonium Nitrate, NH₄NO₃

DATA ANALYSIS:

What are the similarities and differences in the behavior of these compounds? Can you find out any generalization concerning all chemical reactions here? What conclusion can be drawn from these data? **(4 points)**

PART II. QUANTITATIVE

DATA COLLECTION:

- Accurately weigh a 3 to 5 gm sample of MgSO_4 on the analytical balance. Record the exact mass here. For 4 different trials below measure four different weight samples (e.g., 1, 2, 4 & 5 grams respectively).
- Suspend the thermometer into a polystyrene cup/coffee cup. Make sure of the thermometer is not touching the bottom of the cup. Measure 20 mL of distilled water by a volumetric cylinder into the cup and stir for 240 second. Record the temperature in every 20 seconds. After 240 seconds add MgSO_4 with vigorous mixing while continuing to record data for 5 minutes.
- Determine the temperature change, ΔT , for the reaction. This can be done from the difference of the highest temperature minus the slope of the line go through the points from first 240 seconds of data.
- Draw a temperature vs. time graph. Draw the best curve through the points and point out what is happening in each part of the curve.
- Repeat the whole procedure with NaNO_3

DATA TABLE 1:

TRAILS

(I) Mass of MgSO_4 <u>0.5 gm</u>			(II) Mass of MgSO_4 <u>1.0 gm</u>	
Time (s)	Temp($^{\circ}\text{C}$)		Time(s)	Temp($^{\circ}\text{C}$)
20			20	
40			40	
60			60	
80			80	
100			100	
120			120	
140			140	
160			160	
180			180	
200			200	
220			220	

240			240	
260			260	
280			280	
300			300	

(III) Mass of MgSO_4 <u>1.5 gm</u>			(IV) Mass of MgSO_4 <u>2.0 gm</u>	
Time (s)	Temp($^{\circ}\text{C}$)		Time(s)	Temp($^{\circ}\text{C}$)
20			20	
40			40	
60			60	
80			80	
100			100	
120			120	
140			140	
160			160	
180			180	
200			200	
220			220	
240			240	
260			260	
280			280	
300			300	

(I) Mass of NaNO_3 <u>0.5 gm</u>			(II) Mass of NaNO_3 <u>1.0 gm</u>	
Time (s)	Temp($^{\circ}\text{C}$)		Time(s)	Temp($^{\circ}\text{C}$)
20			20	
40			40	
60			60	
80			80	
100			100	
120			120	
140			140	
160			160	
180			180	
200			200	
220			220	
240			240	
260			260	
280			280	
300			300	

(III) Mass of NaNO ₃ <u>1.5 gm</u>			(IV) Mass of NaNO ₃ <u>2.0 gm</u>	
Time (s)	Temp(°C)		Time(s)	Temp(°C)
20			20	
40			40	
60			60	
80			80	
100			100	
120			120	
140			140	
160			160	
180			180	
200			200	
220			220	
240			240	
260			260	
280			280	
300			300	

DATA ANALYSIS

1. What do you understand form the data you recorded and from the other trials?(4 points)

2. Calculate the heat, Q & moles, n , of the reaction. Take help from the equation $Q = C \times M \times \Delta T$. Assume $C = 4.18 \text{ Joules/gram } ^\circ\text{C}$ and M is the mass of water (take the water density as 1.00 grams/cm^3). **(4 points)**

3. Plot the collected data as moles, n vs. Q . Number of moles can be calculated as $n = (\text{mass of sample in gram}) / (\text{molecular weight in grams/mole})$. Try to find an algebraic equation. **(4 points)**

FIGURE: plot here

MENTAL MODEL: Use the chemical equation given above to represent the dissolution reaction in this experiment. Draw a picture(s) which describes what is happening in atomic or in molecular level. How heat release or absorbed can be described from these pictures? **(5 points)**