

Class 11

Sharlene Yang (PID: A16351303)

Candy Project

In today's class we will examine 538 candy.

```
candy <- read.csv("candy-data.csv", row.names = 1)
head(candy)
```

	chocolate	fruity	caramel	peanutyalmondy	nougat	crispedricewafer
100 Grand	1	0	1	0	0	1
3 Musketeers	1	0	0	0	1	0
One dime	0	0	0	0	0	0
One quarter	0	0	0	0	0	0
Air Heads	0	1	0	0	0	0
Almond Joy	1	0	0	1	0	0

	hard	bar	pluribus	sugarpercent	pricepercent	winpercent
100 Grand	0	1	0	0.732	0.860	66.97173
3 Musketeers	0	1	0	0.604	0.511	67.60294
One dime	0	0	0	0.011	0.116	32.26109
One quarter	0	0	0	0.011	0.511	46.11650
Air Heads	0	0	0	0.906	0.511	52.34146
Almond Joy	0	1	0	0.465	0.767	50.34755

Q1. How many different candy types are in this dataset?

```
nrow(candy)
```

```
[1] 85
```

Q2. How many fruity candy types are in the dataset?

```
candy$fruity
```

```
[1] 0 0 0 0 1 0 0 0 0 1 0 1 1 1 1 1 1 1 0 1 1 0 0 0 0 1 0 0 1 1 1 0 0 1 0 0 0  
[39] 0 0 0 1 0 0 1 1 0 0 0 1 1 0 0 0 0 1 0 0 1 0 1 1 0 1 0 0 1 1 1 1 0 0 1 1 1 0  
[77] 0 0 1 0 1 1 1 0 0
```

```
sum(candy$fruity)
```

```
[1] 38
```

Q. What these fruity candy?

We can use the ==

```
rownames(candy [candy$fruity == 1,])
```

```
[1] "Air Heads" "Caramel Apple Pops"  
[3] "Chewey Lemonhead Fruit Mix" "Chiclets"  
[5] "Dots" "Dum Dums"  
[7] "Fruit Chews" "Fun Dip"  
[9] "Gobstopper" "Haribo Gold Bears"  
[11] "Haribo Sour Bears" "Haribo Twin Snakes"  
[13] "Jawbusters" "Laffy Taffy"  
[15] "Lemonhead" "Lifesavers big ring gummies"  
[17] "Mike & Ike" "Nerds"  
[19] "Nik L Nip" "Now & Later"  
[21] "Pop Rocks" "Red vines"  
[23] "Ring pop" "Runts"  
[25] "Skittles original" "Skittles wildberry"  
[27] "Smarties candy" "Sour Patch Kids"  
[29] "Sour Patch Tricksters" "Starburst"  
[31] "Strawberry bon bons" "Super Bubble"  
[33] "Swedish Fish" "Tootsie Pop"  
[35] "Trolli Sour Bites" "Twizzlers"  
[37] "Warheads" "Welch's Fruit Snacks"
```

How often does my favorite candy win

```
candy["Twix", ]$winpercent
```

```
[1] 81.64291
```

Q3. What is your favorite candy in the dataset and what is its winpercent value?

```
candy["Sour Patch Tricksters", ]$winpercent
```

```
[1] 52.82595
```

Q4. What is the winpercent value for “Kit Kat”?

```
candy ["Kit Kat",] $winpercent
```

```
[1] 76.7686
```

Q5. What is the winpercent value for “Tootsie Roll Snack Bars”?

```
candy["Tootsie Roll Snack Bars",] $winpercent
```

```
[1] 49.6535
```

```
library("skimr")  
skim(candy)
```

Table 1: Data summary

Name	candy
Number of rows	85
Number of columns	12
Column type frequency:	
numeric	12
Group variables	None

Variable type: numeric

skim_variable	n_missing	complete_rate	mean	sd	p0	p25	p50	p75	p100	hist
chocolate	0	1	0.44	0.50	0.00	0.00	0.00	1.00	1.00	
fruity	0	1	0.45	0.50	0.00	0.00	0.00	1.00	1.00	
caramel	0	1	0.16	0.37	0.00	0.00	0.00	0.00	1.00	
peanutyalmondy	0	1	0.16	0.37	0.00	0.00	0.00	0.00	1.00	
nougat	0	1	0.08	0.28	0.00	0.00	0.00	0.00	1.00	
crispedricewafer	0	1	0.08	0.28	0.00	0.00	0.00	0.00	1.00	
hard	0	1	0.18	0.38	0.00	0.00	0.00	0.00	1.00	
bar	0	1	0.25	0.43	0.00	0.00	0.00	0.00	1.00	
pluribus	0	1	0.52	0.50	0.00	0.00	1.00	1.00	1.00	
sugarpercent	0	1	0.48	0.28	0.01	0.22	0.47	0.73	0.99	
pricepercent	0	1	0.47	0.29	0.01	0.26	0.47	0.65	0.98	
winpercent	0	1	50.32	14.71	22.45	39.14	47.83	59.86	84.18	

Q6. Is there any variable/column that looks to be on a different scale to the majority of the other columns in the dataset?

Yes, the **winpercent** is on a 0:100 scale and all others appear to be 0:1 scale.


Q7. What do you think a zero and one represent for the `candy$chocolate` column?

A zero means that the candy is not classified as a chocolate candy.

Q8. Plot a histogram of **winpercent** values


```
library("ggplot2")

ggplot(candy) +
  aes(winpercent) +
  geom_histogram(bins= 10)
```



```
hist(candy$winpercent)
```

Histogram of candy\$winpercent



Q9. Is the distribution of winpercent values symmetrical?

No, they are not symmetrical.

Q10. Is the center of the distribution above or below 50%?

Below with the a mean:

```
mean(candy$winpercent)
```

```
[1] 50.31676
```

Q11. On average is chocolate candy higher or lower ranked than fruit candy?

To answer this question I will need to:

- “subset” (aka “select”, “filter”) the candy dataset to just chocolate candy
- get their winpercent values
- calculate the mean of these

Then do the same for fruity candy and compare.

```
#filter to just chocolate rows
chocolate.candy <- candy[as.logical(candy$chocolate),]

#get their winpercent values
chocolate.winpercent <- chocolate.candy$winpercent

#calculate their mean winpercent value
mean(chocolate.winpercent)
```

```
[1] 60.92153
```

```
fruity.candy <- candy[as.logical(candy$fruity),]

fruity.winpercent <- fruity.candy$winpercent

mean(fruity.winpercent)
```

```
[1] 44.11974
```

Chocolate candy is higher ranked than fruity candy.

Q12. Is this difference statistically significant?

```
t.test(chocolate.winpercent, fruity.winpercent)
```

Welch Two Sample t-test

```
data: chocolate.winpercent and fruity.winpercent
t = 6.2582, df = 68.882, p-value = 2.871e-08
alternative hypothesis: true difference in means is not equal to 0
95 percent confidence interval:
 11.44563 22.15795
sample estimates:
mean of x mean of y
 60.92153  44.11974
```

Overall Candy Rankings

There is a base R function called `sort()` for, guess what sorting vectors of input.

```
x <- c(5, 2, 10)

#sort(x, decreasing = TRUE)
sort(x)
```

```
[1] 2 5 10
```

The related function to `sort` that is often more useful is called `order()`. It returns the “indices” of the input that would result in it being sorted.

```
order(x)
```

```
[1] 2 1 3
```

```
x[order(x)]
```

```
[1] 2 5 10
```

Q13. What are the five least liked candy types in this set?

I can order by winpercent

```
ord <- order(candy$winpercent)

head(candy[ord,], 5)
```

	chocolate	fruity	caramel	peanut	almond	nougat
Nik L Nip	0	1	0		0	0
Boston Baked Beans	0	0	0		1	0
Chiclets	0	1	0		0	0
Super Bubble	0	1	0		0	0
Jawbusters	0	1	0		0	0

	crisped	rice	wafer	hard	bar	pluribus	sugar	percent	price	percent
Nik L Nip				0	0	0	1	0.197		0.976
Boston Baked Beans				0	0	0	1	0.313		0.511
Chiclets				0	0	0	1	0.046		0.325
Super Bubble				0	0	0	0	0.162		0.116
Jawbusters				0	1	0	1	0.093		0.511

	winpercent
Nik L Nip	22.44534
Boston Baked Beans	23.41782
Chiclets	24.52499
Super Bubble	27.30386
Jawbusters	28.12744

Q14. What are the top 5 all time favorite candy types out of this set?

```
new_ord <- order(candy$winpercent, decreasing = TRUE)

head(candy[new_ord, ], 5)
```

	chocolate	fruity	caramel	peanut	almond	nougat
Reese's Peanut Butter cup	1	0	0		1	0
Reese's Miniatures	1	0	0		1	0
Twix	1	0	1		0	0
Kit Kat	1	0	0		0	0
Snickers	1	0	1		1	1

	crisped	rice	wafer	hard	bar	pluribus	sugar	percent
Reese's Peanut Butter cup				0	0	0	0	0.720


Reese's Miniatures	0	0	0	0	0.034
Twix	1	0	1	0	0.546
Kit Kat	1	0	1	0	0.313
Snickers	0	0	1	0	0.546

	pricepercent	winpercent
Reese's Peanut Butter cup	0.651	84.18029
Reese's Miniatures	0.279	81.86626
Twix	0.906	81.64291
Kit Kat	0.511	76.76860
Snickers	0.651	76.67378

Q15. Make a first barplot of candy ranking based on winpercent values.

```
library(ggplot2)


ggplot(candy) +
  aes(winpercent, reorder(rownames(candy), winpercent)) +
  geom_col()
```



```
my_cols=rep("black", nrow(candy))
my_cols[as.logical(candy$chocolate)] = "chocolate"
my_cols[as.logical(candy$bar)] = "brown"
```

```
my_cols[as.logical(candy$fruity)] = "pink"
```

```
ggplot(candy) +
  aes(winpercent, reorder(rownames(candy), winpercent)) +
  geom_col(fill= my_cols)
```



Q17. What is the worst ranked chocolate candy?

Sixlets

Q18. What is the best ranked fruity candy?


Starburst

Taking a look at pricepercent

Q. What is the best candy for the least money?


```
my_cols[as.logical(candy$fruity)] = "red"
```

```
ggplot(candy) +
  aes(winpercent, pricepercent) +
  geom_point(col = my_cols)
```



Add some labels

```
ggplot(candy) +
  aes(winpercent, pricepercent, label = rownames(candy)) +
  geom_point(col = my_cols)+
  geom_text()
```




To deal with overlapping labels I can use the `geom_repel` package.

```
library(ggrepel)

ggplot(candy) +
  aes(winpercent, pricepercent, label = rownames(candy)) +
  geom_point(col = my_cols) +
  geom_text_repel(col=my_cols, size=3.3, max.overlaps = 5)
```

Warning: ggrepel: 65 unlabeled data points (too many overlaps). Consider increasing max.overlaps




Exploring the correlation structure

Pearson correlation goes between -1 and +1 with zero indicating no correlations and values close to one being very highly (ani) correlated.

```
library(corrplot)
```

```
corrplot 0.92 loaded
```

```
cij <- cor(candy)
corrplot(cij)
```



Q22. Examining this plot what two variables are anti-correlated (i.e. have minus values)?

Chocolate and fruit are anti-correlated

Q23. Similarly, what two variables are most positively correlated?

Chocolate and winpercent or chocolate and bar

Principal Component Analysis

The base R function for PCA is called `prcomp()` and we can set “scale= TRUE/FALSE”.

```
pca <- prcomp(candy, scale = TRUE)
summary(pca)
```

Importance of components:

	PC1	PC2	PC3	PC4	PC5	PC6	PC7
Standard deviation	2.0788	1.1378	1.1092	1.07533	0.9518	0.81923	0.81530
Proportion of Variance	0.3601	0.1079	0.1025	0.09636	0.0755	0.05593	0.05539
Cumulative Proportion	0.3601	0.4680	0.5705	0.66688	0.7424	0.79830	0.85369

	PC8	PC9	PC10	PC11	PC12
Standard deviation	0.7811	0.77556	0.77455	0.77455	0.77455
Proportion of Variance	0.05539	0.05539	0.05539	0.05539	0.05539
Cumulative Proportion	0.90908	0.96447	1.00000	1.00000	1.00000


Standard deviation	0.74530	0.67824	0.62349	0.43974	0.39760
Proportion of Variance	0.04629	0.03833	0.03239	0.01611	0.01317
Cumulative Proportion	0.89998	0.93832	0.97071	0.98683	1.00000

The main result of PCA- i.e. the new PC plot (projection of candy on out new PC axis) is contained in `pca$x`

```
pc <- as.data.frame(pca$x)

ggplot(pc) +
  aes(PC1, PC2, label = rownames(pc)) +
  geom_point(col = my_cols) +
  geom_text_repel(max.overlaps = 5)
```

Warning: ggrepel: 71 unlabeled data points (too many overlaps). Consider increasing max.overlaps



Q24. What original variables are picked up strongly by PC1 in the positive direction? Do these make sense to you?

Fruity and pluribus are picked up strongly by PC1 in the positive direction.