

engn2770_computational_tutorial

September 16, 2020

1 ENGN 2770: Atomistic Reaction Engineering

1.1 Computational Tutorial

This is the pdf version. The interactive version (which includes necessary environment files) can be accessed from using the link:

https://gesis.mybinder.org/binder/v2/gh/sharma1908shubham/engn2770_computational_tutorial/38dfa226c5862fa9dfa16db6a58bdc1e851a7773

(Note: This can take a minute or two to load and download required environment files)

Once you open the link, you can download the “engn2770_computational_tutorial.ipynb” and save it on your local computer.

Note, you will need jupyter notebook along with other necessary packages if you wish to use it on your local machine(ASE, python, matplotlib, numpy).

1.2 A) Intro to linux bash commands (<https://files.fosswire.com/2007/08/fwunixref.pdf>)

```
[3]: from IPython.display import Image
```

```
[4]: Image(filename='extras/linux_tutorial_page1.png')
```

```
[4]:
```

Job	Command
identify the directory you are in	pwd
see the files in the directory	ls
see the files in the directory with hidden files	ls -a
go to your home directory	cd (or, cd ~/)
change directory to <directory_name>	cd <directory_name>
move one directory up	cd ..
move two directories up	cd ../../
make directory <directory_name>	mkdir <directory_name>
copy a file	cp </path/to/file/file_name> </new/path/to/file/new_file_name>
remove a file / remove a file recursively (careful -- no way to revert if you delete a file/directory)	rm <file_name> / rm -r <file_name>
create an empty file	touch <file_name>

[5]: `Image(filename='extras/linux_tutorial_page2.png')`

[5]:

Job	Command
move a file	mv </path/to/file/file_name> </new/path/to/file/new_file_name>
display the contents of the file	cat <file_name>
copy file from your local computer to CCV	scp -r </path/to/location/on/computer/file_name> <username@ssh.ccv.brown.edu:/path/to/new/location/file_name>
text editors	vim/ vi/ nano/ gedit

Environment Setup:

```
$ cd  
(vi/nano/gedit) ~/.bashrc
```

Inside your .bashrc, type

```
$ source /gpfs/data/ap31/ap31/teach/2020-engn2770/settings/engn2770packages
```

Save your .bashrc, and type

```
$ source ~/.bashrc (This needs to be done only once - i.e whenever you change your ~/.bashrc)
```

```
$ engn2770_loadase (This needs to be loaded every time you will be using ASE)
```

```
$ python3
```

```
$ import ase
```

```
$ ase. file
```

(The output should point to '/users/ap31/data/ap31/teach/2020-engn2770/packages/ase/ase/__init__.py')

Note: The first line in your python script should be always be `#!/usr/bin/env python3`

1.3 B) Python

1.3.1 Basic arithmetic

```
[6]: # Basic arithmetic
a = 2.
b = 3.

print(a + b)
```

5.0

```
[7]: # square of a number
sq = a**2
print(sq)
```

4.0

```
[8]: # division & %
print(4./2)

# remainder of a division
print(15.%8)
print(8.%2) # If this is 0, then 8 is divisible by 2!
```

2.0

7.0

0.0

1.3.2 Functions

```
[9]: # functions

# define add function
def add(a,b):
    return a+b

# define power function
def power(a,pow=2):
    return a**pow

# call add function
add_result = add(2.,3.)
print(add_result)

# call power function
power_result = power(5.,3)
print(power_result)
```

5.0
125.0

```
[10]: sum_result = sum((2.,2.))  
      print(sum_result)  
  
      pow_result = pow(5.,4)  
      print(pow_result)
```

4.0
625.0

1.3.3 numpy, list & arrays

```
[11]: # using numpy (Numerical Python)  
      import numpy as np  
  
      sum_result = np.sum((2.,-3.))  
      print(sum_result)  
  
      power_result = np.power(2.,8)  
      print(power_result)
```

-1.0
256.0

```
[12]: # list  
      things = ['a',7]  
  
      for thing in things:  
          print(thing)
```

a
7

```
[13]: # Basic list commands  
  
      _list = [1,2,3,4]  
  
      #_list.append(5)  
      #_list.pop(2)  
      #_list.reverse()  
  
      print(_list)
```

[1, 2, 3, 4]

```
[14]: import numpy as np

# return evenly spaced numbers over a specified interval

a = np.linspace(1,12,12)
print(a)
print(a.shape)

# convert a 1D array into 2D
a.reshape(3,4)
```

```
[ 1.  2.  3.  4.  5.  6.  7.  8.  9. 10. 11. 12.]
(12,)
```

```
[14]: array([[ 1.,  2.,  3.,  4.],
           [ 5.,  6.,  7.,  8.],
           [ 9., 10., 11., 12.]])
```

1.3.4 For loop & conditional statements

```
[15]: # for loops
for element in range(5):
    print(element, element**2)
```

```
0 0
1 1
2 4
3 9
4 16
```

```
[16]: # For loops with conditional statements

dice = [1,2,3,4,5,6]

type_list = []

for roll in dice:
    if roll % 2 == 0:
        type_list.append('even')
    else:
        type_list.append('odd')

print(type_list)
```

```
['odd', 'even', 'odd', 'even', 'odd', 'even']
```

```
[17]: num = np.linspace(0,10,11)
print(num)
```

```
doubled_list = []

for element in num:
    if element > 5:
        doubled_list.append(element*2)

print(doubled_list)
```

```
[ 0.  1.  2.  3.  4.  5.  6.  7.  8.  9. 10.]
[12.0, 14.0, 16.0, 18.0, 20.0]
```

[18]: *# For loop with conditional statement in one line*

```
num = np.linspace(0,10,11)
print(num)

doubled = [element * 2 for element in num if element>5]
print(doubled)
```

```
[ 0.  1.  2.  3.  4.  5.  6.  7.  8.  9. 10.]
[12.0, 14.0, 16.0, 18.0, 20.0]
```

[19]: `long_words = ['platinum', 'hi', 'carbon', 'welcome', 'hydrogen', 'neptune', 'covid', 'python']`

```
# checks and stores into new list based on length of a word
short_words = [word for word in long_words if len(word) < 6]

short_words
```

[19]: ['hi', 'covid']

1.4 C) Atomic Simulation Environment

[20]: *# Some basic packages we will need throughout*

```
import os
import numpy as np

import ase
from ase.parallel import paropen
from ase.io import read, write
from ase.visualize import view

# Plotting and image visualizing in Jupyter Notebook
import matplotlib
import matplotlib.pyplot as plt
from IPython.display import Image

%matplotlib inline
```

2 Module 1: Atom / Atoms

```
[21]: # import the module (use tab tab for auto-completion)
      from ase import Atom, Atoms
```

```
[22]: # use ? to know more
      #atom = Atom?
```

```
[23]: atom = Atom('O')
      atom.mass = 16
      atom.charge = -2
      atom.position = (0,2,3)
      #atom.symbol
      #atom.x
      #atom.y
```

```
[24]: # use ? to know more
      #Atoms?
```

```
[25]: d = 1.104 # N2 bondlength

      # The following three are equivalent
      N2 = Atoms('N2', [(0, 0, 0), (0, 0, d)])
      N2 = Atoms(numbers=[7, 7], positions=[(0, 0, 0), (0, 0, d)])
      N2 = Atoms([Atom('N', (0, 0, 0)), Atom('N', (0, 0, d))])

      # to save and view a snapshot
      write('extras/n2.png', N2, rotation='-80y')
      Image(filename='extras/n2.png')

      # to have a 3D view in notebook
      #view(N2, viewer='x3d')
```

[25]:



```
[26]: d = 2.9
      L = 10.0

      # construct a Au wire
      wire = Atoms('Au',
                  positions=[[0, L / 2., L / 2.]],
                  cell=[d, L, L],
                  pbc=[1, 0, 0])


      wire.get_chemical_symbols()

      # to save and view a snapshot
      write('extras/wire.png', wire * (6,1,1))
      Image(filename='extras/wire.png')
```

```
#view(wire*(6,1,1), viewer='x3d')

#view(wire*(6,1,1))
```

[26]:



3 Module 2: Build (molecule, bulk, surfaces)

```
[27]: from ase.build import molecule

m = molecule('CH3CH2OH')
print(m.get_chemical_symbols())
print(m.get_chemical_formula())
print(m.get_positions())

# to save and view a snapshot
write('extras/ethanol.png', m, rotation='-20x')
Image(filename='extras/ethanol.png',width=150)

#view(m, viewer='x3d')

#view(m)
```

```
['C', 'C', 'O', 'H', 'H', 'H', 'H', 'H', 'H']
C2H6O
[[ 1.168181 -0.400382  0.        ]
 [ 0.        0.559462  0.        ]
 [-1.190083 -0.227669  0.        ]
 [-1.946623  0.381525  0.        ]
 [ 0.042557  1.207508  0.886933]
 [ 0.042557  1.207508 -0.886933]
 [ 2.115891  0.1448    0.        ]
 [ 1.128599 -1.037234  0.885881]
 [ 1.128599 -1.037234 -0.885881]]
```


[27]:



4 Optimize lattice constant

```
[28]: from ase.build import bulk
from ase.calculators.emt import EMT
from ase.io import Trajectory

a0 = 3.6
cu = bulk('Cu', 'fcc', a=a0, cubic=True)

print(cu.get_cell())


# to save and view a snapshot
write('extras/cu.png', cu)
Image(filename='extras/cu.png',width=150)

#view(cu, viewer='x3d')

#view(cu)
```

Cell([3.6, 3.6, 3.6])

[28]:



```
[29]: # create an empty list to store energy
energy_list = []

# start with a good guess
# and create a list with fluctuations in a.

a0 = 3.6
a_list = a0 + np.linspace(-0.2, 0.2, 101)

# For loop to use each and every value of a
# and calculate & store the potential energy
```

```

for a in a_list:
    cu = bulk('Cu', 'fcc', a=a, cubic=True)
    cu.set_calculator(EMT())
    e = cu.get_potential_energy()
    energy_list.append(e)

```

```


[30]: # plotting using matplotlib (https://matplotlib.org/)

```

```

fig, ax = plt.subplots(figsize=(7,7))
plt.scatter(a_list, energy_list, marker='o', s=10)
plt.xlabel('$a$, $\mathrm{\AA}$', fontsize=15)
plt.xticks(fontsize=15)
plt.ylabel('potential energy, eV', fontsize=15)
plt.yticks(fontsize=15)
fig.savefig('extras/lattice_constant.png')
plt.show()

```



```
[31]: # index corresponding to the lowest energy

index = energy_list.index(min(energy_list))
lc = a_list[index]
print('lattice constant:%0.3f'%lc)
```

lattice constant:3.588

5 Module 3: Surface, Constraints

```
[32]: from ase.build import fcc111
      from ase.constraints import FixAtoms
```

```
[33]: #fcc111?
```

```
[34]: # Create a slab using build
slab = fcc111('Cu', size=(3,3,3), a=3.588, vacuum=10)
slab.set_pbc((1,1,0))


# Use of constraints, here, Fix the bottom most layer
indices=[atom.index for atom in slab if atom.tag == 3]
constraint = FixAtoms(indices)
slab.set_constraint(constraint)

# Note that the constraint is not visible through
# the following method

write('extras/cu_slab.png', slab, rotation='10z,-80x')
Image(filename='extras/cu_slab.png')

#view(slab, viewer='x3d')
#view(slab)
```

```
[34]:
```



6 Module 4: Add an adsorbate

method 1

```
[35]: from ase.build import add_adsorbate


# create a slab
slab = fcc111('Cu', size=(3,3,3), a=3.588, vacuum=10)
slab.set_pbc((1,1,0))

# create an atom and use add_adsorbate to add it to the slab
adsorbate = Atom('O')
add_adsorbate(slab, adsorbate, 1.8, 'ontop')

write('extras/slab_with_adsorbate.png', slab * (2, 2, 1))
Image(filename='extras/slab_with_adsorbate.png')
```

```
#view(slab, viewer='x3d')
#view(slab)
```

[35]:



method 2

[36]:

```
a = 3.558

# create a slab
slab = fcc111('Cu', size=(3,3,3), a=3.588, vacuum=10)
slab.set_pbc((1,1,0))


# use the slab atom position to add it to the slab
adsorbate = Atom('O')
adsorbate.position = slab[18].position + (0,0,2)

slab_with_add = slab + adsorbate

write('extras/slab_with_adsorbate.png', slab_with_add)
Image(filename='extras/slab_with_adsorbate.png')

#view(slab_with_add, viewer='x3d')
#view(slab_with_add)
```

[36]:



7 Module 5: Structure optimization using EMT

```
[37]: from ase.constraints import FixAtoms
      from ase.build import fcc111, add_adsorbate
      from ase.calculators.emt import EMT
      from ase.optimize import BFGS

      # Construct a slab with optimized lattice constant
      slab = fcc111('Cu', size=(3,3,3), a=3.588, vacuum=10)
      # Periodic boundary conditions
      slab.set_pbc((1,1,0))

      # Construct & add an adsorbate
      adsorbate = Atom('O')
      add_adsorbate(slab, adsorbate, 1.8, 'ontop')

      # Fix atoms
      indices=[atom.index for atom in slab if atom.tag == 3]
      constraint = FixAtoms(indices)
      slab.set_constraint(constraint)

      # Calculator
      calc = EMT()
      slab.set_calculator(calc)

      if os.path.exists('qn.log'):
          os.remove('qn.log')

      # Optimization
      opt = BFGS(slab, logfile='qn.log', trajectory='qn.traj')
      opt.run(fmax=0.01)
```

[37]: True

```
[38]: #relaxed_slab = read('qn.traj')
      #view(relaxed_slab)
```

```
[39]: myfile = open("qn.log", 'r')
      txt = myfile.read()
      print(txt)
```

	Step	Time	Energy	fmax
BFGS:	0	17:33:42	6.520552	0.9096
BFGS:	1	17:33:42	6.505566	0.4225
BFGS:	2	17:33:42	6.501307	0.0559
BFGS:	3	17:33:42	6.501107	0.0437
BFGS:	4	17:33:42	6.500731	0.0845
BFGS:	5	17:33:42	6.500402	0.0943
BFGS:	6	17:33:43	6.500170	0.0693
BFGS:	7	17:33:43	6.500026	0.0385
BFGS:	8	17:33:43	6.499856	0.0668
BFGS:	9	17:33:43	6.499623	0.0791
BFGS:	10	17:33:43	6.499416	0.0562
BFGS:	11	17:33:43	6.499321	0.0183
BFGS:	12	17:33:43	6.499287	0.0128
BFGS:	13	17:33:43	6.499261	0.0155
BFGS:	14	17:33:43	6.499236	0.0165
BFGS:	15	17:33:44	6.499223	0.0082

```
[40]: from ase.constraints import FixAtoms
      from ase.build import fcc111, add_adsorbate
      from ase.calculators.emt import EMT
      from ase.optimize import BFGS

      # Construct a slab with optimized lattice constant
      slab = fcc111('Cu', size=(3,3,3), a=3.588, vacuum=10)
      # periodic boundary condition
      slab.set_pbc((1,1,0))

      # Construct & add an adsorbate
      adsorbate = Atom('O')
      add_adsorbate(slab, adsorbate, 1.8, 'hcp')

      # Fix atoms constraint to fix the last atomic layer
      indices=[atom.index for atom in slab if atom.tag == 3]
      constraint = FixAtoms(indices)
      slab.set_constraint(constraint)

      # Calculator
      calc = EMT()
      slab.set_calculator(calc)

      if os.path.exists('qn.log'):
          os.remove('qn.log')

      # Optimization
      opt = BFGS(slab, logfile='qn.log', trajectory='qn.traj')
      opt.run(fmax=0.01)
```

[40]: True

```
[41]: myfile = open("qn.log", 'r')
      txt = myfile.read()
      print(txt)
```

	Step	Time	Energy	fmax
BFGS:	0	17:33:44	6.570523	1.7805
BFGS:	1	17:33:44	6.508532	1.5692
BFGS:	2	17:33:44	6.388970	0.6331
BFGS:	3	17:33:44	6.368169	0.3576
BFGS:	4	17:33:44	6.363440	0.2807
BFGS:	5	17:33:44	6.346981	0.1420
BFGS:	6	17:33:45	6.345193	0.1026
BFGS:	7	17:33:45	6.343991	0.0898
BFGS:	8	17:33:45	6.342511	0.0919
BFGS:	9	17:33:45	6.340466	0.0910
BFGS:	10	17:33:45	6.339097	0.1304
BFGS:	11	17:33:45	6.338357	0.1316
BFGS:	12	17:33:45	6.337813	0.1097
BFGS:	13	17:33:46	6.337098	0.0630
BFGS:	14	17:33:46	6.336492	0.0408
BFGS:	15	17:33:46	6.336258	0.0134
BFGS:	16	17:33:46	6.336218	0.0105
BFGS:	17	17:33:46	6.336205	0.0090

8 Module 6: Calculate Barriers using nudged elastic band (NEB)

```
[42]: from ase.neb import NEB
      from ase.optimize import BFGS
      from ase.calculators.emt import EMT

      # Create a bare slab
      slab = fcc111('Cu', size=(3,3,3), a=3.588, vacuum=10)
      slab.set_pbc((1,1,0))

      indices=[atom.index for atom in slab if atom.tag == 3]
      constraint = FixAtoms(indices)
      slab.set_constraint(constraint)

      slab.set_calculator(EMT())

      opt = BFGS(slab, logfile='bare_slab.log', trajectory='bare_slab.traj')
      opt.run(fmax=0.01)

      # create an adsorbate
      adsorbate = Atom('O')

      #####
```



```

##### SLAB 1: Cu with O in fcc (position 1) #####
#####

initial = slab + adsorbate
initial[-1].position = initial[18].position + (1.3,0.8,1)

initial.set_calculator(EMT())
opt = BFGS(initial, logfile='initial.log', trajectory='initial.traj')
opt.run(fmax=0.01)

#####
##### SLAB 1: Cu with O in fcc (position 2) #####
#####

final = slab + adsorbate
final[-1].position = final[19].position + (1.3, 0.8, 1)

final.set_calculator(EMT())
opt = BFGS(final, logfile='final.log', trajectory='final.traj')
opt.run(fmax=0.01)

n = 10

images = [initial]
for i in range(n):
    image = initial.copy()
    image.set_calculator(EMT())
    image.get_potential_energy()
    image.get_forces()
    images.append(image)

images.append(final)

neb = NEB(images)
neb.interpolate()

qn = BFGS(neb, logfile='neb.log', trajectory='neb.traj')
qn.run(fmax=0.05)

```

[42]: True