

MA2C03: ASSIGNMENT 1
DUE BY FRIDAY, NOVEMBER 17
AT LECTURE OR IN THE MATHS OFFICE ROOM 0.6

Please write down clearly both your name and your student ID number on everything you hand in. Please attach a cover sheet with a declaration confirming that you know and understand College rules on plagiarism. Details can be found on <http://tcd-ie.libguides.com/plagiarism/declaration>.

1) For any sets A and B , define $A \Delta B$, the **symmetric difference** of A and B to be the set $(A - B) \cup (B - A)$. Prove that intersection \cap distributes over Δ : $A \cap (B \Delta C) = (A \cap B) \Delta (A \cap C)$ for all sets A , B , and C using the proof methods employed in lecture. Venn diagrams, truth tables, or diagrams for simplifying statements in Boolean algebra such as Veitch diagrams are **NOT** acceptable and will not be awarded any credit.

2) Let \mathbb{R} be the set of real numbers. For $x, y \in \mathbb{R}$, $x \sim y$ iff $x - y \in \mathbb{Q}$, i.e., if the difference $x - y$ is a rational number. Determine:

- (i) Whether or not the relation \sim is *reflexive*;
- (ii) Whether or not the relation \sim is *symmetric*;
- (iii) Whether or not the relation \sim is *anti-symmetric*;
- (iv) Whether or not the relation \sim is *transitive*;
- (v) Whether or not the relation \sim is an *equivalence relation*;
- (vi) Whether or not the relation \sim is a *partial order*.

Justify your answers.

3) Let A be a set, and let $\mathcal{A} = \{A_\alpha \mid \alpha \in I\}$, where I is an indexing set, be any partition of the set A . Define a relation R on A as follows: $x, y \in A$ satisfy xRy iff $x, y \in A_\alpha$ for some $\alpha \in I$. In other words, xRy iff x and y belong to the same set of the partition. Prove that R is an equivalence relation and that the partition R defines on A is precisely the given partition \mathcal{A} . (Hint: Recall we discussed in lecture the one-to-one correspondence between partitions and equivalence relations, and this is the proof direction I sketched in lecture without providing the details.)

4) Where is the fallacy in the following argument by induction? Justify your answer.

Statement: For every non-negative integer k , $2 \times k = 0$.

“Proof:” We give a proof using strong induction on k . Denote by $P(k)$ the statement “if k is non-negative integer, then $2 \times k = 0$.”

Base case: Show $P(0)$. Obviously, $2 \times 0 = 0$.

Inductive step: Assume $P(n)$ is true for every n such that $0 \leq n \leq k$ (the strong induction hypothesis). We have to show that $P(k+1)$ also holds. We write $k+1 = i+j$, where i and j are non-negative integers. By the inductive hypothesis,

$$2(k+1) = 2(i+j) = 2i + 2j = 0 + 0 = 0.$$

Therefore, by induction, $P(k)$ is true for all $k \in \mathbb{N}$, so $2 \times 1 = 0$.

5) Use mathematical induction to prove that

$$1^3 + 2^3 + \cdots + n^3 = (1 + 2 + \cdots + n)^2.$$

(Hint: Recall we proved in lecture that $1 + 2 + \cdots + n = \frac{n(n+1)}{2}$.)

6) Let $f : [-2, 0] \rightarrow [0, 1]$ be the function defined by $f(x) = \frac{1}{x^2 + 6x + 9}$ for all $x \in [-2, 0]$. Determine whether or not this function is injective and whether or not it is surjective. Justify your answers. Recall that $[-2, 0]$ is the set of all real numbers between -2 and 0 with the endpoints of -2 and 0 included in the set.

7) Let $A = \{(x, y) \in \mathbb{R}^2 \mid 2x - 3y = 0\}$ with the operation of addition given by $(x_1, y_1) + (x_2, y_2) = (x_1 + x_2, y_1 + y_2)$.

- (a) Is $(A, +)$ a semigroup? Justify your answer.
- (b) Is $(A, +)$ a monoid? Justify your answer.
- (c) Is $(A, +)$ a group? Justify your answer.