


Fig. 10 | Time-resolved signed VDFs
Latest corrected spectra / D4_s0 · Focused velocity view, full grids retained in NPZ


Every saved time row; time increases downward; native velocity cells; shared signed linear color scales. Focus changes axis limits only; see full-range companion.
Input correction: (a) subtract 1691; (b,c) marked pixels filled from nearest original valid value in the same row.
Signed VDF values retained without smoothing, clipping or renormalization; no VDF ground truth.


Fig. 10 | Signed VDFs at representative times
Latest corrected spectra / D4_s0 · Focused velocity view, full grids retained in NPZ

























Nearest saved rows to 2, 3 and 4 ns; actual times and source row indices in legends. Focus changes axis limits only; see full-range companion.
Input correction: (a) subtract 1691; (b,c) marked pixels filled from nearest original valid value in the same row.
Signed VDF values retained without smoothing, clipping or renormalization; no VDF ground truth.

Fig. 10 | Time-resolved signed VDFs
Latest corrected spectra / D4_s0 · Focused velocity view, full grids retained in NPZ




Every saved time row; time increases downward; native velocity cells; shared signed linear color scales. Focus changes axis limits only; see full-range companion.
Input correction: (a) subtract 1691; (b,c) marked pixels filled from nearest original valid value in the same row.
Signed VDF values retained without smoothing, clipping or renormalization; no VDF ground truth.

Fig. 10 | Signed VDFs at representative times
Latest corrected spectra / D4_s0 · Focused velocity view, full grids retained in NPZ




Nearest saved rows to 2, 3 and 4 ns; actual times and source row indices in legends. Focus changes axis limits only; see full-range companion.
Input correction: (a) subtract 1691; (b,c) marked pixels filled from nearest original valid value in the same row.
Signed VDF values retained without smoothing, clipping or renormalization; no VDF ground truth.

Fig. 10 | Time-resolved signed VDFs
Latest corrected spectra / D4_s0 · Focused velocity view, full grids retained in NPZ




Every saved time row; time increases downward; native velocity cells; shared signed linear color scales. Focus changes axis limits only; see full-range companion.
Input correction: (a) subtract 1691; (b,c) marked pixels filled from nearest original valid value in the same row.
Signed VDF values retained without smoothing, clipping or renormalization; no VDF ground truth.

Fig. 10 | Signed VDFs at representative times
Latest corrected spectra / D4_s0 · Focused velocity view, full grids retained in NPZ




Nearest saved rows to 2, 3 and 4 ns; actual times and source row indices in legends. Focus changes axis limits only; see full-range companion.
Input correction: (a) subtract 1691; (b,c) marked pixels filled from nearest original valid value in the same row.
Signed VDF values retained without smoothing, clipping or renormalization; no VDF ground truth.

Fig. 10 | Time-resolved signed VDFs
Latest corrected spectra / D4_s0 · Focused velocity view, full grids retained in NPZ




Every saved time row; time increases downward; native velocity cells; shared signed linear color scales. Focus changes axis limits only; see full-range companion.
Input correction: (a) subtract 1691; (b,c) marked pixels filled from nearest original valid value in the same row.
Signed VDF values retained without smoothing, clipping or renormalization; no VDF ground truth.

Fig. 10 | Signed VDFs at representative times
Latest corrected spectra / D4_s0 · Focused velocity view, full grids retained in NPZ




Nearest saved rows to 2, 3 and 4 ns; actual times and source row indices in legends. Focus changes axis limits only; see full-range companion.
Input correction: (a) subtract 1691; (b,c) marked pixels filled from nearest original valid value in the same row.
Signed VDF values retained without smoothing, clipping or renormalization; no VDF ground truth.

Fig. 10 | Time-resolved signed VDFs
Latest corrected spectra / D4_s0 · Full velocity range



Every saved time row; time increases downward; native velocity cells; shared signed linear color scales.
Input correction: (a) subtract 1691; (b,c) marked pixels filled from nearest original valid value in the same row.
Signed VDF values retained without smoothing, clipping or renormalization; no VDF ground truth.

Fig. 10 | Signed VDFs at representative times
Latest corrected spectra / D4_s0 · Full velocity range



Nearest saved rows to 2, 3 and 4 ns; actual times and source row indices in legends.
Input correction: (a) subtract 1691; (b,c) marked pixels filled from nearest original valid value in the same row.
Signed VDF values retained without smoothing, clipping or renormalization; no VDF ground truth.