

# Norm 2 optimization under unit vectors

Sharon Rotgaizer

April 27, 2024

It is relatively easy to optimize convex functions without any constraints, but in CS, we need many times to solve problems using the constraint of  $\|x\| = 1$ . We will see how to solve two fundametal problems.

## 1 simple case

Let  $x \in \mathbb{R}^2$ ,  $A \in \mathbb{R}^{n \times 2}$ ,  $b \in \mathbb{R}^n$  and  $f(x) = \|Ax - b\|_2^2$ . Our problem is:

$$\min_{\|x\|=1} f(x) = \min_{\|x\|=1} \|Ax - b\|_2^2 = \min_{\|x\|=1} \sum_{i=1}^n (A_{i1}x_1 + A_{i2}x_2 - b_i)^2$$

Using polar representation, we get rid of  $\|x\| = 1$  by choosing  $x_1 \leftarrow \cos(x)$ ,  $x_2 \leftarrow \sin(x)$ .

$$\min_{\|x\|=1} f(x) = \min_x \sum_{i=1}^n (A_{i1} \cos(x) + A_{i2} \sin(x) - b_i)^2$$

Lets derive!

$$f'(x) = 2 \sum_{i=1}^n (A_{i1} \cos(x) + A_{i2} \sin(x) - b_i)(A_{i2} \cos(x) - A_{i1} \sin(x)) = 0$$

For each inner expression:

$$\begin{aligned} E &= (A_{i1} \cos(x) + A_{i2} \sin(x) - b_i)(A_{i2} \cos(x) - A_{i1} \sin(x)) \\ &= A_{i1}A_{i2} \cos(2x) + 0.5(A_{i2}^2 - A_{i1}^2) \sin(2x) - A_{i2}b_i \cos(x) - A_{i1}b_i \sin(x) \end{aligned}$$

And totally:

$$f'(x) = \cos(2x) \sum_{i=1}^n A_{i1}A_{i2} + 0.5 \sin(2x) \sum_{i=1}^n (A_{i2}^2 - A_{i1}^2) - \cos(x) \sum_{i=1}^n A_{i2}b_i - \sin(x) \sum_{i=1}^n A_{i1}b_i$$

Denote the following:

$$A = \frac{1}{2} \sum_{i=1}^n A_{i1}A_{i2} \quad B = \frac{1}{4i} \sum_{i=1}^n (A_{i2}^2 - A_{i1}^2) \quad C = \frac{1}{2} \sum_{i=1}^n A_{i2}b_i \quad D = \frac{1}{2i} \sum_{i=1}^n A_{i1}b_i$$

And using Euler expressions, we get:

$$f'(x) = A(e^{2ix} + e^{-2ix}) + B(e^{2ix} - e^{-2ix}) - C(e^{ix} + e^{-ix}) + D(e^{ix} - e^{-ix}) = 0$$

Let  $\theta \leftarrow e^{ix}$ , we are left with

$$\theta^4(A + B) + \theta^3(D - C) - \theta(C + D) + (A - B) = 0$$

This gives ap to four solutions and once filtering solutions which are not on the unit vectors, solution to  $x$  will be the real and imaginary coordinated of  $\theta_i = \text{cis}(x)$ .