5-Pair with Difference-O(n^2)Time Complexity,O(1) Space Complexity

Given an array A of sorted integers and another non negative integer k, find if there exists 2 indices i and j such that A[j] - A[i] = k, i != j.

Input Format:

First Line n - Number of elements in an array

Next n Lines - N elements in the array

k - Non - Negative Integer

Output Format:

1 - If pair exists

0 - If no pair exists

Explanation for the given Sample Testcase:

YES as 5 - 1 = 4

So Return 1.

For example:

Input	Result
3 1 3 5 4	1

Answer:(penalty regime: 0 %)

Feedback

```
#include <stdio.h>


int main()
{
   int n, k;


   scanf("%d", &n);


   int a[n];


   for (int i = 0; i < n; i++)
   {
```

```
    scanf("%d", &a[i]);
  }

  scanf("%d", &k);

  for (int i = 0; i < n; i++)
  {
    for (int j = 0; j < n; j++)
    {
      if (i != j && a[j] - a[i] == k)
      {
        printf("1");
        return 0;
      }
    }
  }

  printf("0");

  return 0;
}
```

	Input	Expected	Got	
	3 1 3 5 4	1	1	
	10	1	1	

	Input	Expected	Got	
	1 4 6 8 12 14 15 20 21 25 1			
	10 1 2 3 5 11 14 16 24 28 29 0	0	0	
	10 0 2 3 7 13 14 15 20 24 25 10	1	1	

Passed all tests!

Correct

Marks for this submission: 1.00/1.00.