# THE EFFECTS OF TELLURIC LINES IN RADIAL VELOCITY SEARCHES FOR PLANETS WITH IODINE CELL AS CALIBRATORS $^1$

Sharon Xuesong Wang (王雪凇)<sup>2,3</sup>, Jason T. Wright<sup>2,3</sup>, Chad Bender<sup>2,3</sup>, Andrew W. Howard<sup>4</sup>, Geoffrey W. Marcy<sup>5</sup>, Howard Isaacson<sup>5</sup>, and Suvrath Mahadevan<sup>2,3</sup>

### ABSTRACT

Tellurics are bad and you really don't want them. Here's how to get rid of them. Subject headings: instrumentation

### 1. INTRODUCTION

We are going to cite Artigau et al. (2014); Cunha et al. (2014) and Rothman et al. (2013).

### 2. Data and data analysis tools

Introduction to Keck/HIRES CPS data, data reduction, Doppler code. Our adoption of Doppler code.

Introduction to the standard star(s) in this paper? ZZZ Table: standard star stellar properties? esp. coordinates to highlight BC ranges?

### 3. EFFECTS OF TELLURIC LINES

### 3.1. Simulated Data

This is where TERRASPEC got introduced. ZZZ Plots:

- telluric line atlas in the optical? just to demonstrate where they are.
- telluric free simulations to demonstrate RV precision, vs. year, vs. BC
- telluric injected simulations to demonstrate effects, vs. year, vs. BC
- $\mbox{RV}$  vs. BC for chunks with and without telluric lines before and after injection

# 3.2. On-Sky Data

Should this be the first?

ZZZ Plots:

- RV vs. year/time, RV vs. BC
- RV from chunks with and without telluric lines, vs.  ${\rm BC}$

# 3.3. Where and How Telluric Lines Enter the Data

Should this subsection be moved to the first? It enters the epoch data as well as the empirically derived stellar template.

ZZZ Plots:

- fitting plots showing data (with telluric highlighted) and DSST (tellurics highlighted) and the mismatch
  - <sup>1</sup> Based on observations observations obtained at the Keck Observatory, which is operated by the University of California. The Keck Observatory was made possible by the generous financial support of the W. M. Keck Foundation.

<sup>2</sup> Department of Astronomy and Astrophysics, 525 Davey Laboratory, The Pennsylvania State University, University Park, PA 16802, USA; Send correspondence to xxw131@psu.edu and jtwright@astro.psu.edu

jtwright@astro.psu.edu <sup>3</sup> Center for Exoplanets and Habitable Worlds, 525 Davey Laboratory, The Pennsylvania State University, University Park, PA 16802, USA

<sup>4</sup> Hawaii, USA

 $^5\,\mathrm{Department}$  of Astronomy, University of California, Berkeley, CA 94720, USA

# 4. CORRECTING FOR THE EFFECTS OF TELLURIC LINES

# 4.1. Masking out the Telluric Lines

### ZZZ Plot:

- Illustration of choice of mask
- RV vs. BC before and after for all, telluric, and non-telluric chunks  $\,$
- simulation results showing no aliases introduced by masking
- 4.2. Removing the Telluric Contamination from Stellar Templates

### ZZZ Plot:

- selected chunk before and after cleaning
- ?RV vs. BC using cleaned template, no masking, single mask?, double mask?
- 4.3. Modeling Telluric Lines in the Star+Iodine Spectra ZZZ Plot:
- ${\rm RV}$  vs. BC using cleaned template plus full telluric modeling
- RV vs. BC using cleaned template plus full modeling plus some masking?

### 4.4. Results for A Different RV Standard Star

I think this is a must! At least one more star, hopefully two.

## 5. SUMMARY AND FUTURE WORK

Gaussian processes!

The authors thank John A. Johnson for providing a copy of his Doppler code and his help with our incorporation of the code into the HET pipeline. The authors also thank Debra Fischer for her assistance in this regard.

This work was partially supported by funding from the Center for Exoplanets and Habitable Worlds, which is supported by the Pennsylvania State University, the Eberly College of Science, and the Pennsylvania Space Grant Consortium.

The authors appreciate the significant Keck observing time and associated funding support from NASA for the study of long period planets and mulitplanet systems. J.T.W. and S.X.W. acknowledge support from NASA Origins of Solar Systems grant NNX10AI52G.

The work herein is based on observations obtained at the W. M. Keck Observatory, which is operated jointly by the University of California and the California Institute of Technology. The Keck Observatory was made possible by the generous financial support of the W.M. Keck Foundation. We wish to recognize and acknowledge the very significant cultural role and reverence that the summit of Mauna Kea has always had within the indigenous Hawaiian community. We are most fortunate to have the opportunity to conduct observations from this

mountain.

The Hobby-Eberly Telescope is a joint project of the University of Texas at Austin, the Pennsylvania State University, Stanford University, Ludwig Maximillians Universität München, and Georg August Universität Göttingen. The HET is named in honor of its principal benefactors, William P. Hobby and Robert E. Eberly.

This work has made use NASA's Astrophysics Data System Bibliographic Services.

### REFERENCES

Artigau, É., Astudillo-Defru, N., Delfosse, X., et al. 2014, in Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series, Vol. 9149, Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series, 5 Cunha, D., Santos, N. C., Figueira, P., et al. 2014, A&A, 568, A35 Rothman, L. S., Gordon, I. E., Babikov, Y., et al. 2013, J. Quant. Spec. Radiat. Transf., 130, 4