

IBD Endoscopic Activity Scores

Date of Scoring: _____

Endoscopist: _____

UCEIS (UC)

(Score the worst affected segment)

	0	1	2	3
Vascular pattern	Normal	Patchy obliteration	Obliterated	-
Bleeding	None	Mucosal	Luminal mild (some liquid blood in lumen)	Luminal moderate-severe (frank blood, visible oozing)
Erosions/ulcers	None	Erosions (<5mm)	Superficial ulcer	Deep ulcer

Total = ____ (0-8)

UCEIS \geq 7 predicts need for salvage therapy and colectomy (AUC = 0.85, sensitivity 60% specificity 85%)

Mayo Endoscopic Subscore (UC)

(Score the worst affected segment)

Mayo 0

Normal

Normal mucosa

Mayo 1

Mild

Mild erythema, friability, reduced vascular pattern

Mayo 2

Moderate

Marked erythema, lack of vascular pattern, friability, erosions

Mayo 3

Severe

Severe activity, spontaneous bleeding, large ulcers

Rutgeerts (Post-op CD recurrence)

i0

No lesions in ileum

Remission

i1

\leq 5 anastomotic aphthous lesions

Remission

i2

>5 aphthous lesions with normal areas in between, ulcers up to 1cm

Substantial recurrence

i3

Diffuse aphthous ileitis

Advanced recurrence

i4

Diffuse inflammation with large ulcers or narrowing/stenosis

Advanced recurrence

IBD Endoscopic Activity Scores

Date of Scoring: _____

Endoscopist: _____

SES-CD

		N/A	0	1	2	3
Ileum	Ulcer size	Not assessed	None	<5mm	0.5-2cm	>2cm
	Ulcerated surface	Not assessed	None	<10%	10-30%	>30%
	Affected surface	Not assessed	None	<50%	50-75%	>75%
	Strictures	Not assessed	None	Single	Multiple	Can't pass
Right colon	Ulcer size	Not assessed	None	<5mm	0.5-2cm	>2cm
	Ulcerated surface	Not assessed	None	<10%	10-30%	>30%
	Affected surface	Not assessed	None	<50%	50-75%	>75%
	Strictures	Not assessed	None	Single	Multiple	Can't pass
Transverse	Ulcer size	Not assessed	None	<5mm	0.5-2cm	>2cm
	Ulcerated surface	Not assessed	None	<10%	10-30%	>30%
	Affected surface	Not assessed	None	<50%	50-75%	>75%
	Strictures	Not assessed	None	Single	Multiple	Can't pass
Left colon	Ulcer size	Not assessed	None	<5mm	0.5-2cm	>2cm
	Ulcerated surface	Not assessed	None	<10%	10-30%	>30%
	Affected surface	Not assessed	None	<50%	50-75%	>75%
	Strictures	Not assessed	None	Single	Multiple	Can't pass
Rectum	Ulcer size	Not assessed	None	<5mm	0.5-2cm	>2cm
	Ulcerated surface	Not assessed	None	<10%	10-30%	>30%
	Affected surface	Not assessed	None	<50%	50-75%	>75%
	Strictures	Not assessed	None	Single	Multiple	Can't pass

Total = _____ (0-56)

(Inactive ≤2, Mild 3-6, Moderate 7-15, Severe ≥16)