

NexRay AI

Medical Assistant Platform • Clinical Decision Support Report
23 July 2026 | 08:24 PM

Session ID: 13	Modules Used: Symptom Checker	Date: 23 July 2026 08:24 PM
----------------	-------------------------------	-------------------------------

Symptom Analysis

Primary Condition	COVID-19 (SARS-CoV-2 Infection)
Likelihood	High
Assessment	The symptom cluster is highly consistent with COVID-19: respiratory symptoms (persistent cough, sore throat, runny/blocked nose, shortness of breath), systemic symptoms (chills, fatigue, muscle aches, headache), and gastrointestinal involvement (nausea, vomiting, diarrhea). The presence of anosmia/ageusia (loss or change in taste/smell) is particularly characteristic of COVID-19. The combination of upper and lower respiratory tract involvement with multi-system involvement strongly suggests COVID-19 over common viral respiratory infections.

■ Urgency Level: Urgent — Same day attention required

Recommended Tests

- SARS-CoV-2 RT-PCR (nasopharyngeal/oropharyngeal swab) - gold standard
- SARS-CoV-2 Rapid Antigen Test (if RT-PCR unavailable)
- Full blood count and differential
- C-reactive protein or procalcitonin
- Chest X-ray (if respiratory distress is significant)

Suggested Treatment

- Supportive care: adequate hydration, rest, and nutrition
- Paracetamol or ibuprofen for fever, myalgia, and headache
- Symptomatic management of cough with appropriate cough suppressants or expectorants
- Isolation precautions to prevent transmission to contacts
- Monitor oxygen saturation; consider supplemental oxygen if SpO2 <94%
- Consider antivirals (e.g., remdesivir, molnupiravir, or paxlovid) if severe or high-risk patient

Next Steps

- Confirm diagnosis with SARS-CoV-2 testing urgently
- Assess severity: evaluate oxygen saturation, respiratory rate, and degree of dyspnea
- Implement infection control measures and isolate patient
- Monitor for warning signs of severe disease (increasing dyspnea, persistent chest pain, confusion)
- Consider referral to higher facility if oxygen requirement develops or respiratory distress worsens

Summary: This presentation is highly suggestive of COVID-19 infection given the distinctive combination of respiratory symptoms, systemic symptoms, gastrointestinal involvement, and notably the loss/change in taste or smell. Urgent RT-PCR testing is recommended to confirm diagnosis. Management should focus on supportive care, isolation, and close monitoring for disease progression, particularly given the respiratory compromise indicated by shortness of breath.

■ **DISCLAIMER:** This report is generated by NexRay AI and is intended solely as a clinical decision-support tool. All findings, conditions, treatments, and recommendations are AI-generated suggestions and do not constitute a formal medical diagnosis or prescription. The clinical judgment of the attending medical professional must be applied before any action is taken.