

NexRay AI

Medical Assistant Platform • Clinical Decision Support Report
23 July 2026 | 09:40 PM

Session ID: 21	Modules Used: X-Ray Analysis	Date: 23 July 2026 09:40 PM
----------------	------------------------------	-------------------------------

X-Ray Analysis

Detected Region: Chest (Posteroanterior View)

Finding	Confidence
Right-sided pleural effusion	High
Normal left lung	High
Normal cardiac silhouette	High
Normal mediastinum	High

■ Urgency Level: Urgent — Same day attention required

Recommended Tests

- Chest ultrasound or CT chest with contrast to characterize the pleural effusion and assess for underlying lung pathology
- Thoracentesis with fluid analysis (cell count, protein, LDH, culture, cytology) to determine the etiology of the effusion
- Complete blood count and metabolic panel
- ECG to evaluate for cardiac cause
- Echocardiogram if cardiac pathology is suspected

Suggested Treatment

- Investigate underlying cause of pleural effusion (cardiac, infectious, malignant, or other etiology)
- Supportive care and oxygen therapy if hypoxemic
- Diuretics if pleural effusion is related to cardiac or renal disease
- Consider thoracentesis for diagnostic and/or therapeutic purposes if the effusion is causing respiratory compromise

Next Steps

- Clinical correlation with patient history, symptoms, and vital signs to determine the underlying cause
- Perform thoracentesis if patient is symptomatic or if diagnostic evaluation is needed
- Obtain imaging (ultrasound or CT) to further evaluate the nature of the effusion and underlying lung parenchyma
- Monitor oxygen saturation and respiratory status

Summary: Chest X-ray shows a moderate right-sided pleural effusion with otherwise normal cardiac and left lung appearances. The left lung is clear with no evidence of consolidation or acute pathology.

■ **DISCLAIMER:** This report is generated by NexRay AI and is intended solely as a clinical decision-support tool. All findings, conditions, treatments, and recommendations are AI-generated suggestions and do not constitute a formal medical diagnosis or prescription. The clinical judgment of the attending medical professional must be applied before any action is taken.

