

NexRay AI

Medical Assistant Platform • Clinical Decision Support Report
23 July 2026 | 10:26 PM

Session ID: 24	Modules Used: Symptom Checker	Date: 23 July 2026 10:26 PM
----------------	-------------------------------	-------------------------------

Symptom Analysis

Possible Condition	Confidence
COVID-19 (SARS-CoV-2)	92%
Influenza (Flu)	78%
Malaria with secondary respiratory infection	45%

■ Urgency Level: Urgent — Same day attention required

Recommended Tests

- COVID-19 RT-PCR or rapid antigen test (nasopharyngeal/oropharyngeal swab)
- Malaria rapid diagnostic test (RDT) or blood smear
- Full blood count (FBC) with differential
- Chest X-ray (if shortness of breath is significant)
- Influenza rapid test (if COVID-19 negative and clinical suspicion remains high)

Suggested Treatment

- If COVID-19 confirmed: supportive care including rest, hydration, paracetamol or ibuprofen for fever and body aches; monitor oxygen saturation closely given shortness of breath
- If malaria co-infection suspected: commence appropriate antimalarial therapy (artemisinin-based combination therapy preferred in West Africa)
- Manage respiratory symptoms: monitor oxygen saturation, consider oxygen supplementation if SpO2 <94%, assess for need for escalated care
- Consider empiric broad-spectrum antibiotic coverage if secondary bacterial infection suspected based on clinical deterioration

Next Steps

- Prioritize COVID-19 testing immediately to guide isolation precautions
- Perform malaria screening given West African context and fever presentation
- Monitor respiratory status closely; document oxygen saturation and respiratory rate
- Obtain chest imaging if shortness of breath worsens or does not improve with supportive care
- Review test results and adjust treatment plan accordingly within 24 hours

Summary: Patient presents with a constellation of symptoms highly suggestive of COVID-19, particularly given the specific triad of fever, dry cough, and loss of taste/smell. Influenza remains a differential, and malaria should not be excluded in the West African context. The presence of shortness of breath elevates clinical concern and warrants prompt respiratory assessment. Immediate testing for COVID-19 and malaria is recommended, with supportive care and close monitoring of respiratory status.

■ **DISCLAIMER:** This report is generated by NexRay AI and is intended solely as a clinical decision-support tool. All findings, conditions, treatments, and recommendations are AI-generated suggestions and do not constitute a formal medical diagnosis or prescription. The clinical judgment of the attending medical professional must be applied before any action is taken.