

Tylenol - Acetaminophen tablet Johnson & Johnson Consumer Inc.

Tylenol ®

Drug Facts

Active ingredient (in each tablet)

Acetaminophen 325 mg

Purpose

Pain reliever/fever reducer

Uses

temporarily relieves minor aches and pains due to: headache

muscular aches

backache

minor pain of arthritis

the common cold

toothache

premenstrual and menstrual cramps

temporarily reduces fever

Warnings Liver warning

This product contains Acetaminophen. The maximum daily dose of this product is 10 tablets (3,250 mg) in 24 hours for adults or 5 tablets (1,625 mg) in 24 hours for children. Severe liver damage may occur if

adult takes more than 4,000 mg of Acetaminophen in 24 hours

child takes more than 5 doses in 24 hours, which is the maximum daily amount taken with other drugs containing Acetaminophen

adult has 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

skin reddening blisters

rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

with any other drug containing Acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains Acetaminophen, ask a doctor or pharmacist. if you are allergic to Acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if the user has liver disease

Ask a doctor or pharmacist before use if the user is taking the blood thinning

drug warfarin

Stop use and ask a doctor if

pain gets worse or lasts more than 10 days in adults

pain gets worse or lasts more than 5 days in children under 12 years fever gets worse or lasts more than 3 days

new symptoms occur

redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.**

Overdose warning

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

do not take more than directed (see overdose warning)

adults and children 12 years and over

children 6 years to under 12 years

take 2 tablets every 4 to 6 hours while symptoms last do not take more than 10 tablets in 24 hours, unless directed by a doctor

do not use for more than 10 days unless directed by a doctor

take 1 tablet every 4 to 6 hours while symptoms last
do not take more than 5 tablets in 24 hours

do not use for more than 5 days unless directed by a

years

doctor

children
under 6 ask a doctor years

Other information

store between 20-25°C (68-77°F)

**do not use if carton is opened. Do not use if foil inner seal imprinted with
"Tylenol" is broken or missing**

Inactive ingredients

magnesium stearate, modified starch, powdered cellulose, pregelatinized starch, sodium
starch glycolate

Questions or comments?

call **1-877-895-3665** (toll-free) or **215-273-8755** (collect)

PRINCIPAL DISPLAY PANEL

NDC 50580-496-98 Tylenol[®] Acetaminophen Pain Reliever

Fever Reducer Actual Size Regular Strength

100 Tablets 325 mg each