

Question 4

In [1]:

```
def eucl_dist(a,b):
    distance = 0
    for i in range(len(a)):
        distance += (a[i]-b[i])**2
    return round(float(distance)**0.5,3)
```

In [30]:

```
def find_center(cluster):
    C_x = 0
    C_y = 0
    for i in cluster:
        C_x += i[0]
        C_y += i[1]
    center = [float(C_x)/len(cluster), float(C_y)/len(cluster)]
    return center
```

In [35]:

```
def cluster(data,center1,center2):
    C1 = []
    C2 = []
    for i in data:
        if eucl_dist(i,center1) < eucl_dist(i,center2):
            C1.append(i)
        else:
            C2.append(i)
    print C1
    print C2
    print find_center(C1)
    print find_center(C2)
    return find_center(C1),find_center(C2)
```

Q.a

In [32]:

```
data = [[1,1],[1,2],[2,1],[5,1],[3,2],[5,2],[3,3]]
```

In [36]:

```
center1, center2 = cluster(data,data[0],data[2])
```

```
[[1, 1], [1, 2]]
[[2, 1], [5, 1], [3, 2], [5, 2], [3, 3]]
[1.0, 1.5]
[3.6, 1.8]
```

In [39]:

```
center1, center2 = cluster(data,center1, center2)
```

```
[[1, 1], [1, 2], [2, 1]]
[[5, 1], [3, 2], [5, 2], [3, 3]]
[1.3333333333333333, 1.3333333333333333]
[4.0, 2.0]
```

In [41]:

```
center1, center2 = cluster(data,center1, center2)
```

```
[[1, 1], [1, 2], [2, 1]]
[[5, 1], [3, 2], [5, 2], [3, 3]]
[1.3333333333333333, 1.3333333333333333]
[4.0, 2.0]
```

We can find there is no change in the **3rd** iteration. The k-means finished.

Q.b

In [42]:

```
center1, center2 = cluster(data,data[1],data[5])
```

```
[[1, 1], [1, 2], [2, 1]]
[[5, 1], [3, 2], [5, 2], [3, 3]]
[1.3333333333333333, 1.3333333333333333]
[4.0, 2.0]
```

In [43]:

```
center1, center2 = cluster(data,center1, center2)
```

```
[[1, 1], [1, 2], [2, 1]]
[[5, 1], [3, 2], [5, 2], [3, 3]]
[1.3333333333333333, 1.3333333333333333]
[4.0, 2.0]
```

We can find there is no change in the **2nd** iteration. The k-means finished.

Q.c

Randomly set one point as 1st center, find the longest point from the center as the 2nd center

In [51]:

```
import random as rd
```

In [64]:

```
a = data[rd.randint(0,6)]
```

In [65]:

```
for i in data:  
    print eucl_dist(i,a)
```

```
2.828  
2.236  
2.236  
2.828  
1.0  
2.236  
0.0
```

We can find P7 is the random select data, P3 is the farrest from P7

In [66]:

```
b= data[2]
```

In [67]:

```
center1, center2 = cluster(data,a, b)
```

```
[[5, 1], [3, 2], [5, 2], [3, 3]]  
[[1, 1], [1, 2], [2, 1]]  
[4.0, 2.0]  
[1.3333333333333333, 1.3333333333333333]
```

In [68]:

```
center1, center2 = cluster(data,center1, center2)
```

```
[[5, 1], [3, 2], [5, 2], [3, 3]]  
[[1, 1], [1, 2], [2, 1]]  
[4.0, 2.0]  
[1.3333333333333333, 1.3333333333333333]
```

It get to the center in the **1st** step.