

Thesis

Shay Ben Moshe

September 15, 2018

Contents

1	Overview of Chromatic Homotopy Theory	2
1.1	The Balmer Spectrum	2
1.2	MU and Complex Orientations	4
1.3	BP, Morava K-Theory and Morava E-Theory	7
1.4	$\mathrm{Spec} \mathbb{S}_{(p)}$ and $\mathrm{Spec} \mathbb{S}$	9
1.5	The Stacky Point of View and the Relationship Between Morava K-Theory and Morava E-Theory	11
1.6	Landweber Exact Functor Theorem	12
1.7	Lubin-Tate Deformation Theory	14
2	Atiyah-Segal	17
2.1	The Atiyah-Segal Theorem	17
2.2	Examples	19
2.3	Recollections from Character Theory	21
3	HKR Character Theory	23
3.1	Towards a Definition of the Character Map	24
3.2	The E -Cohomology of BA	25
3.3	The Rings $L_r(E^*)$ and $L(E^*)$	26

1 Overview of Chromatic Homotopy Theory

Our goal is to motivate the introduction of Morava K-theory $K(n)$ and Morava E-theory $E(n)$, and other variants of Morava E-theory $E(k, \Gamma)$, and their connection to formal group laws. There are different views on what chromatic homotopy theory is. **TODO write some more**

1.1 The Balmer Spectrum

We will start with an algebraic motivation. Let R be a noetherian ring. Consider the symmetric monoidal stable ∞ -category $\mathrm{Ch}(R)$ of chain complexes on R . **TODO be more specific** It is then natural to ask how much information about R is encoded in the category $\mathrm{Ch}(R)$. We will try to recover $\mathrm{Spec} R$, as a topological space, from $\mathrm{Ch}(R)$.

Remark 1.1.1. Balmer's work actually recovers the structure sheaf as well. **TODO reference**

Definition 1.1.2. A *perfect complex* is a complex that is quasi-isomorphic to a bounded complex of finitely-generated projective modules. These objects are the compact objects in $\mathrm{Ch}(R)$, thus they can be defined categorically. Their full subcategory is denoted by $\mathrm{Ch}_{\mathrm{perf}}(R)$.

Definition 1.1.3. Let \mathcal{C} be some symmetric monoidal stable ∞ -category. A full subcategory \mathcal{T} is *thick* if:

- $0 \in \mathcal{T}$,
- it is closed under cofibers,
- it is closed under retracts.

Example 1.1.4. Consider the case $\mathcal{C} = \mathrm{Ch}_{\mathrm{perf}}(R)$ (e.g. over \mathbb{Z} , chain complexes quasi-isomorphic to bounded chain complexes of finitely-generated free abelian groups). Let $K \in \mathrm{Ch}(R)$, and define $\mathcal{T}_K = \{A \in \mathrm{Ch}_{\mathrm{perf}}(R) \mid A \otimes K \cong 0\}$. We claim that \mathcal{T}_K is thick. Clearly $0 \in \mathcal{T}_K$. Let $A \rightarrow B$ be a morphism between two complexes in \mathcal{T} . Since tensor is left, tensoring the cofiber with K is given by $\mathrm{cofib}(A \rightarrow B) \otimes K \cong \mathrm{cofib}(A \otimes K \rightarrow B \otimes K) \cong \mathrm{cofib}(0 \rightarrow 0) \cong 0$, therefore the cofiber is indeed in \mathcal{T}_K . Lastly, if $A \rightarrow B \rightarrow A$ is the identity and $B \otimes K \cong 0$, we get that $\mathrm{id}_{A \otimes K}$ factors through 0, which implies that $A \otimes K$ is 0, so that $A \in \mathcal{T}_K$.

Definition 1.1.5. A thick subcategory \mathcal{T} is an *ideal* if $A \in \mathcal{T}, B \in \mathcal{C} \implies A \otimes B \in \mathcal{T}$. Furthermore, it is a *prime ideal* if it is a proper subcategory, and $A \otimes B \in \mathcal{T} \implies A \in \mathcal{T} \text{ or } B \in \mathcal{T}$. The *spectrum* of the category is defined similarly to the classical spectrum of a ring: As a set, $\text{Spec } \mathcal{C} = \{\mathcal{P} \text{ prime ideal}\}$. For any family of objects $S \subseteq \mathcal{C}$ we define $V(S) = \{\mathcal{P} \in \text{Spec } \mathcal{C} \mid S \cap \mathcal{P} = \emptyset\}$. We topologize $\text{Spec } \mathcal{C}$ with the Zariski topology by declaring those to be the closed subsets. We also denote $\text{Supp}(A) = V(\{A\})$.

Example 1.1.6. We continue the example of \mathcal{T}_K . Clearly if $A \otimes K \cong 0$ then also $A \otimes B \otimes K \cong 0$, so it is an ideal. Let \mathfrak{p} be a prime ideal in R in the usual sense, and take $K = R_{\mathfrak{p}}$ (concentrated at degree 0), then $A \otimes K = A_{\mathfrak{p}}$ (level-wise localization). **TODO consider actually proving** We will omit the proof that \mathcal{T}_K is a prime, but we shall prove something weaker, namely only the case where A, B are bounded complexes of finitely generate projective modules (and not merely quasi-isomorphic to such complexes). Assume then that $0 = (A \otimes B)_{\mathfrak{p}} = A_{\mathfrak{p}} \otimes_{R_{\mathfrak{p}}} B_{\mathfrak{p}}$. Assume by negation that $A_{\mathfrak{p}}, B_{\mathfrak{p}} \neq 0$, i.e. $(A_n)_{\mathfrak{p}}, (B_m)_{\mathfrak{p}} \neq 0$ but $(A_n)_{\mathfrak{p}} \otimes_{R_{\mathfrak{p}}} (B_m)_{\mathfrak{p}} = 0$ for some n, m . Since the localization of a projective module is again a projective module, and a projective over a local ring is free, and clearly if the tensor of two free modules vanish then one of them vanishes, it follows that $(A_n)_{\mathfrak{p}} = 0$ or $(B_m)_{\mathfrak{p}} = 0$, which is a contradiction. Therefore $\mathcal{T}_{\mathfrak{p}}$ is a prime ideal.

Theorem 1.1.7. *The map $\text{Spec } R \rightarrow \text{Spec}(\text{Ch}_{\text{perf}}(R))$, given by $\mathfrak{p} \mapsto \mathcal{T}_{\mathfrak{p}} = \{A \mid A_{\mathfrak{p}} = 0\}$ is a homeomorphism.*

TODO reference

Proposition 1.1.8. *Prime ideals pullback: Let $F : \mathcal{C} \rightarrow \mathcal{D}$ be a reduced symmetric monoidal functor that preserves cofibers, between two symmetric monoidal stable ∞ -categories, and let \mathcal{P} be a prime ideal in \mathcal{D} , then $F^*\mathcal{P} = \{A \in \mathcal{C} \mid F(A) \in \mathcal{P}\}$ is a prime ideal.*

Proof. Clearly $F(0) = 0 \in \mathcal{P}$ since F is reduced, so $0 \in F^*\mathcal{P}$. Since F preserves cofibers, for $A, B \in F^*\mathcal{P}$, i.e. $F(A), F(B) \in \mathcal{P}$, and a map $A \rightarrow B$ we get $F(\text{cofib}(A \rightarrow B)) = \text{cofib}(F(A) \rightarrow F(B)) = \text{cofib}(F(A) \rightarrow F(B)) \in \mathcal{P}$. Let $A \rightarrow B \rightarrow A$ be a retract, that is the composition is the identity, s.t. $B \in F^*\mathcal{P}$. We know that $F(A) \rightarrow F(B) \rightarrow F(A)$ is also a retract by functoriality, thus $F(A) \in \mathcal{P}$, that is $A \in F^*\mathcal{P}$. We conclude that $F^*\mathcal{P}$ is indeed a thick subcategory.

Let $A \in F^*\mathcal{P}, B \in \mathcal{C}$, since F is monoidal, $F(A \otimes B) = F(A) \otimes F(B) \in \mathcal{P}$, so $A \otimes B \in F^*\mathcal{P}$, that is $F^*\mathcal{P}$ is an ideal.

Lastly, assume that $A \otimes B \in F^*\mathcal{P}$, again since F is monoidal, $F(A \otimes B) = F(A) \otimes F(B) \in \mathcal{P}$, so $A \in F^*\mathcal{P}$ or $B \in F^*\mathcal{P}$, that is $F^*\mathcal{P}$ is a prime ideal. \square

Now, recall that $\mathrm{Ch}(R) \cong \mathrm{Mod}_{\mathrm{HR}}$, therefore we can reinterpret the above theorem as $\mathrm{Spec} R \cong \mathrm{Spec}(\mathrm{Mod}_{\mathrm{HR}}^{\mathrm{comp}})$ (where the comp denotes the compact objects in the category). We shall turn this theorem into a definition:

Definition 1.1.9. Let R be an \mathbb{E}_∞ -ring. We define the *spectrum* of R to be $\mathrm{Spec} R = \mathrm{Spec}(\mathrm{Mod}_R^{\mathrm{comp}})$.

A natural question to ask then is what is the topological space $\mathrm{Spec} \mathbb{S}$. Recall that $\mathrm{Mod}_{\mathbb{S}} = \mathrm{Sp}$, the category of spectra, and that the compact objects in spectra are the finite spectra $\mathrm{Sp}^{\mathrm{fin}}$. So, unwinding the definitions, the question can be rephrased as finding the prime ideals in $\mathrm{Sp}^{\mathrm{fin}}$, and their topology. Chromatic homotopy theory provides an answer to this question.

1.2 MU and Complex Orientations

Throughout this section, let E be a multiplicative cohomology theory (that is, equipped with a map $E \otimes E \rightarrow E$ and $1 \in E_0$, which is associative and unital after taking homotopy groups).

Consider the map $S^2 \rightarrow \mathrm{BU}(1)$ classifying the universal complex line bundle. Concretely, under the identifications $S^2 \cong \mathbb{CP}^1$ and $\mathrm{BU}(1) \cong \mathbb{CP}^\infty$, this map can be realized as the inclusion $\mathbb{CP}^1 \subseteq \mathbb{CP}^\infty$. This map induces a map

$$\tilde{E}^2(\mathrm{BU}(1)) \rightarrow \tilde{E}^2(S^2) \cong \tilde{E}^0(S^0) \cong E^0(*) = E_0.$$

Since E is unital, there is a canonical generator $1 \in E_0$.

Definition 1.2.1. E is called *complex oriented* if the map $\tilde{E}^2(\mathrm{BU}(1)) \rightarrow E_0$ is surjective, equivalently, if 1 is in the image of that map. A choice of a lift $x \in \tilde{E}^2(\mathrm{BU}(1))$ of $1 \in E_0$ is called a *complex orientation*.

Example 1.2.2. Let R be some ring, and consider HR . It is known that $\mathrm{HR}^*(\mathbb{CP}^n) \cong R[x]/(x^{n+1})$ and $\mathrm{HR}^*(\mathbb{CP}^\infty) \cong R[[x]]$, where $|x| = 2$, and the maps induced by the inclusions of projective spaces maps x to x . Therefore we see that $x \in \mathrm{HR}^2(\mathrm{BU}(1))$ is mapped to $x \in \mathrm{HR}^2(S^2) = R\{x\}$, which is mapped to the generator of the reduced part of $\mathrm{HR}^0(S^0) = R \oplus R$. Hence, x is a complex orientation for HR .

Example 1.2.3. Let K be complex K -theory, then we know that $K_* = \mathbb{Z} [\beta^{\pm 1}]$ where β is the Bott element, with $|\beta| = 2$. It is also known (by the Atiyah-Hirzebruch spectral sequence) that $K^*(\mathbb{CP}^n) \cong K_*[t] / (t^{n+1})$ and $K^*(\mathbb{CP}^\infty) \cong K_*[[t]]$ (here $|t| = 0$), where the maps induced by the inclusions of projective spaces maps t to t . We deduce that $\beta^{-1}t \in K^2(BU(1))$ is mapped to $\beta^{-1}t \in K^2(S^2) = \mathbb{Z} \{\beta^{-1}t\}$, and further mapped to $t \in K^0(S^0) = \mathbb{Z} \{t\}$, which is indeed the generator of the reduced part. Therefore $x = \beta^{-1}t$ is complex orientation for K . **TODO write the reduced thing more clearly**

Example 1.2.4. Recall that MU is constructed as the colimit $MU = \text{colim } MU(n)$. Also, $MU(1) \cong \Sigma^{\infty-2}BU(1)$. Therefore we get a canonical map $\Sigma^{\infty-2}BU(1) \rightarrow MU$, which gives a cohomology class $x_{MU} \in MU^2(BU(1))$.

Proposition 1.2.5 ([Rav86, 4.1.3]). x_{MU} is a complex orientation for MU .

Theorem 1.2.6 (TODO reference). MU is the universal complex oriented cohomology theory, in the following sense: For any multiplicative cohomology theory E , then there is a bijection between (homotopy classes of) multiplicative maps $MU \rightarrow E$ and complex orientations on E . The bijection is given in one direction by pulling back x_{MU} along a multiplicative map.

Assume that E is complex oriented with a complex orientation x .

Proposition 1.2.7 ([Rav86, 4.1.4]). As E_* -algebras, $E^*(BU(1)) \cong E^*[[x]]$ and $E^*(BU(1) \times BU(1)) \cong E^*[[y, z]]$.

TODO maybe indicate the use of AHSS

There is a multiplication map for the group $U(1)$, i.e. $U(1) \times U(1) \rightarrow U(1)$. We can take the B of this map, and since it commutes with products we get a map $BU(1) \times BU(1) \rightarrow BU(1)$, which is the universal map the classifies the tensor product of vector bundles. Therefore we get a map $E^*(BU(1)) \rightarrow E^*(BU(1) \times BU(1))$, which is completely determined by the image of $x \in E^*[[x]]$ in $E^*[[y, z]]$ as above. We conclude that a choice of a complex orientation on E gives rise to an element $F_E(y, z) \in E^*[[y, z]]$.

Proposition 1.2.8 ([Rav86, 4.1.4]). F_E is a formal group law on E_* .

Definition 1.2.9. The *height* of E is simply the height of F_E .

Example 1.2.10. We continue with HR from 1.2.2. It is known that the tensor of complex line bundles induces the map

$$R[[x]] = HR^*(BU(1)) \rightarrow HR^*(BU(1) \times BU(1)) = R[[y, z]],$$

given by $x \mapsto y + z$. This is the additive formal group law. It is immediate that $[p] = px$. So for $R = \mathbb{Q}$ we get that the height of $H\mathbb{Q}$ is 0, while for $R = \mathbb{F}_p$ we have $px = 0$ so the height of $H\mathbb{F}_p$ is ∞ .

Example 1.2.11. We return to the example of complex K-theory 1.2.3. It is known that the tensor of complex line bundles induces the map

$$K_*[[t]] = K^*(BU(1)) \rightarrow K^*(BU(1) \times BU(1)) = K_*[[u, v]],$$

given by $t \mapsto u + v + uv$. Note that to comply with the definition of the formal group law, we should use the isomorphism $K^*(BU(1)) \cong K_*[[x]]$, i.e. the element $x = \beta t$. By multiplying by β^{-1} (recall that the map is of K_* -modules) we get that

$$x = \beta^{-1}t \mapsto \beta^{-1}u + \beta^{-1}v + \beta^{-1}uv = y + z + \beta yz = F_K(y, z).$$

By induction we prove that the n -series is $[n](x) = \beta^{-1}(1 + \beta x)^n - \beta^{-1}$. This is clear for $n = 1$, and we have:

$$\begin{aligned} [n+1](x) &= x + [n](x) + \beta x [n](x) \\ &= x + \beta^{-1}(1 + \beta x)^n - \beta^{-1} + x(1 + \beta x)^n - x \\ &= \beta^{-1}(1 + \beta x)(1 + \beta x)^n - \beta^{-1} \\ &= \beta^{-1}(1 + \beta x)^{n+1} - \beta^{-1} \end{aligned}$$

TODO consider discussing the computation of $BU(1)$, maybe as part of complex K-theory example?

Example 1.2.12. By taking the cofiber of the multiplication-by- p map, we get a spectrum K/p , mod- p K-theory, with coefficients $(K/p)_* = \mathbb{F}_p[\beta^{\pm 1}]$. It is evident that $F_{K/p}(y, z) = y + z + \beta yz$ as well. From the result above, it follows that

$$[p](x) = \beta^{-1}(1 + \beta x)^p - \beta^{-1} = \beta^{-1}(1^p + \beta^p x^p) - \beta^{-1} = \beta^{p-1}x^p,$$

which shows that the height is exactly 1.

A formal group law on E_* is the same data as a map from the Lazard ring L , so the complex orientation gives a map $L \rightarrow E_*$. In particular, since MU is complex oriented, there is a canonical map $L \rightarrow MU_*$.

Theorem 1.2.13 (Quillen, [Rav86, 4.1.6]). *The canonical map $L \rightarrow MU_*$ is an isomorphism.*

TODO consider adding something about the $X(n)$, and the obstruction to lifting to a complex orientation, with obstructions living in odd E_k , so even are automatically orientable

1.3 BP, Morava K-Theory and Morava E-Theory

A good principle in homotopy theory (and in many other areas in math) is to study it one prime at a time. This is possible in homotopy theory due to the arithmetic square **TODO reference**. So, let us fix a prime p . We can form $\mathrm{MU}_{(p)}$, the p -localization of MU .

Theorem 1.3.1 ([Ada74, II 15]). *There exists a unique map of ring spectra $\varepsilon : \mathrm{MU}_{(p)} \rightarrow \mathrm{MU}_{(p)}$ (depending on the prime p) satisfying:*

- ε is an idempotent, i.e. $\varepsilon^2 = \varepsilon$,
- ε_* sends $[\mathbb{CP}^n] \in \pi_*(\mathrm{MU}_{(p)})$ to itself if $n = p^r - 1$ and to 0 otherwise.

The map $\varepsilon : \mathrm{MU}_{(p)} \rightarrow \mathrm{MU}_{(p)}$ gives a cohomology operation, for every X we have $\varepsilon^* : \mathrm{MU}_{(p)}^*(X) \rightarrow \mathrm{MU}_{(p)}^*(X)$. Denote by $\mathrm{BP}_{(p)}^*(X)$ the image of ε^* .

Theorem 1.3.2 ([Ada74, II 16], [Rav86, 4.1.12]). *BP is a cohomology theory, represented by an associative commutative ring spectrum BP (depending on the prime p), which is a retract of $\mathrm{MU}_{(p)}$. The homotopy groups of BP are $\mathrm{BP}_* = \mathbb{Z}_{(p)}[v_1, v_2, \dots]$ where $|v_n| = 2(p^n - 1)$.*

For convenience we denote $v_0 = p$ (and indeed $|v_0| = 2(p^0 - 1) = 0$). Since BP is a retract of MU , it comes with a map $\mathrm{MU} \rightarrow \mathrm{BP}$, that is, a complex orientation.

Proposition 1.3.3. [**TODO reference**] *The p -series of the formal group law associated to BP is $[p](x) = \sum v_n x^{p^n}$.*

Remark 1.3.4 ([Rav92, B.5]). The formal group law on BP has a similar interpretation to that of MU , namely it is the universal p -typical formal group law. Moreover, the idempotent $\varepsilon : \mathrm{MU}_{(p)} \rightarrow \mathrm{MU}_{(p)}$ induces an idempotent on homotopy groups, which can be described as the map that takes a formal group law to the canonical p -typical formal group law isomorphic to it.

Once we have BP, we can turn to the definition of Morava K-theory and Morava E-theory

Definition 1.3.5. Let $0 < n < \infty$. *Morava K-theory at height n and prime p , denoted by $\mathrm{K}(p, n)$ or $\mathrm{K}(n)$ when the prime is clear from the context, is the spectrum obtained by killing $p = v_0, \dots, v_{n-1}, v_{n+1}, \dots$ in BP and inverting v_n . Therefore $\mathrm{K}(n)_* = \mathbb{F}_p[v_n^{\pm 1}]$. We also define $\mathrm{K}(0) = \mathrm{H}\mathbb{Q}$ and*

$K(\infty) = \mathrm{HF}_p$. Similarly, *Morava E-theory* at height n and prime p , denoted by $E(p, n)$ or $E(n)$, is the spectrum obtained by killing v_{n+1}, v_{n+2}, \dots in BP and inverting v_n . Therefore $E(n)_* = \mathbb{Z}_{(p)}[v_1, \dots, v_{n-1}, v_n^{\pm 1}]$.

Since Morava K-theory and E-theory are obtained from BP by cofibers and filtered colimits, they are equipped with a map from BP, hence also with a complex orientation. Then, from 1.3.3, we get:

Corollary 1.3.6. *The p -series associated to the formal group laws of $K(n)$ and $E(n)$ are $v_n x^{p^n}$ and $v_0 x + \dots v_n x^{p^n}$ respectively. Therefore the height of $K(n)$ is exactly n . (Note that by the example of HR, this is also true for $K(0)$ and $K(\infty)$.)*

We want to describe some properties of Morava K-theory. To do so we first need some definitions.

Definition 1.3.7. Let R be an evenly graded ring. R is called a *graded field* if it satisfies one of the equivalent conditions:

- every non-zero homogenous element is invertible,
- it is a field F concentrated at degree 0, or $F[\beta^{\pm 1}]$ for β of positive even degree.

An \mathbb{A}_∞ -ring E is a *field* if E_* is a graded field.

Example 1.3.8. $K(n)$ is a field for $0 \leq n \leq \infty$.

Proposition 1.3.9. *A field E has Kunneth, i.e. $E_*(X \otimes Y) \cong E_*(X) \otimes_{E_*} E_*(Y)$ for any spectra X, Y .*

Proposition 1.3.10 ([Lur10, 24]). *Let $E \neq 0$ be a complex oriented cohomology theory, whose formal group law has height exactly n , then $E \otimes K(n) \neq 0$. Let E be a field s.t. $E \otimes K(n) \neq 0$, then E admits the structure of a $K(n)$ -module. (Here $0 \leq n \leq \infty$.)*

Example 1.3.11. As we have seen in 1.2.12, mod- p K-theory, K/p , has height exactly 1 and coefficients $(K/p)_* = \mathbb{F}_p[\beta^{\pm 1}]$. It is also known that K and K/p , are A_∞ ring spectra, from which it follows that K/p is a field. We deduce that K/p is a $K(1)$ -module. Since $|\beta| = 2$ and $|v_1| = 2(p-1)$ it is free of rank $p-1$.

From this we also deduce some form of uniqueness for Morava K-theory:

Corollary 1.3.12. *Let E be a field with $E_* \cong \mathbb{F}_p[v_n^{\pm 1}]$, which is also complex oriented of height exactly n . Then $E \cong K(n)$ (as spectra).*

1.4 $\mathrm{Spec} \mathbb{S}_{(p)}$ and $\mathrm{Spec} \mathbb{S}$

We are now in a position to describe the topological space $\mathrm{Spec} \mathbb{S}$. However, it will be easier to state it first for $\mathrm{Spec} \mathbb{S}_{(p)}$, and then pullback prime ideals. We know that $\mathrm{Mod}_{\mathbb{S}_{(p)}} = \mathrm{Sp}_{(p)}$, and its compact objects are $\mathrm{Sp}_{(p)}^{\mathrm{fin}}$, the p -localizations of finite spectra.

Proposition 1.4.1. *Let \mathcal{T}_E be the E -acyclics, i.e.*

$$\mathcal{T}_E = \ker E_* = \left\{ X \in \mathrm{Sp}_{(p)}^{\mathrm{fin}} \mid E_*(X) = 0 \right\} = \left\{ X \in \mathrm{Sp}_{(p)}^{\mathrm{fin}} \mid X \otimes E_* = 0 \right\}.$$

Then \mathcal{T}_E is thick.

Proof. The proof follows the same lines of 1.1.4 for the case $\mathrm{Ch}_{\mathrm{perf}}(R)$. \square

Definition 1.4.2. We define $\mathcal{C}_{p,n} = \mathcal{T}_{K(n)}$, the $K(n)$ -acyclics. By the above proposition, it is thick. Also, $\mathcal{C}_{p,\infty} = \{0\}$, which is trivially thick. When the prime is clear from the context, we write \mathcal{C}_n in place of $\mathcal{C}_{p,n}$.

Proposition 1.4.3 ([Lur10, 26]). *For $X \in \mathrm{Sp}_{(p)}^{\mathrm{fin}}$, if $K(n)_*(X) = 0$ then $K(n-1)_*(X) = 0$.*

Definition 1.4.4. We say that a spectrum $X \in \mathrm{Sp}_{(p)}^{\mathrm{fin}}$ is of *type n* (possibly ∞) if its first non-zero Morava K -theory homology is $K(n)$.

Corollary 1.4.5. \mathcal{C}_n is the full subcategory of finite p -local spectra of type $> n$, that is $\mathcal{C}_n = \left\{ X \in \mathrm{Sp}_{(p)}^{\mathrm{fin}} \mid \forall m \leq n : K(m)_*(X) = 0 \right\}$. Thus $\mathcal{C}_{n+1} \subseteq \mathcal{C}_n$.

Proposition 1.4.6 (TODO reference). *The inclusions $\mathcal{C}_{n+1} \subset \mathcal{C}_n$ are proper.*

Proposition 1.4.7. *If $X \in \mathrm{Sp}_{(p)}^{\mathrm{fin}}$ is not contractible, then is of finite type. Therefore $\bigcap_{n < \infty} \mathcal{C}_n = \{0\} = \mathcal{C}_{\infty}$.*

Proof. Let X be non-contractible. Then $H\mathbb{Z}_*(X) \neq 0$. Let m be the first non-zero degree. Using the universal coefficient theorem and the fact that the spectrum is p -local we get that $(H\mathbb{F}_p)_m(X) \neq 0$, thus $(H\mathbb{F}_p)_*(X) \neq 0$. Since X is finite, $(H\mathbb{F}_p)_*(X)$ is bounded. The Atiyah-Hirzebruch spectral sequence for X with cohomology $K(n)$ has E^2 -page given by $E_{p,q}^2 = H_p(X; K(n)_q)$. Since $K(n)_q = \mathbb{F}_p$ for $q = 0 \bmod 2(p^n - 1)$ and 0 otherwise, we see that the rows $q = 0 \bmod 2(p^n - 1)$ are $(H\mathbb{F}_p)_*(X)$, and the

others are 0. Therefore if we take n such that the period $2(p^n - 1)$ is larger than the bound on $(\mathrm{H}\mathbb{F}_p)_*(X)$, then all differentials have either source or target 0. Thus, the spectral sequence collapses at the E^2 -page, and since $(\mathrm{H}\mathbb{F}_p)_*(X) \neq 0$, we get that $K(n)(X) \neq 0$, i.e. X has type $< n$. \square

Proposition 1.4.8. \mathcal{C}_n is a prime ideal.

Proof. For $X, Y \in \mathrm{Sp}_{(p)}^{\mathrm{fin}}$, by Kunneth we have

$$K(n-1)_*(X \otimes Y) = K(n-1)_*(X) \otimes K(n-1)_*(Y).$$

Assume that $X \in \mathcal{C}_n$, that is $K(n-1)_*(X) = 0$. It follows that $K(n-1)_*(X \otimes Y) = 0$, i.e. $X \otimes Y \in \mathcal{C}_n$, so \mathcal{C}_n is an ideal. Assume that $X \otimes Y \in \mathcal{C}_n$, that is $K(n-1)_*(X \otimes Y) = 0$, therefore one of the terms in the RHS of the equation must vanish (since they are graded vector spaces), so \mathcal{C}_n is a prime ideal. \square

Theorem 1.4.9 (Thick Subcategory Theorem [HS98]). *If \mathcal{T} is a proper thick subcategory of $\mathrm{Sp}_{(p)}^{\mathrm{fin}}$, then $\mathcal{T} = \mathcal{C}_n$ for some $0 \leq n \leq \infty$.*

Remark 1.4.10. The proof relies on a major theorem called the Nilpotence Theorem.

Corollary 1.4.11. $\mathrm{Spec} \mathbb{S}_{(p)} = \{\mathcal{C}_0, \mathcal{C}_1, \dots, \mathcal{C}_\infty\}$, and the closed subsets in the topology are chains $\{\mathcal{C}_k, \mathcal{C}_{k+1}, \dots, \mathcal{C}_\infty\}$ for some $0 \leq k \leq \infty$.

Proof. Follows immediately from the previous results. \square

We now want to describe $\mathrm{Spec} \mathbb{S}$. Note that the p -localization functor $L_{(p)}$ is a Bousfield localization. As such, it is left (its right adjoint is the inclusion), and in particular preserves cofibers. It also clearly reduced, i.e. sends 0 to 0. Now, $L_{(p)}$ is smashing, that is $L_{(p)}X = X \otimes \mathbb{S}_{(p)}$, so it is also symmetric monoidal. As we have seen in 1.1.8, under these conditions we can pullback primes. Since $L_{(p)}$ is smashing and $K(p, n)$ is p -local for every $0 \leq n \leq \infty$, we have that $K(n)_*(X_{(p)}) = K(n)_*(X)$. Therefore

$$\mathcal{P}_{p,n} = L_{(p)}^* \mathcal{C}_{p,n} = \left\{ X \in \mathrm{Sp}^{\mathrm{fin}} \mid K(p, n)_*(X) = 0 \right\}$$

and

$$\mathcal{P}_{p,\infty} = L_{(p)}^* \mathcal{C}_{p,\infty} = \left\{ X \in \mathrm{Sp}^{\mathrm{fin}} \mid X_{(p)} = 0 \right\}$$

are prime ideals. Moreover, by definition $K(p, 0) = \mathrm{H}\mathbb{Q}$, which implies that $\mathcal{P}_{p,0} = \left\{ X \in \mathrm{Sp}^{\mathrm{fin}} \mid \mathrm{H}\mathbb{Q}_*(X) = 0 \right\}$. We see that $\mathcal{P}_{p,0}$ is independent of p , and we denote it by $\mathrm{Sp}_{\mathrm{tor}}^{\mathrm{fin}}$.

Theorem 1.4.12 (TODO explain/reference). $\mathrm{Spec} \mathbb{S} = \left\{ \mathrm{Sp}_{\mathrm{tor}}^{\mathrm{fin}} \right\} \cup \left(\bigcup_p \{ \mathcal{P}_{p,1}, \dots, \mathcal{P}_{p,\infty} \} \right)$, and the closed subsets in the topology are finite unions of chains $\{ \mathcal{P}_{p,k}, \mathcal{P}_{p,k+1}, \dots, \mathcal{P}_{p,\infty} \}$ for some $0 \leq k \leq \infty$ (i.e. they may include $\mathrm{Sp}_{\mathrm{tor}}^{\mathrm{fin}}$). **TODO diagram**

TODO regarding the topology, maybe I should prove that the pullback is also continuous?

Remark 1.4.13. Thick subcategories are interesting for another reason, unrelated to the Balmer spectrum point of view, namely they give a very powerful proof method. Say we have a property that is satisfied by 0, and is closed under cofibers and retracts. It follows that the collection of objects that satisfy it is thick. Then, for example, by the thick subcategory theorem 1.4.9, it is enough to find one object in $\mathcal{C}_n \setminus \mathcal{C}_{n-1}$ that satisfies the property, to show that all objects in \mathcal{C}_n satisfy it.

1.5 The Stacky Point of View and the Relationship Between Morava K-Theory and Morava E-Theory

First we will describe, without being precise, another point of view on what chromatic homotopy theory is about.

There is a stack of formal groups with strict isomorphisms, denoted by $\mathcal{M}_{\mathrm{fg}}^s$. It can be described as the stack that sends a ring R to the groupoid of formal group laws, with strict isomorphisms between them. Quillen theorem 1.2.13 tells us that MU_* is the Lazard ring, that is the universal ring that carries the universal formal group law. It turns out that this theorem has a second part, which says that $(\mathrm{MU} \otimes \mathrm{MU})_*$ is the universal ring that carries two formal group laws and a strict isomorphism between them. Therefore, $\mathcal{M}_{\mathrm{fg}}^s$ is represented by $(\mathrm{MU}_*, (\mathrm{MU} \otimes \mathrm{MU})_*)$.

The geometric points of the stack $\mathcal{M}_{\mathrm{fg}}^s$ describe precisely the same as $\mathrm{Spec} \mathbb{S}$, that is because for an algebraically closed field of characteristic 0 there is a unique (up to isomorphism) formal group law which is of height 0 namely the additive formal group law, and for characteristic p there is a unique (up to isomorphism) formal group law of each height $1 \leq n \leq \infty$.

For a spectrum X , $\mathrm{MU}_*(X)$ is a $(\mathrm{MU}_*, (\mathrm{MU} \otimes \mathrm{MU})_*)$ -comodule, which is the same as a sheaf over $\mathcal{M}_{\mathrm{fg}}^s$. From this point of view, chromatic homotopy theory lets us study a spectrum by decomposing it over the stack $\mathcal{M}_{\mathrm{fg}}^s$.

We can restrict ourselves to the stack only over rings of characteristic p ,

$\mathcal{M}_{\text{fg},p}^s$, which is then represented by $\left((\text{MU}_{(p)})_*, (\text{MU}_{(p)} \otimes \text{MU}_{(p)})_*\right)$. **TODO I think it's true, is it?** Similarly to MU , BP is universal ring with the universal p -typical formal group law, and $(\text{BP} \otimes \text{BP})_*$ is the universal ring with two p -typical formal group laws and an isomorphism between them **TODO I didn't say this before**. Since every formal group law is isomorphic to a p -typical one, we know that the stack $\mathcal{M}_{\text{fg},p}^s$ is also represented by $(\text{BP}_*, (\text{BP} \otimes \text{BP})_*)$.

It is now reasonable that $K(n)$, obtained from BP by killing the v_m 's for $m \neq n$ and inverting v_n , sees the n -th level, and that $E(n)$ obtained in the same way but only killing v_m for $m > n$, sees the levels $\leq n$.

Let us now claim a precise statement, formalizing this description.

Theorem 1.5.1 (TODO reference). *$E(n)$ and $K(0) \oplus \cdots \oplus K(n)$ are Bousfield equivalent. That is, they have the same acyclics, locals, and their localization functors are the same.*

TODO chromatic square and chromatic tower, maybe another subsection?

1.6 Landweber Exact Functor Theorem

As we have seen, a complex orientation on a cohomology theory, which is described by a map $\text{MU} \rightarrow E$, has an associated formal group law, which is described by the map $L = \text{MU}_* \rightarrow E_*$. Note that this formal group law is of degree -2 , by virtue of the grading on $L = \text{MU}_*$. **TODO say more?** One can ask whether the converse is true, namely given a graded ring R and a formal group law F of degree -2 given by $L \rightarrow R$, is there a complex oriented cohomology theory whose coefficients are R and the associated formal group law is F .

A strategy is to define $(E_{R,F})_*(X) = \text{MU}_*(X) \otimes_{\text{MU}_*} R$. Unfortunately, this is not always a homology theory. However there is a condition that one can check, which guarantees that it is.

Definition 1.6.1. $L \rightarrow R$ is called *Landweber flat* if for every prime p , the image of the sequence $p = v_0, v_1, v_2, \dots$ in R , which are the coefficients of the p -series, is regular. That is, for each p and $n \geq 0$, v_n is not a zero divisor in $R/(v_0, v_1, \dots, v_{n-1})$.

Remark 1.6.2. If p is invertible in R , then p is invertible, and R/p is already 0, so we don't need to check v_1, v_2, \dots .

Theorem 1.6.3 (Landweber Exact Functor Theorem (LEFT), [Lur10, 15, 16]). *If $L \rightarrow R$ is Landweber flat, then $E_{R,F}$ defined above is a homology theory. Moreover, there are no phantom maps between such spectra, so $E_{R,F}$ is represented by a spectrum. This spectrum is complex oriented, has coefficients R and associated formal group law F .*

Example 1.6.4. Morava E-theory is Landweber flat, since by 1.3.6, the p -series has coefficients $p = v_0, v_1, \dots, v_n$. p is not a zero divisor in $E(n)_* = \mathbb{Z}_{(p)}[v_1, \dots, v_{n-1}, v_n^{\pm 1}]$. Then v_i is not a zero divisor in $E(n)_* / (p, v_1, \dots, v_{i-1}) \cong \mathbb{F}_p[v_i, \dots, v_{n-1}, v_n^{\pm 1}]$. After v_n the ring becomes 0 and we are done. For other primes, by 1.6.2 we are done.

Example 1.6.5. Morava K-theory $K(n)$ for $n > 0$ is not Landweber flat since $p = v_0$ is 0 in $K(n)_* = \mathbb{F}_p[v_n^{\pm 1}]$.

Example 1.6.6. $\mathbb{H}\mathbb{Z}$ is not Landweber flat since although $p = v_0$ is invertible, as we have seen in 1.2.10 the p -series is px , so v_1 is 0 in $\mathbb{Z}/p = \mathbb{F}_p$.

We can also ask the following question: given complex oriented cohomology theory $\text{MU} \rightarrow E$, such that $L \rightarrow E_*$ is Landweber flat, is $E_{R,F}$ equivalent to E ? The answer is yes, at least in some cases.

Theorem 1.6.7. *Let E be as above, which is also evenly graded (i.e. E_* is an evenly graded ring), then there is an equivalence $E_{R,F} \rightarrow E$.*

Proof. This is a slight variation on [Lur10, 18, proposition 11]. First note that for every spectrum X we have $\text{MU} \otimes X \rightarrow E \otimes X$, which induces $\text{MU}_*(X) \rightarrow E_*(X)$, a map of MU_* -modules. Moreover, since $E_* \rightarrow E_*(X)$ is a map of E_* -module, the map $\text{MU}_* \rightarrow E_*$ makes it a map of MU_* -modules. Together this gives a map $(E_{R,F})_*(X) = \text{MU}_*(X) \otimes_{\text{MU}_*} E_* \rightarrow E_*(X)$. This map is a map of homology theories. **TODO should I explain why?** By [Lur10, 17, theorem 6] **TODO does it follow?**, this map lifts to a map of spectra $E_{R,F} \rightarrow E$. Since by construction when $X = \mathbb{S}$ the map above is $E_* \rightarrow E_*$ which is an isomorphism, we see that the map $E_{R,F} \rightarrow E$ is an equivalence. \square

Example 1.6.8. We return to complex K-theory, from 1.2.3 and 1.2.11. We can take the completion at the element $p \in K_*$, which gives the spectrum K_p^\wedge . This spectrum has coefficients $(K_p^\wedge)_* = (K_*)_p^\wedge = (\mathbb{Z}[\beta^{\pm 1}])_p^\wedge = \mathbb{Z}_p[\beta^{\pm 1}]$.

The formal group law, as we have seen, is given by $F_{K_p^\wedge}(y, z) = y + z + \beta yz$. We claim that $F_{K_p^\wedge}/K_p^\wedge$ is Landweber flat. Clearly $p = v_0$ is not a zero divisor in $\mathbb{Z}_p[\beta^{\pm 1}]$. As we have seen in 1.2.12, mod- p the p -series is $\beta^{p-1}x^p$, so that $v_1 = \beta^{p-1}$ which is not a zero divisor $\mathbb{F}_p[\beta^{\pm 1}]$. Modulo v_1 the ring is already 0, and we are done. For other primes, by 1.6.2 we are done. Therefore, by 1.6.7 we get that $K_p^\wedge \cong E_{K_p^\wedge, F_{K_p^\wedge}}$.

1.7 Lubin-Tate Deformation Theory

The Morava E-theory we have considered until now $E(n)$, also called Johnson-Wilson spectrum was constructed from BP. As we noted, it is Landweber flat, which indicates that there is another approach to constructing it. Indeed there is a way to construct a related spectrum, which will be called the Lubin-Tate spectrum.

To that end, we first define the category CompRing as the category of complete local rings. The objects are complete local rings (R, \mathfrak{m}) , we also denote by $\pi : R \rightarrow R/\mathfrak{m}$ the projection. Morphisms $\varphi : (R, \mathfrak{m}) \rightarrow (S, \mathfrak{n})$ are local homomorphisms, i.e. a homomorphism $\varphi : R \rightarrow S$ s.t. $\varphi(\mathfrak{m}) \subseteq \mathfrak{n}$. In particular it induces a homomorphism $\varphi/\mathfrak{m} : R/\mathfrak{m} \rightarrow S/\mathfrak{n}$, which satisfies $\varphi/\mathfrak{m} \circ \pi_R = \pi_S \circ \varphi$.

We fix k be a perfect field of characteristic p (i.e. the Frobenius is an isomorphism), and Γ a formal group law over k of height $n < \infty$. Lubin and Tate [LT66] considered a moduli problem associated to Γ/k , described by a functor $\text{Def} : \text{CompRing} \rightarrow \text{Grpds}$.

Definition 1.7.1. Let (R, \mathfrak{m}) be a complete local ring and denote by $\pi : R \rightarrow R/\mathfrak{m}$ the quotient. A *deformation* of Γ/k to (R, \mathfrak{m}) , is (G, i) , where G is a formal group law over R , $i : k \rightarrow R/\mathfrak{m}$ is a homomorphism of fields, such that $i^*\Gamma = \pi^*G$. A \star -*isomorphism* between two deformations to (R, \mathfrak{m}) , $f : (G_1, i_1) \rightarrow (G_2, i_2)$, is defined only when $i_1 = i_2$, and consists of an isomorphism $f : G_1 \rightarrow G_2$, such that $\pi^*f : i^*\Gamma = \pi^*G_1 \rightarrow \pi^*G_2 \rightarrow i^*\Gamma$ is the identity, i.e. $f(x) = x \bmod \mathfrak{m}$. These assemble to a groupoid $\text{Def}(R, \mathfrak{m})$, whose objects are deformations to (R, \mathfrak{m}) , and morphisms are \star -isomorphisms.

Remark 1.7.2. $\text{Def}(R, \mathfrak{m})$ can be seen as the pullback of the groupoids $\text{FGL}(R)$ and $\coprod_{i:k \rightarrow R/\mathfrak{m}} \{\Gamma\}$ over $\text{FGL}(R/\mathfrak{m})$ (where the maps are $G \mapsto q^*G$ and $i \mapsto i^*\Gamma$ respectively).

Proposition 1.7.3 (/definition). *The construction $\text{Def}(R, \mathfrak{m})$ is functorial.*

Proof. Let $\varphi : (R, \mathfrak{m}) \rightarrow (S, \mathfrak{n})$ be a local homomorphism.

For a deformation (G, i) to (R, \mathfrak{m}) , we define $\text{Def}(\varphi)(G, i) = (\varphi^*G, \varphi/\mathfrak{m} \circ i)$. Note that φ^*G is a formal group law over S , and $\varphi/\mathfrak{m} \circ i : k \rightarrow R/\mathfrak{m} \rightarrow S/\mathfrak{n}$ is a homomorphism. Moreover, $(\varphi/\mathfrak{m} \circ i)^* \Gamma = (\varphi/\mathfrak{m})^* i^* \Gamma = (\varphi/\mathfrak{m})^* \pi_R^* G = (\varphi/\mathfrak{m} \circ \pi_R)^* G = (\pi_S \circ \varphi)^* G = \pi_S^* \varphi^* G$, which shows that $\text{Def}(\varphi)(G, i)$ is a deformation to (S, \mathfrak{n}) .

For a \star -isomorphism $f : (G_1, i) \rightarrow (G_2, i)$, which is the data of an isomorphism $f : G_1 \rightarrow G_2$ such that $\pi_R^* f = \text{id}_{i^* \Gamma}$ is the identity, we need to define a \star -isomorphism $\text{Def}(\varphi)(G, i_1) \rightarrow \text{Def}(\varphi)(G, i_2)$. Take it to be the isomorphism $\varphi^* f : \varphi^* G_1 \rightarrow \varphi^* G_2$, which satisfies $\pi_S^* \varphi^* f = (\varphi/\mathfrak{m})^* \pi_R^* f = (\varphi/\mathfrak{m})^* \text{id}_{i^* \Gamma} = \text{id}_{(\varphi/\mathfrak{m})^* i^* \Gamma} = \text{id}_{(\varphi/\mathfrak{m} \circ i)^* \Gamma}$. The identity $\text{id}_G : (G, i) \rightarrow (G, i)$ is clearly sent to $\text{id}_{\varphi^* G}$, and compositions are sent to compositions.

This shows that $\text{Def}(\varphi) : \text{Def}(R, \mathfrak{m}) \rightarrow \text{Def}(S, \mathfrak{n})$ is indeed a functor. Moreover, it is clear that $\text{Def}(\text{id}_R)$ is the identity and compositions are sent to compositions, which shows that $\text{Def} : \text{CompRing} \rightarrow \text{Grpds}$ is indeed a functor. \square

Remark 1.7.4. We recall quickly that the Witt vectors Wk is a ring of characteristic 0, with maximal ideal (p) , and residue field $Wk/p \cong k$. For example, $W\mathbb{F}_p = \mathbb{Z}_p$.

Theorem 1.7.5 ([Rez98, 4.4, 5.10], originally due to [LT66]). *The functor Def lands in discrete groupoids (i.e. $\text{Def}(R, \mathfrak{m})$ has 0 or 1 morphisms between objects). Furthermore the functor Def is co-represented, that is there exists a universal deformation, and the complete local ring can be chosen (non-canonically) to be $Wk[[u_1, \dots, u_{n-1}]]$.*

Let us unravel what that means. First note that the quotient of $Wk[[u_1, \dots, u_{n-1}]]$ by the maximal ideal (p, u_1, \dots, u_{n-1}) is k . The universal deformation can be chosen such that the formal group law over it Γ_U over $Wk[[u_1, \dots, u_{n-1}]]$ satisfies $\pi^* \Gamma_U$ is Γ . The universality means that for (R, \mathfrak{m}) , the assignment

$$\text{hom}_{\text{CompRing}}(Wk[[u_1, \dots, u_{n-1}]], R) \rightarrow \text{Def}(R, \mathfrak{m}), \quad \varphi \mapsto \varphi^* \Gamma_U$$

is an equivalence.

Now, we can form the graded ring $Wk[[u_1, \dots, u_{n-1}]] [u^{\pm 1}]$ where $|u| = 2$. We can define the formal group law $(u\Gamma_U)(x, y) = u^{-1}\Gamma_U(uy, uz)$, which is of degree -2 , thus we get a map $L \rightarrow Wk[[u_1, \dots, u_{n-1}]] [u^{\pm 1}]$.

Proposition 1.7.6 ([Rez98, 6.9]). $L \rightarrow Wk[[u_1, \dots, u_{n-1}]] [u^{\pm 1}]$ is Landweber flat.

Using LEFT 1.6.3, we immediately get:

Corollary 1.7.7. *There is a complex oriented cohomology theory $E(k, \Gamma) = E_{Wk[[u_1, \dots, u_{n-1}]] [u^{\pm 1}], u\Gamma_U}$, called Lubin-Tate spectrum, with coefficients $E(k, \Gamma)_* = Wk[[u_1, \dots, u_{n-1}]] [u^{\pm 1}]$ and associated formal group law $u\Gamma_U$.*

Example 1.7.8. We continue the complex K-theory saga from 1.6.8. Take the field $k = \mathbb{F}_p$ and the formal group law $\Gamma(y, z) = y + z + yz$, of height $n = 1$. By the above construction, the ring of the universal deformation is $W\mathbb{F}_p = \mathbb{Z}_p$. The universal formal group law of the universal deformation can be taken to be $\Gamma_U(y, z) = y + z + yz$ (this follows from the proof at [Rez98, 5.10], since here $n = 1$ so there are no u_i 's). We look at the ring $\mathbb{Z}_p[u^{\pm 1}]$, and at the formal group law over it $(u\Gamma_U)(y, z) = u^{-1}(uy + uz + u^2yz) = y + z + uyz$. It is then clear that the isomorphism $\mathbb{Z}_p[u^{\pm 1}] \rightarrow \mathbb{Z}_p[\beta^{\pm 1}]$, sends $u\Gamma_U$ to $F_{K_p^\wedge}$. It follows that $K_p^\wedge \cong E_{K_p^\wedge, F_{K_p^\wedge}} \cong E(\mathbb{F}_p, \Gamma)$, i.e. p -complete K-theory K_p^\wedge is a Lubin-Tate spectrum at height 1.

This concludes the construction of the Lubin-Tate variant of Morava E-theory. To compare this version with the previous, we define another variant of Morava E-theory.

Definition 1.7.9. Let $I \leq E(n)_* = \mathbb{Z}_{(p)}[v_1, \dots, v_{n-1}, v_n^{\pm 1}]$ be the ideal $I = (p = v_0, v_1, \dots, v_{n-1})$. We define the spectrum $\widehat{E(n)} = E(n)_I^\wedge$, called completed Johnson-Wilson spectrum.

Theorem 1.7.10. *The following three forms of Morava E-theory are Bousfield equivalent:*

1. Johnson-Wilson $E(n)$,
2. completed Johnson-Wilson $\widehat{E(n)}$,
3. Lubin-Tate $E(k, \Gamma)$.

TODO reference Hovey Strickland Morava K-Theories and Localization 5.3

2 Atiyah-Segal

We now leave the realm of chromatic homotopy theory. One aspect of algebraic topology is to try to capture properties of spaces using algebraic invariants. One of the most fruitful such invariants is complex K-theory, denoted K , and one of the most important spaces in homotopy theory is BG , so it is natural to ask for a description of $K(BG)$ (by Bott periodicity, we will consider only $K = K^0$). Atiyah and Segal [AS69] gave a description of this, and more, in the case that G is a compact Lie group, in terms of representations.

From now we fix a compact Lie group G . Also, a representation means a finite dimensional unitary representation. We should also note that beyond this part, we will be mostly interested in finite groups.

2.1 The Atiyah-Segal Theorem

We denote by $R(G)$ the *representation ring* of G , that is the collection of virtual representations of G (which can be written as a formal difference $V - U$) up to isomorphism, where the addition is given by direct sum and the product is given by tensor product. This is an augmented ring $\varepsilon : R(G) \rightarrow \mathbb{Z}$ by the virtual dimension (i.e. $\varepsilon(V - U) = \dim(V - U) = \dim V - \dim U$). The *augmentation ideal* is $I = \ker \varepsilon = \{V - U \in R(G) \mid 0 = \dim(V - U)\}$.

Atiyah and Segal showed that one can describe $K(BG)$ in these terms, namely, it is the completion of $R(G)$ at the ideal I :

Theorem 2.1.1 ([AS69]). $K(BG) \cong R(G)_I^\wedge$.

We will not prove the theorem, but we will indicate some of the key ingredients.

First of all, to show that objects are isomorphic, we need a map. Before giving the map actually used in the proof, we describe an easier way to see where this map comes from. Recall that $K(X) \cong [X, BU \times \mathbb{Z}]$. The data of a representation of G is the same thing as a homomorphism $G \rightarrow U(n)$. Since B is a functor, we get a map $BG \rightarrow BU(n)$, and by composing with the injection $BU(n) \cong BU(n) \times \{n\} \rightarrow BU \times \mathbb{Z}$, we get a map $BG \rightarrow BU \times \mathbb{Z}$, that is, an element of $K(BG)$. Therefore we get a map $R(G) \rightarrow K(BG)$. The theorem shows that it is a ring homomorphism which exhibits $K(BG)$ as the completion of $R(G)$ at I .

There is an alternative description of this map. In [Seg68], Segal described a variant of K theory, called equivariant K-theory K_G . This variant assigns to a G -space the ring of virtual G -bundles, that is, bundles equipped with an action of G which is compatible with the action on the base G -space. Note that K_G is no longer homotopy invariant, since it also takes into account the action of G . First we note the following:

Proposition 2.1.2. $K_G(*) = R(G)$ (where $*$ is equipped with a trivial G action).

Proof. This follows from the definitions, since a vector bundle over a point is just a vector space, and it is equipped with a G action over the point, which is just a G representation. \square

For any G -space X , the projection map $\text{pr} : X \rightarrow X/G$ allows us to pullback vector bundles on X/G to G -bundles on X . In other words, it induces a map $\text{pr}^* : K(X/G) \rightarrow K_G(X)$.

Proposition 2.1.3 ([Seg68, 2.1]). *Suppose the action of G on X is free. Then pr^* admits an inverse, given by taking a bundle $E \rightarrow X$ to $E/G \rightarrow X/G$. In particular, $K(X/G) \cong K_G(X)$.*

Now, we have a map of G -spaces given by $EG \rightarrow *$. By the above we get:

$$R(G) \cong K_G(*) \rightarrow K_G(EG) \cong K(EG/G) = K(BG)$$

It can be shown that this is the same map $R(G) \rightarrow K(BG)$ described before, which exhibits $K(BG)$ as the I -completion of $R(G)$. Atiyah and Segal use this map and variants to prove the theorem.

Here is a sketch of the proof given by Atiyah and Segal. First of all, we note that the theorem is proven for the entire K^* rather than just for $K = K^0$. Also note that $R^*(G) = K_G^*(*)$ is a 2-periodic version of the representation ring (because K_G^* also satisfies Bott periodicity). We have the corresponding 2-periodic version of the augmentation ideal, which we denote by I^* . They use the Milnor join construction $EG_n = \underbrace{G * \cdots * G}_{n \text{ times}}$

and $BG_n = EG_n/G$, which has the property that $\text{colim } EG_n \rightarrow \text{colim } BG_n$ is a model for $EG \rightarrow BG$. Then, for any compact G -space X there is a similar map to the map above: using $X \times EG_n \rightarrow X$ we get a map $K_G^*(X) \rightarrow K_G^*(X \times EG_n)$. All of these are $R^*(G)$ -modules, and Atiyah

and Segal show that this map factors through the quotient by $(I^*)^n$, to give a map $K_G^*(X) / (I^*)^n \rightarrow K_G^*(X \times EG_n)$. The two sides assemble into pro-rings, and the maps assemble to a map between the pro-rings:

$$\{K_G^*(X) / (I^*)^n\}_n \rightarrow \{K_G^*(X \times EG_n)\}_n$$

What they actually prove is the strong form:

Theorem 2.1.4 ([AS69]). *If $K_G^*(X)$ is finite over $R^*(G)$, then the above map of pro-rings is an isomorphism.*

Their proof has another interesting aspect. Although it is a statement about the K_G of some class of G -spaces, for one specific group G , their proof involves the K_G 's of several groups. In particular, to prove the result for example for a finite group, their proof involves more general compact Lie groups. The proof consists of four steps. In every step we show the theorem holds for a more general type of group:

- $G = U(1)$ (circle group),
- $G = U(1)^n$ (torus group),
- $G = U(n)$,
- G a general compact Lie group (this step is proven by G embedding in $U(n)$).

We note that the first formulation of the Atiyah-Segal theorem 2.1.1, is indeed a private case of the second formulation 2.1.4. Take the case $X = *$. By definition, $K_G^*(*)$ is finite over $R^*(G) = K_G^*(*)$, so 2.1.4 holds and we have an isomorphism of pro-rings $\{K_G^*(*) / (I^*)^n\}_n \rightarrow \{K_G^*(EG_n)\}_n$. In particular, after computing the limits $\lim : \text{Pro Ring} \rightarrow \text{Ring}$, we get an isomorphism of rings $K_G^*(*)_{I^*}^\wedge \rightarrow K_G^*(EG)$. Taking only the 0-th cohomology gives the desired isomorphism:

$$R(G)_I^\wedge \cong K_G^*(*)_I^\wedge \rightarrow K_G^*(EG) \cong K(EG/G) = K(BG)$$

2.2 Examples

We compute a few examples in detail, to make the isomorphism more vivid.

2.2.1 $U(1)$, the Circle Group

Take $G = U(1)$, the circle group. It is known that the irreducible representations are of dimension 1 and labeled by an integer $m \in \mathbb{Z}$, i.e. they are homomorphisms $\rho_m : U(1) \rightarrow U(1)$ given by $\rho_m(e^{i\theta}) = e^{mi\theta}$. In particular, $\rho_0 = 1$ is the trivial representation. It is then clear that for $m \geq 0$, $\rho_1^{\otimes m} = \rho_m$ and $\rho_{-1}^{\otimes -m} = \rho_{-m}$. Therefore the representation ring generated under (virtual) direct sums and tensor products by ρ_1 and ρ_{-1} . Moreover, $\rho_1 \otimes \rho_{-1} = 1$. Therefore we conclude that $R(U(1)) = \mathbb{Z}[\rho, \rho^{-1}]$.

The augmentation map is the homomorphism $\varepsilon : R(U(1)) \rightarrow \mathbb{Z}$ which sends $1, \rho$ and ρ^{-1} to 1. Recall that the augmentation ideal is $I = \ker \varepsilon$. We set $t = \rho - 1$, which clearly belongs to I . We can also write then $R(U(1)) = \mathbb{Z}[t, (1+t)^{-1}]$. Note that ε factors to a map $R(U(1))/I \rightarrow \mathbb{Z}$, which is already an isomorphism, so by the first isomorphism theorem indeed $I = (t)$.

We compute the completion $R(U(1))_I^\wedge$. Note that in $\mathbb{Z}[t]/t^n$, $1+t$ is already invertible. The reason is that the formal power series for the inverse is finite since large enough powers of t are zero, $\frac{1}{1-(-t)} = \sum_{m=0}^{n-1} (-t)^m$ is an inverse. Therefore we see that $R(U(1))/I^n \cong \mathbb{Z}[t]/t^n$, and clearly the maps in the limit diagram send t to t . We get that $R(U(1))_I^\wedge = \lim \mathbb{Z}[t]/t^n \cong \mathbb{Z}[[t]]$.

In [Hat17, proposition 2.24], it is shown that $K(\mathbb{CP}^n) \cong \mathbb{Z}[L]/(L-1)^{n+1}$, where L is the canonical line bundle on \mathbb{CP}^n . In 1.2.3 we denoted $t = L - 1$ (warning: there we looked at K^* , now we focus on K), which allows us to rewrite this as $K(\mathbb{CP}^n) \cong \mathbb{Z}[t]/t^{n+1}$. As we noted, the limit is $K(\mathbb{CP}^\infty) \cong \mathbb{Z}[[t]]$. We thus see that $K(BU(1)) \cong \mathbb{Z}[[t]]$ where $t = L - 1$ is the canonical line bundle minus 1.

The identity map $\rho : U(1) \rightarrow U(1)$, is mapped to the identity $BU(1) \rightarrow BU(1)$ (by functoriality of B), which tautologically corresponds the universal line bundle L on $BU(1)$. We therefore see that the Atiyah-Segal map $R(U(1)) \rightarrow K(BU(1))$, sends ρ to L , and therefore $t = \rho - 1$ to $t = L - 1$. Which indeed shows that the map admits $K(BU(1)) \cong \mathbb{Z}[[t]]$ as the $I = (t)$ -completion of $R(U(1))$.

2.2.2 $\mathbb{Z}/2$, Cyclic Group of Order 2

Take $G = \mathbb{Z}/2$. Here we have only two irreducible representations, the trivial, and $\rho(0) = 1, \rho(1) = -1$. Also, it is clear that $\rho \otimes \rho$ is the trivial

representation. Therefore, $R(\mathbb{Z}/2) = \mathbb{Z}[\rho]/(\rho^2 - 1)$. Similarly to before, the augmentation $\varepsilon : R(\mathbb{Z}/2) \rightarrow \mathbb{Z}$ sends $1, \rho$ to 1, so clearly $(\rho - 1) \subseteq I$, and for the same reasoning as in the previous example this is actually an equality. We change coordinates to $t = \rho - 1$, and we have $R(\mathbb{Z}/2) = \mathbb{Z}[t]/(t^2 + 2t)$, and $I = (t)$.

We move to computing the completion $R(\mathbb{Z}/2)_I^\wedge$. Modulo $t^2 + 2t$, i.e. $t^2 = -2t$, we have that $t^n = (-2)^{n-1}t$. Thus $I^n = ((-2)^{n-1}t) = (2^{n-1}t)$, so $R(\mathbb{Z}/2)/I^n = \mathbb{Z}[t]/(t^2 + 2t, 2^{n-1}t)$. We will not describe the ring structure on the completion, but since the forgetful functor from rings to abelian groups is right adjoint, to compute the completion as an abelian group, we can compute the limit in abelian groups. As an abelian group, $R(\mathbb{Z}/2)/I^n$ is isomorphic to $\mathbb{Z} \oplus \mathbb{Z}/2^{n-1}$. It is then clear that as an abelian group, $R(\mathbb{Z}/2)_I^\wedge$ is isomorphic to $\mathbb{Z} \oplus \mathbb{Z}_2$.

Using the Atiyah-Segal theorem we conclude that as abelian groups

$$K(\mathbb{RP}^\infty) = K(B\mathbb{Z}/2) \cong \mathbb{Z} \oplus \mathbb{Z}_2.$$

2.3 Recollections from Character Theory

We restrict ourselves to the case of finite groups G . We recall that representations of groups can be studied by their characters. Specifically the character map $\chi : R(G) \rightarrow \mathbb{Z}[\chi_{\rho_i}]$, defined by $\chi_\rho = \text{tr } \rho$, is an isomorphism, where the ring on the right is the ring of functions generated by the irreducible characters (the multiplication of two characters is a character so it is indeed closed under multiplication).

We also recall that characters are class functions, that is, they are constant on conjugacy classes. Let L be some field containing all the values of all characters. Then a natural place to study characters is in the ring of class functions with values in L , denote by $\text{Cl}(G; L)$. Let us phrase this in a way that will be useful in the next section. G is equipped with a G action by conjugation, $\gamma \cdot g = \gamma^{-1}g\gamma$. Equip L with the trivial G action. Then $\text{Cl}(G; L) = \text{hom}_{G\text{Set}}(G, L)$.

We can of course extend the range of the character map to get an injection $\chi : R(G) \rightarrow \text{Cl}(G; L)$. The first classical theorem regarding the relationship between characters and class functions is:

Theorem 2.3.1. *After tensoring with L , the character map $\chi \otimes L : R(G) \otimes L \rightarrow \text{Cl}(G; L)$ becomes an isomorphism.*

Proof. Similarly to the proof in [Ser77, 9.1] for $L = \mathbb{C}$, we can view $\text{Cl}(G; L)$ as a vector space over L , and the characters are linearly independent, so by counting them we see that the image of $\chi \otimes L$ has the dimension of the whole vector space and we are done. \square

By definition the value of a character is the trace of a linear transformation $\chi_\rho(g) = \text{tr } \rho(g) = \sum \lambda_i$ where λ_i are the eigenvalues (which exist since the representation is unitary). Since $g^{|G|} = e$, we get $\rho(g^{|G|}) = \rho(e) = \text{id}$, but then we get that the eigenvalues of $\rho(g^{|G|})$ are on the one hand $\lambda_i^{|G|}$ and on the other hand they are all 1, which means that all the eigenvalues are roots of unity. Therefore $L = \mathbb{Q}^{\text{ab}} = \mathbb{Q}(\zeta_\infty)$ is always a valid choice for L (regardless of G). To be concrete, we will take this choice.

The Galois group of \mathbb{Q}^{ab} is $\text{Gal}(\mathbb{Q}^{\text{ab}}/\mathbb{Q}) \cong \hat{\mathbb{Z}}^\times$. For every $m \in \hat{\mathbb{Z}}^\times$ we also denote by $m \in \text{Gal}(\mathbb{Q}^{\text{ab}}/\mathbb{Q})$ the corresponding element, which can be described as the homomorphism which raises a root of unity to the power of m . Similarly it acts on G by taking g to g^m . Then, for every such m and g we have that $\chi_\rho(g^m) = \text{tr } \rho(g) = \sum \lambda_i^m = m.(\sum \lambda_i) = m.(\chi_\rho(g))$. We replace g with $g^{m^{-1}}$ (m is invertible), and rewrite this as $\chi_\rho(g) = m.(\chi_\rho(g^{m^{-1}}))$. Similarly to this equality, we can define an action of $\text{Gal}(\mathbb{Q}^{\text{ab}}/\mathbb{Q})$ on $\text{Cl}(G; \mathbb{Q}^{\text{ab}})$, by taking a class function f to $m.f$ defined by $(m.f)(g) = m.(f(g^{m^{-1}}))$.

Let us rewrite this action in another way, which will be helpful in the next section. We note that $G \cong \text{hom}_{\text{TopGrp}}(\hat{\mathbb{Z}}, G)$ (continuous homomorphisms). We get an action of $\text{Aut}(\hat{\mathbb{Z}}) \cong \hat{\mathbb{Z}}^\times$ on G by pre-composition. Concretely, $m \in \hat{\mathbb{Z}}^\times$ acts by sending $g \in G$ to g^m . Since $\hat{\mathbb{Z}}^\times$ acts on \mathbb{Q}^{ab} , we get an action on $\text{Cl}(G; \mathbb{Q}^{\text{ab}}) = \text{hom}_{G\text{Set}}(G, \mathbb{Q}^{\text{ab}})$ by acting with m^{-1} in the source and with m in the target. It is evident that this is the same action from the previous paragraph.

As we just saw, the characters are in the fixed points $\text{Cl}(G; \mathbb{Q}^{\text{ab}})^{\text{Gal}(\mathbb{Q}^{\text{ab}}/\mathbb{Q})}$. Also, since the rationals are fixed by the action of the Galois group, rational linear combinations of characters are in the fixed points. We therefore conclude that the character map after tensoring with \mathbb{Q} lands in the fixed points, i.e. $\chi \otimes \mathbb{Q} : \text{R}(G) \otimes \mathbb{Q} \rightarrow \text{Cl}(G; \mathbb{Q}^{\text{ab}})^{\text{Gal}(\mathbb{Q}^{\text{ab}}/\mathbb{Q})}$. Moreover, the second classical theorem is:

Theorem 2.3.2 ([Ser77, Theorem 25]). *The map $\chi \otimes \mathbb{Q} : \text{R}(G) \otimes \mathbb{Q} \rightarrow$*

$\mathrm{Cl}(G; \mathbb{Q}^{\mathrm{ab}})^{\mathrm{Gal}(\mathbb{Q}^{\mathrm{ab}}/\mathbb{Q})}$ is an isomorphism.

To conclude, 2.3.1 tells us that $R(G) \otimes \mathbb{Q}^{\mathrm{ab}} \cong \mathrm{Cl}(G; \mathbb{Q}^{\mathrm{ab}})$, and 2.3.2 tells us that $R(G) \otimes \mathbb{Q} \cong \mathrm{Cl}(G; \mathbb{Q}^{\mathrm{ab}})^{\mathrm{Gal}(\mathbb{Q}^{\mathrm{ab}}/\mathbb{Q})}$.

3 HKR Character Theory

As we have seen in the previous section, Atiyah and Segal gave a description of $K(BG)$ in terms of the representation ring. We have also seen in the section on chromatic homotopy theory, that complex K-theory is related to Morava K-theory at height 1 by 1.3.11, and to Morava E-theory at height 1 by 1.7.8. Representations can be studied using their characters, and one may wonder if a similar construction can be used to study higher analogues of complex K-theory evaluated at BG .

Hopkins, Kuhn and Ravenel showed in [HKR00] that it is indeed possible. Their paper contains a lot of results, but we will concentrate on theorem C. Fix some finite group G . Similarly to the proof Atiyah-Segal theorem, the actual proof of theorem C involves a general construction, even to prove the specific case we are interested, but it will be easier to state it first for the specific case. Let $E = E(k, \Gamma)$ be the Lubin-Tate spectrum from 1.7.7, for some field k of characteristic p , and Γ a formal group law over k of height n . **TODO they write it specifically for $k = \mathbb{F}_{p^n}$, but we don't really need that, right?** There is some ring $L = L(E^*)$ (which depends on the spectrum E). It is then possible define some generalized characters $\mathrm{Cl}_{n,p}(G; L)$, which are completely algebraic and combinatorial (besides the definition of the ring L). Lastly, there is a character map $\chi_{n,p}^G : E^*(BG) \rightarrow \mathrm{Cl}_{n,p}(G; L)$. This character map has similar formal properties to the ordinary character map, namely, similarly to 2.3.1, after tensoring with L , the character map

$$\chi_{n,p}^G \otimes L : E^*(BG) \otimes L \rightarrow \mathrm{Cl}_{n,p}(G; L)$$

becomes an isomorphism. Similarly to 2.3.2, there is an action of $\mathrm{Aut}(\mathbb{Z}_p^n) \cong (\mathbb{Z}_p^\times)^n$ on $\mathrm{Cl}_{n,p}(G; L)$, and it turns out that the character map lands in the fixed points. Moreover, we can merely rationalize, which is given by inverting p , the source, rather tensoring with L . It turns out that after rationalization and restricting the codomain to the fixed points, the map

becomes an isomorphism, that is

$$p^{-1}\chi_{n,p}^G : p^{-1}E^*(BG) \rightarrow \text{Cl}_{n,p}(G; L)^{\text{Aut}(\mathbb{Z}_p^n)}$$

is an isomorphism.

TODO write something about the structure of this section

3.1 Towards a Definition of the Character Map

Following [HKR00], we denote by $\Lambda_r = (\mathbb{Z}/p^r)^n$ and $\Lambda = \mathbb{Z}_p^n$.

An element $g \in G$ is called *p-power-torsion* if $g^{p^a} = e$ for some a . Note that a conjugation of a *p-power-torsion* element is again *p-power-torsion*. We also define $r_0 \in \mathbb{N}$ to be the minimal r s.t. every *p-power-torsion* element g is p^{r_0} -torsion, i.e. satisfies $g^{p^{r_0}} = e$.

Consider the set $G_{n,p}$ of n -tuples (g_1, \dots, g_n) of commuting *p-power-torsion* elements. This set has an action of G by conjugating all the elements in a tuple by the same element, i.e. $\gamma \in G$ acts by $\gamma \cdot (g_1, \dots, g_n) = (\gamma^{-1}g_1\gamma, \dots, \gamma^{-1}g_n\gamma)$. Concretely, for $r \geq r_0$, we have $G_{n,p} = \text{hom}_{\text{Grp}}(\Lambda_r, G)$, with the G -action by conjugation at the values. In a similar fashion, $G_{n,p} = \text{hom}_{\text{TopGrp}}(\Lambda, G)$ (the homomorphisms are required to be continuous).

Let R be a ring. Equip it with the trivial G -action. We define the class functions $\text{Cl}_{n,p}(G; R) = \text{hom}_{G\text{Set}}(G_{n,p}, R)$, that is functions to R which are invariant under the G action. This is a ring, by defining the operations point-wise. Note that this is a purely algebraic/combinatorial construction, just a copy of R for every orbit of $G_{n,p}/G$, that is $\text{Cl}_{n,p}(G; R) \cong \bigoplus_{[\alpha] \in G_{n,p}/G} R$.

We would like to construct a character map $E^*(BG) \rightarrow \text{Cl}_{n,p}(G; R)$, for some R . We will try to unravel what this means, and find appropriate R 's at the same time. By the above, this is a homomorphism $E^*(BG) \rightarrow \bigoplus_{[\alpha] \in G_{n,p}/G} R$. That is, for every $[\alpha] \in G_{n,p}/G$ we need to provide a homomorphism $E^*(BG) \rightarrow R$. Choose a representative $\alpha \in G_{n,p} = \text{hom}_{\text{Grp}}(\Lambda_r, G)$ (for $r \geq r_0$). Since B is a functor we get $B\alpha : B\Lambda_r \rightarrow BG$, and then we can take E -cohomology to get a homomorphism $B\alpha^* : E^*(BG) \rightarrow E^*(B\Lambda_r)$. If we had a homomorphism $E^*(B\Lambda_r) \rightarrow R$, such that its composition with $B\alpha^*$ is independent of the choice of r and α , we would indeed get a well-defined character map.

3.2 The E -Cohomology of BA

We postpone the discussion of the rings, to give some properties of E -cohomology.

First we wish to describe the $E^*(B\mathbb{Z}/m)$. Let $\psi_m : \mathbb{Z}/m \rightarrow U(1)$ be the homomorphism $\psi(1) = e^{2\pi i/m}$. This induces $B\psi^* : E^*(BU(1)) \rightarrow E^*(B\mathbb{Z}/m)$. Denote by $x \in E^2(B\mathbb{Z}/m)$ the cohomology class $B\psi^*(x)$.

Proposition 3.2.1 ([HKR00, 5.8]). *The E^* -cohomology of $B\mathbb{Z}/m$ is given by $E^*(B\mathbb{Z}/m) = E^*[[x]] / ([m](x))$. Write $m = sp^t$ with s coprime to p , then this is a free E^* -module of rank p^{nt} .*

TODO I don't understand the proof, try to understand and write here

Proposition 3.2.2. *Let Y be a space s.t. $E^*(Y)$ is a free E^* -module of finite rank. Then Y satisfies Kunneth with respect to any X , that is, the map $E^*(X) \otimes_{E_*} E^*(Y) \rightarrow E^*(X \times Y)$ is an isomorphism.*

Proof. Look at the functors $X \mapsto E^*(X) \otimes E^*(Y)$ and $X \mapsto E^*(X \times Y)$. Both of them are manifestly homotopy invariant. Since $E^*(Y)$ is free, it is also flat, and so both functors satisfy Mayer-Vietoris. Both functors send arbitrary wedges to arbitrary products, since tensor with a free finite rank modules commutes with arbitrary products. We conclude that they are both cohomology theories. Moreover, they agree on $X = *$, and therefore are isomorphic. \square

Using both propositions we can bootstrap to arbitrary finite abelian groups.

Proposition 3.2.3 ([HKR00, 5.8]). *Let A be an abelian group, and write $|A| = sp^t$ for s coprime to p . Then $E^*(BA)$ is a free E^* -module of rank p^t , and BA satisfies Kunneth.*

Proof. A finite abelian group is the product of finite cyclic groups. Since B commutes with products, we can induct on the number of components in the product. \square

Recall that the formal group law on E_* was defined by taking the $E^*(B-)$ to the multiplication map $U(1) \times U(1) \rightarrow U(1)$. By pre-composing with the

diagonal, and doing this for k -copies of $U(1)$, we see that the multiplication-by- k map $U(1) \xrightarrow{m} U(1)$ induces the map $E^*[[x]] \rightarrow E^*[[x]]$ given by $x \mapsto [k](x)$.

Let $\varphi : \mathbb{Z}/m \rightarrow \mathbb{Z}/l$ be the homomorphism given by $1 \mapsto k$. The composition $\psi_l \circ \varphi$ is given by $1 \mapsto e^{2\pi i k/l}$. We then get the commutative diagram below by taking the upper arrow to be the multiplication-by- k map.

$$\begin{array}{ccc} U(1) & \xrightarrow{k} & U(1) \\ \psi_m \uparrow & & \uparrow \psi_l \\ \mathbb{Z}/m & \xrightarrow{\varphi} & \mathbb{Z}/l \end{array}$$

By taking $E^*(B-)$ we get the commutative diagram:

$$\begin{array}{ccc} E^*[[x]] & \xleftarrow{x \mapsto [k](x)} & E^*[[x]] \\ \downarrow x \mapsto x & & \downarrow x \mapsto x \\ E^*[[x]] / ([m](x)) & \xleftarrow{\quad} & E^*[[x]] / ([l](x)) \end{array}$$

We then see that the bottom map is also given by $x \mapsto [k](x)$. **TODO does it really follow?**

3.3 The Rings $L_r(E^*)$ and $L(E^*)$

As we have seen, to construct the character map we needed a ring R together with homomorphisms $E^*(B\Lambda_r) \rightarrow R$. It is not too hard to define the rings $L_r = L_r(E^*)$ and $L = L(E^*)$ which will be used, but it is harder to show their properties. We will give a quick definition, and claim, first without proving, the properties we need.

Recall that $E^*(BU(1)) = E^*[[x]]$, where x is the complex orientation. For any homomorphism $\alpha : \Lambda_r \rightarrow U(1)$, we can take $E^*(B-)$ to get $B\alpha^* : E^*(U(1)) \rightarrow E^*(B\Lambda_r)$. Let $S_r = \{B\alpha^*(x) \mid \alpha : \Lambda_r \rightarrow U(1), \alpha \neq 1\} \subseteq E^*(B\Lambda_r)$. We define $L_r = S_r^{-1}E^*(B\Lambda_r)$. Also, $E^*(B\Lambda_r)$ clearly has an $\text{Aut}(\Lambda_r)$ action. **TODO pre composition? post composition?** This action lifts to an action on L_r .

The projections $\Lambda_{r+1} \rightarrow \Lambda_r$ induce maps $E^*(B\Lambda_r) \rightarrow E^*(B\Lambda_{r+1})$. By post composing with the localization map, we get $E^*(B\Lambda_r) \rightarrow L_{r+1}$. It turns out that this map lifts to a map from the localization of the source $L_r \rightarrow L_{r+1}$.

Moreover, the projection $\mathrm{Aut}(\Lambda_{r+1}) \rightarrow \mathrm{Aut}(\Lambda_r)$ gives a $\mathrm{Aut}(\Lambda_{r+1})$ action on L_r , and the map $L_r \rightarrow L_{r+1}$ is equivariant with respect to that action. The colimit of this tower is the desired ring, that is $L = L(E^*) = \mathrm{colim} L_r$. Since it has a compatible $\mathrm{Aut}(\Lambda_r)$ action for all r , we get an $\mathrm{Aut}(\Lambda)$ action compatible with all of the $\mathrm{Aut}(\Lambda_r)$ actions.

References

- [Ada74] J. F. Adams. *Stable homotopy and generalised homology*. Chicago Press, Illinois and London, 1974.
- [AS69] M. F. Atiyah and G. B. Segal. Equivariant K-theory and completion. *J. Differential Geometry*, 3:1-18, 1969.
- [Hat17] A. Hatcher. *Vector Bundles and K-Theory*. November 2017. Version 2.2.
- [HKR00] Michael J. Hopkins, Nicholas J. Kuhn, and Douglas C. Ravenel. Generalized group characters and complex oriented cohomology theories. *J. Am. Math. Soc.* 13, no. 3, 553-594, 2000.
- [HS98] M. Hopkins and J. H. Smith. Nilpotence and stable homotopy theory II. *Annals of Mathematics*, 148(1), second series, 1-49, 1998.
- [LT66] J. Lubin and J. Tate. Formal moduli for one-parameter formal lie groups. *Bull. Soc. Math. France* 94, 49-59. MR 0238854, 1966.
- [Lur10] J. Lurie. Chromatic homotopy theory. *252x course notes*, <http://www.math.harvard.edu/~lurie/252x.html>, 2010.
- [Rav86] D. C. Ravenel. *Complex Cobordism and Stable Homotopy Groups of Spheres*. Academic Press, New York, 1986.
- [Rav92] D. C. Ravenel. *Nilpotence and periodicity in stable homotopy theory*. Annals of Mathematics Studies, vol. 128, Princeton University Press, Princeton, NJ, 1992.
- [Rez98] C. Rezk. Notes on the Hopkins-Miller theorem. *Homotopy theory via algebraic geometry and group representations (Evanston, IL, 1997)*, *Contemp. Math.*, vol. 220, Amer. Math. Soc., Providence, RI, 1998, pp. 313-366. MR1642902, 1998.

- [Seg68] G. B. Segal. Equivariant K-theory. *Inst. Hautes études Sci. Publ. Math. No. 34*, 129-151, 1968.
- [Ser77] J.-P. Serre. *Linear representations of finite groups*. Graduate Texts in Mathematics, vol. 42, Springer-Verlag, New York, Heidelberg, Berlin, 1977.