



# Bayesian estimation of global glacial isostatic adjustment for sea level rise re-evaluation

Zhe Sha<sup>1</sup>, Maike Schumacher<sup>1</sup>,  
Jonathan Rougier<sup>2</sup> and Jonathan Bamber<sup>1</sup>

<sup>1</sup>School of Geographical Sciences, <sup>2</sup>School of Mathematics  
University of Bristol

26 July 2017

### *GlobalMass*

- combine satellite and in-situ data related to different aspects of the sea level budget,
- attribute global sea level rise to its component parts.



European Research Council  
Established by the European Commission

Funded by the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme under grant agreement No 69418.

## The sea level budget *enigma*

$$\Delta\text{sea level}(t) = \Delta\text{barystatic}(t) + \Delta\text{steric}(t) + \text{GIA}$$

mass                      density                      ocean basins


- GIA: glacial isostatic adjustment
- inconsistencies between the discipline-specific estimates

## GlobalMass Aims

- simultaneous global estimates of all the components
- close the sea level budget

# The Bayesian hierarchical model

## Framework concept (Zammit-Mangion et al., 2015)



# First step – modelling GIA

## What is GIA?




Figure courtesy of the Canadian Geodetic Survey, Natural Resources Canada.


### GIA - glacial isostatic adjustment

- vertical displacement of the Earth's crust once burdened by ice-age glaciers ( $\sim 20,000$  yrs ago)
- constant over the time scale of this study (1981-2020)
- Regional discrepancies in estimates from physical models, e.g. ICE-6G (Peltier et al., 2015)

# The GIA process

true GIA process  
 $\mathbf{Y} : \mathbb{S}^2 \mapsto \mathbb{R}$

prior mean trend  
 $\mu : \mathbb{S}^2 \mapsto \mathbb{R}$



After de-trend, the residuals become stationary and assume

$$\mathbf{X} := \mathbf{Y} - \mu \sim \mathcal{GP}(\mathbf{0}, \kappa(\theta)) \quad (1)$$

$\kappa(\theta)$  – Matérn covariance function with parameter  $\theta = (\rho, \sigma^2)$  for the length-scale and variance.

# The GPS observations


## Vertical movement in the Earth's surface

GPS times series signals at Station  $i$ ,

- an uplift rate  $Z_i$  (mm/yr)
- a measurement error  $\varepsilon_i \sim \mathcal{N}(0, e_i^2)$
- $e_i = \text{s.d.}(Z_i)$  estimated from the time series.




A GPS station in Antarctic.



# Forward modelling

From process to observations



\*GRACE: the Gravity Recovery And Climate Experiment satellites

$\mathcal{A}_i$  maps the latent process over the appropriate spatial **foot print** to the  $i^{th}$  observation,

$$Z_i = \mathcal{A}_i Y + \varepsilon_i, \quad (2)$$

$\mathcal{A}_i$  is usually a linear operator

- point → point
- area → point
- area → area

# The BHM for GIA

Denote by  $\mathcal{A}^T = [\mathcal{A}_1^T, \dots, \mathcal{A}_N^T]$ , then we can have the vector form for equation (2)

$$\mathbf{Z} = \mathcal{A}\mathbf{Y} + \boldsymbol{\varepsilon} \quad (3)$$

## Bayesian hierarchical model for GIA

Combine the observation equation (3) and the process equation (1)

$$\begin{cases} \tilde{\mathbf{Z}} = \mathcal{A}\mathbf{X} + \boldsymbol{\varepsilon}, \boldsymbol{\varepsilon} \sim \mathcal{N}(\mathbf{0}, \text{diag}(e_1^2, e_2^2, \dots, e_N^2)) \\ \mathbf{X} \sim \mathcal{GP}(\mathbf{0}, \kappa(\theta)) \\ \theta \sim \pi(\theta) \end{cases} \quad (4)$$

# Predicting GIA

## The predictive distribution of the latent process

### Predictive Distribution of GIA

The prediction is based on the posterior marginal distribution of the latent process

$$\pi(\mathbf{X}|\tilde{\mathbf{Z}}) = \int_{\Theta} \pi(\mathbf{X}, \boldsymbol{\theta}|\tilde{\mathbf{Z}}) d\boldsymbol{\theta} \quad (5)$$

– Integrating the uncertainty of the parameters.

Other choices include the “plug-in” predictive distribution

$\pi_p(\mathbf{X}|\tilde{\mathbf{Z}}) = \pi_p(\mathbf{X}|\tilde{\mathbf{Z}}, \hat{\boldsymbol{\theta}})$ , where  $\hat{\boldsymbol{\theta}}$  are the estimated values of the parameters, e.g. posterior mean or mode.

# Predicting GIA

## Point-wise prediction of the latent process

### Point-wise predicted mean and uncertainty

For point-wise update at  $X_i$  on a fine grid, we use

$$\text{predicted mean: } X_i^* = \mathbb{E}(X_i | \tilde{\mathbf{Z}})$$

$$\text{predicted uncertainty: } u_i^* = \text{s.d.}(X_i | \tilde{\mathbf{Z}})$$

$$\text{where } \pi(X_i | \tilde{\mathbf{Z}}) = \int_{\mathcal{X}_{-i}} \pi(\mathbf{X} | \tilde{\mathbf{Z}}) d\mathbf{X}_{-i}.$$

When using the “plug-in” predictive distribution,  $\pi_p(X_i | \tilde{\mathbf{Z}})$  is Gaussian and the Bayesian update can be written in closed form.

# The GMRF approximation

- Bayesian update of the GP on a grid of  $m$  points scales as  $\mathcal{O}(m^3)$ ,  $m \sim 10^5$  for a  $1^\circ$  global grid.
- Gaussian Markov random field (GMRF) with sparse precision matrix has a much better scaling property.

## The SPDE approach (Lindgren et al., 2011)

Denote by  $\mathbf{x}$  the GMRF approximation of  $\mathbf{X}$  on a triangulation of  $m$  vertices with basis functions  $\{\phi_i\}_{i \in \mathbb{N}}$ , then for  $s \in \mathbb{S}^2$

$$\mathbf{X}(s) \approx \phi_i(s)^T \mathbf{x} \quad (6)$$

For a grid  $\mathcal{S} \subset \mathbb{S}$  of locations, we find the basis matrix  $\mathbf{C}$  and write  $\mathbf{X}(\mathcal{S}) \approx \mathbf{Cx}$ .

# The GMRF approximation


Now the BHM in equation (4) becomes

$$\begin{cases} \tilde{\mathbf{Z}} = \mathcal{A}\mathbf{C}\mathbf{x} + \varepsilon, \quad \varepsilon \sim \mathcal{N}(\mathbf{0}, \text{diag}(e_1^2, e_2^2, \dots, e_N^2)) \\ \mathbf{x} \sim \mathcal{N}(\mathbf{0}, \mathbf{Q}^{-1}(\theta)) \\ \theta \sim \pi(\theta) \end{cases} \quad (7)$$






$\mathbf{x}$  is GMRF approximation defined by the precision matrix  $\mathbf{Q}$ .

# Choosing the mesh on a sphere

semi-regular mesh




irregular mesh



Parameter estimation using different meshes.


# Mesh effect

The predicted uncertainties using regular mesh




# Mesh effect

The predicted uncertainties using irregular mesh



# First solution for GIA



# Conclusions and outlook

- Testing the statistical framework for the GIA.
- Requiring improved mesh.
  - Adaptive – fast computation
  - Regular – reliable approximation
- Improving GPS data quality.
- Extending to the full system.

# References

- Lindgren, F., Rue, H., and Lindström, J. (2011). An explicit link between Gaussian fields and Gaussian Markov random fields: the stochastic partial differential equation approach. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 73(4):423–498.
- Peltier, W. R., Argus, D. F., and Drummond, R. (2015). Space geodesy constrains ice age terminal deglaciation: The global ICE-6G\_C (VM5a) model. *Journal of Geophysical Research: Solid Earth*, 120(1):450–487.
- Zammit-Mangion, A., Rougier, J., Schön, N., Lindgren, F., and Bamber, J. (2015). Multivariate spatio-temporal modelling for assessing Antarctica's present-day contribution to sea-level rise. *Environmetrics*, 26(3):159–177.

# Questions

---

# Thank you!