

Status of Black Rockfish (*Sebastes melanops*) off Oregon and federal waters in 2023



by
Jason M. Cope¹
Alison D. Whitman²
Aaron M. Berger¹
Leif R. Rasmussen²

¹Northwest Fisheries Science Center, U.S. Department of Commerce, National Oceanic and Atmospheric Administration, National Marine Fisheries Service, 2725 Montlake Boulevard East, Seattle, Washington 98112

²Oregon Department of Fish and Wildlife, 2040 Southeast Marine Science Drive, Newport, Oregon 97365

© Pacific Fishery Management Council, 2023

Correct citation for this publication:

Cope, J.M., A.D. Whitman, A.M. Berger, L.R. Rasmuson. 2023. Status of Black Rockfish (*Sebastodes melanops*) off Oregon and federal waters in 2023. Pacific Fishery Management Council, Portland, Oregon. 209 p.

Contents

Disclaimer	1
Executive summary	i
Stock	i
Removals	i
Data and assessment	iii
Stock biomass and dynamics	iii
Recruitment	vi
Exploitation status	vi
Ecosystem considerations	x
Reference points	x
Management performance	xiii
Unresolved problems and major uncertainties	xiii
Scientific uncertainty	xiii
Harvest Projections and Decision Table	xiv
Research and data needs	xvi
1 Introduction	1
1.1 Basic Information	1
1.2 Life History	2
1.3 Ecosystem Considerations	3
1.4 Historical and Current Fishery Information	4
1.5 Summary of Management History and Performance	4
1.6 Canadian and Alaska fisheries	6
2 Data and Model Inputs	6
2.1 Removals	6
2.1.1 Commercial Landings and Discards	6
2.1.2 Recreational Landings and Discards	8
2.2 Length and Age Composition Data	11
2.2.1 Length Compositions	11

2.2.2	Age Compositions	12
2.3	Mean weights	13
2.4	Fishery-Dependent Abundance Indices	13
2.4.1	Commercial Index of Abundance - Nearshore Fishery Logbook CPUE	13
2.4.2	Oregon Recreational Indices of Abundance	14
2.5	Fishery-Independent Data	17
2.5.1	Abundance Indices	17
2.6	Biological Data and Parameters	20
2.6.1	Natural Mortality	21
2.6.2	Growth (Length-at-Age)	22
2.6.3	Ageing Bias and Precision	23
2.6.4	Length-Weight Relationship	24
2.6.5	Maturation and Fecundity	24
2.6.6	Stock-recruitment function and compensation	25
2.6.7	Sex Ratio	26
2.7	Environmental and Ecosystem Data	26
2.8	Data Considered but Not Included	26
3	Assessment Model	27
3.1	Summary of Previous Assessments and Reviews	27
3.1.1	History of Modeling Approaches Used for this Stock	27
3.1.2	Most Recent STAR Panel and SSC Recommendations	28
3.2	Model description	29
3.2.1	Modelling Platform	29
3.2.2	Bridging the assessment model from Stock Synthesis 3.24 to 3.30	29
3.3	Model Structure and Assumptions	30
3.4	Model Structure, Evaluation, and Specification	30
3.4.1	Fleet and survey designations	30
3.5	Model likelihood components	31
3.6	Reference model exploration, key assumptions and specification	31
3.6.1	Data Weighting	33

3.6.2	Model Changes from the Last Assessment	34
3.7	Reference Model Diagnostics and Results	34
3.7.1	Model Convergence and Acceptability	34
3.8	Base Model Results	34
3.8.1	Reference Model Outputs	36
3.9	Characterizing uncertainty	38
3.9.1	Sensitivity Analyses	38
3.9.2	Likelihood Profiles	41
3.9.3	Retrospective Analysis	42
3.9.4	Unresolved Problems and Major Uncertainties	42
4	Management	43
4.1	Reference Points	43
4.2	Management performance	44
4.3	Unresolved Problems and Major Uncertainties	44
4.4	Harvest Projections and Decision Tables	44
4.5	Evaluation of Scientific Uncertainty	45
4.6	Research and Data Needs	45
5	Acknowledgments	47
6	References	48
7	Tables	55
8	Figures	105
9	Appendix A: Detailed Fit to Length Composition Data	165
10	Appendix B: Fit to Conditional-Age-at-Length Composition Data	172
11	Appendix C: Fit to Conditional-Age-at-Length Composition Data	184
12	Appendix D: Passive Fit to Marginal Age Composition Data	202

13 Appendix E: Numbers at Age Plot	206
13.1 Females	206
13.2 Males	206
14 Appendix F: Numbers at Length Plot	208
14.1 Females	208
14.2 Males	209

Disclaimer

These materials do not constitute a formal publication and are for information only. They are in a pre-review, pre-decisional state and should not be formally cited or reproduced. They are to be considered provisional and do not represent any determination or policy of NOAA or the Department of Commerce.

Executive summary

Stock

This assessment reports the status of Black Rockfish (*Sebastes melanops*) off Oregon state using data through 2022. Black Rockfish are also found in California and Washington waters of the U.S. West Coast, and those are treated in separate area-based stock assessments given different management considerations and exploitation histories as discussed at the preliminary stock assessment workshop in February 2023 (Council 2023). There is substantial biogeographic separation in the populations off Oregon and Washington, thus justifying separation of those populations into different management units and stock assessments. Black Rockfish are also caught from the waters off British Columbia and Alaska, but there have not been any formal assessments of stock status for those areas in recent years. Genetic studies of stock structure indicate fish in Alaska are more differentiated than those along the contiguous west coast of the United States, and that genetic diversity varies in a non-systematic way from California to Oregon (Hess et al. 2023).

Removals

Black Rockfish are caught by a wide variety of gear types in Oregon and in recent decades have been a very important target species for recreational charter-boats and private sport anglers (i.e., ocean boat fleet). In recent years the recreational fishery has accounted for most of the Black Rockfish catches (Figure i). Black Rockfish are also be an important component of nearshore commercial fisheries, either as incidental catch by the troll fishery for salmon or as directed catch by fixed gear fisheries for groundfish. Further, there are nearshore commercial fisheries that catch and sell fish live for the restaurant trade.

Historical catch reconstructions for Oregon show that catch of Black Rockfish was limited until the development of trawl fisheries around World War II in the 1940s. Trawl landings were consistent through the 1970s and landings from other gear types were minimal (Figure i). There has been almost no trawl-caught landings of Black Rockfish in recent years (Table i). Modern-era, detailed catch estimates are available from Pacific Fishery Information Network (PacFIN) database for commercial landings (1987 - 2022) and from the Recreational Fishery Information Network (RecFIN) (2001 - 2022). Historical recreational catch estimates were provided by ODFW (1979 - 2000). Generally, the historical catch series were derived by applying available estimates or assumed values for the proportion of Black Rockfish landings in reported landings of rockfish. Observer data, which are available since the early 2000s, indicate low levels of discarding of Black Rockfish, generally less than 2% of total catch. While Black Rockfish are unlikely to have ever comprised a large percentage of overall rockfish landings due to their low abundance compared to other rockfish species, it seems plausible that they have been more than a trivial component due to their nearshore distribution for many years. Black Rockfish were one of only four rockfish species mentioned by scientific name in reports of rockfish landings in Oregon during the 1940s.

Overall, removals of Black Rockfish remained relatively low (less than 100 mt) until the mid to late 1970s when landings quickly quadrupled with the expansion of the recreational

fishery. Since the 1980s, removals have consistently fluctuated between 300 and 600 mt (no major trend), comprising mostly of removals from the ocean boat recreational fleet and the non-trawl commercial fleet (Figure i).

Table i: Recent landings by fleet and total landings summed across fleets.

Year	Trawl wdis	Non-Trawl wdis	Ocean	Shore	Total Landings
2013	0.08	107.47	321.31	13.28	442.14
2014	0.71	122.92	342.55	13.28	479.45
2015	0.51	122.12	458.61	13.28	594.52
2016	0.64	105.95	404.65	13.28	524.53
2017	0.26	125.21	402.67	13.28	541.43
2018	0.03	122.98	278.78	13.28	415.07
2019	0.01	119.19	305.07	13.28	437.55
2020	0.04	101.79	320.45	13.28	435.56
2021	0.00	101.53	320.73	13.28	435.54
2022	0.00	118.11	394.61	13.28	526.00




Figure i: Landings by fleet used in the reference model where catches in metric tons by fleet are stacked.

Data and assessment

The first Black Rockfish stock assessment along the west coast of the United States that included the majority of Oregon waters was completed in 1993, covering the area south of Cape Falcon, Oregon to north of Point Piedros Blancos, California (Sampson 2007). Subsequent assessments that included Oregon waters were done in 2003 and 2007. In 2015, the last assessment prior to the current one was completed that included Oregon waters only as one of three (also Washington and California) separate assessment areas delineated by state lines (Cope et al. 2016). Similarly, this assessment treats Oregon waters as a single assessment area. The previous two assessments used Stock Synthesis software, as does this one (version 3.30.21.00).

This assessment integrates data and information from multiple sources into one modeling framework. The stock assessment model for Black Rockfish is informed by catch data from two commercial fleets and two recreational fleets, six abundance indices, five sets of length composition data, and three sets of conditional age-at-length compositions. It also uses multiple ageing error matrices to incorporate ageing imprecision and applies fixed parameterizations of weight-at-length, maturity-at-length, fecundity-at-length, the Beverton-Holt stock-recruitment steepness value, and recruitment variability. Life history parameters were sex-specific (i.e., a two-sex model) with natural mortality fixed at external estimates and growth and recruitment parameters estimated. Additional parameters that were estimated include initial population scale ($\ln R_0$), selectivity for each fishery and survey, and extra survey variance. The base model was tuned to account for the weighting of the length and age data and index variances (which was estimated), as well as the specification of the recruitment bias adjustments. Derived quantities include, among other things, the time series of spawning biomass, age and size structure, and current and projected future stock status. The model covers the years 1892 to 2022, with a 12 year forecast beginning in 2023.

Within model uncertainty is explicitly included in this assessment by parameter estimation uncertainty, while among model uncertainty is explored through sensitivity analyses addressing alternative input assumptions such as data treatment and weighting, and model specification sensitivity to the treatment of life history parameters, selectivity, recruitment, and survey catchability. A reference model was selected that best fit the observed data while concomitantly balancing the desire to capture the central tendency across those sources of uncertainty, ensure model realism and tractability, and promote robustness to potential model misspecification.

Stock biomass and dynamics

Spawning output (in millions of eggs; meggs) instead of spawning biomass is used to report the functionally mature population scale because fecundity is nonlinearly related to body female weight. The estimated spawning output at the beginning of 2023 was 900 meggs (~95 percent asymptotic intervals: 855 to 944 meggs, Table ii and Figure ii), which when compared to unfished spawning output (1,633) meggs gives a relative stock status level of 55 percent (~95 percent asymptotic intervals: 53 to 57 percent, Figure iii). Overall, spawning output declined with the onset of increasing commercial removals in the 1960s and continued

to decline with the increase in recreational catches through the 1990s, even dropping below the target relative stock size from 1993 to 2008, before steadily increasing back above target since that time. The largest of the estimated recruitment pulses occurred in 2008 and was followed by several above average recruitment years in the early 2010s, which contributed to the increase in spawning output through the mid to late 2010s. The minimum relative stock size of 25 percent of unfished levels is estimated to have occurred in 1999. Accordingly, the stock has not been below the minimum stock size threshold (i.e., never overfished based on median estimates). Currently the stock is estimated above the management target of $SO_{40\%}$ in 2023 and is estimated to have remained above the target since 2009 (Table ii and Figure iii).

Table ii: Estimated recent trend in spawning output and the fraction unfished and the 95 percent intervals.

Year	Spawning Output	Lower Interval	Upper Interval	Fraction Unfished	Lower Interval	Upper Interval
2013	719.26	634.23	804.29	0.44	0.40	0.48
2014	726.10	641.98	810.22	0.44	0.41	0.48
2015	740.31	656.93	823.70	0.45	0.41	0.49
2016	744.99	662.79	827.20	0.46	0.42	0.49
2017	759.97	679.34	840.59	0.47	0.43	0.50
2018	773.49	695.31	851.66	0.47	0.44	0.51
2019	805.69	730.63	880.75	0.49	0.46	0.53
2020	838.15	767.58	908.72	0.51	0.48	0.55
2021	868.88	804.45	933.31	0.53	0.50	0.56
2022	894.62	838.74	950.51	0.55	0.52	0.57
2023	899.77	855.05	944.49	0.55	0.53	0.57




Figure ii: Estimated time series of spawning output (circles and line: median; light broken lines: 95 percent intervals) for the base model.




Figure iii: Estimated time series of fraction of unfished spawning output (circles and line: median; light broken lines: 95 percent intervals) for the base model.

Recruitment

Recruitment is informed by the data and estimated from 1980 to 2017, before and after which it is assumed taken from the stock-recruit relationship (Table iii and Figure iv). The highest recruitment years occurred in 1999, 2000, 2008, 2013, and 2016. The large 2008 and 2016 year classes, as well as several above average year classes in the early 2010s, contributed to the recent increase in Black Rockfish biomass. Recruitment is informed by composition data and six relative abundance indices. The 2015 stock assessment did not estimate deviations from the stock-recruitment curve. While the Black Rockfish stock has been reduced to levels that theoretically would provide some information on how recruitment compensation changes across spawning biomass levels (i.e., inform the steepness parameter), the assessment model could not adequately estimate a reasonable steepness parameter. Thus, recruitment is based on a fixed assumption about steepness ($h = 0.72$) and recruitment variability ($\sigma_R = 0.6$).

Table iii: Estimated recent trend in recruitment and recruitment deviations and the 95 percent intervals.

Year	Recruit- ment	Lower Interval	Upper Interval	Recruit- ment Deviations	Lower Interval	Upper Interval
2013	5030.30	4417.04	5728.71	0.56	0.42	0.71
2014	4638.54	4014.69	5359.33	0.48	0.32	0.64
2015	2255.93	1741.04	2923.10	-0.24	-0.51	0.02
2016	6224.71	5294.64	7318.15	0.77	0.59	0.95
2017	1595.64	991.03	2569.12	-0.64	-1.12	-0.15
2018	3436.75	3317.86	3559.90	0.00	0.00	0.00
2019	3462.37	3346.58	3582.16	0.00	0.00	0.00
2020	3486.56	3374.00	3602.87	0.00	0.00	0.00
2021	3508.07	3398.78	3620.87	0.00	0.00	0.00
2022	3525.14	3419.31	3634.25	0.00	0.00	0.00
2023	3528.45	3426.30	3633.64	0.00	0.00	0.00

Exploitation status

Fishing intensity, as measured by the SPR rate (1 - SPR), has fluctuated around the estimated SPR rate fishing intensity target of 0.50 (1 - SPR_{50%}) since 1980. Fishing was at or above the target rate from 1989 to 2005 and has been slightly below it over the past 5 years (Table iv and Figures vi and vii). The steepness value of 0.72 indicates that a lower value of SPR (or equivalently a higher fishing intensity than SPR_{50%}) would be consistent with the biomass-based target of (SO_{40%}) for sustainable removals. Trends in fishing intensity largely mirrored that of landings until the 1990s, after which recruitment pulses countered the catches somewhat to lower overall fishing intensity (Figure vi). The maximum fishing




Figure iv: Estimated time series of age-0 recruits (1000s) for the base model with 95 percent intervals.




Figure v: Estimated time series of recruitment deviations.

intensity was 0.7 in 1992, which is well above the target SPR-based harvest rate of 0.50. The current level of 0.42 for 2022 is below that target. Fishing intensity over the past decade has ranged between 0.4 and 0.56 and the exploitation rate (range of 0.04 - 0.08, Table iv) has come down since the time series high of 0.12 in 1992. Current estimates indicate that Black Rockfish spawning output is greater than the target biomass level ($SO_{40\%}$), though fishing intensity remains near the target F_{MSY} proxy harvest rate of $1 - SPR_{50\%}$ (Figure vii).

Table iv: Estimated recent trend in the 1-SPR where SPR is the spawning potential ratio the exploitation rate, and the 95 percent intervals.

Year	1-SPR	Lower Interval	Upper Interval	Exploitation Rate	Lower Interval	Upper Interval
2013	0.49	0.46	0.51	0.06	0.06	0.06
2014	0.50	0.48	0.52	0.06	0.06	0.07
2015	0.56	0.54	0.58	0.08	0.07	0.08
2016	0.52	0.49	0.54	0.07	0.06	0.07
2017	0.52	0.50	0.54	0.07	0.07	0.07
2018	0.43	0.41	0.45	0.05	0.05	0.05
2019	0.44	0.42	0.45	0.05	0.05	0.05
2020	0.43	0.42	0.44	0.05	0.05	0.05
2021	0.42	0.42	0.43	0.05	0.05	0.05
2022	0.48	0.48	0.48	0.06	0.06	0.07




Figure vi: Estimated 1 - relative spawning ratio (SPR) by year for the base model. The management target is plotted as a red horizontal line and values above this reflect harvest in excess of the proxy harvest rate.

Ecosystem considerations

This stock assessment does not explicitly incorporate trophic interactions, habitat factors or environmental factors into the assessment model. More predation, diet and habitat work, and mechanistic linkages to environmental conditions would be needed to incorporate these elements into the stock assessment and should remain a priority. McClure et al. (2023) report the climate vulnerability for several west coast groundfishes, including Black Rockfish. Black Rockfish demonstrated both high biological sensitivity and high climate exposure risk, to give it an overall high vulnerability score to climate change. This result should also be considered with the fact that, like many rockfishes, periods of low productivity is not unusual to Black Rockfish and their extended longevity (though admittedly this seems shorter than previously believed and should be reconsidered) has historically allowed them to wait for advantageous productivity periods. Additional stressors such as fishing and climate change that possibly truncate longevity could bring significant challenges to population sustainability.

Reference points

Reference points were based on the rockfish FMSY proxy ($SPR_{50\%}$), target relative biomass (40%), and estimated selectivity and catch for each fleet (Table v). The Black Rockfish population in Oregon at the start of 2023 is estimated to be 1.38 times (above) the target biomass, and fishing intensity during 2022 is estimated to be 0.96 times (below) the fishing intensity target (Figure vii). The yield values are lower than the previous assessment for similar reference points due to updated life history estimates. The proxy MSY values of management quantities are the most conservative compared to the estimated MSY and MSY relative to 40% biomass. Sustainable total yield, removals, using the proxy $SPR_{50\%}$ is 455 mt. The spawning output equivalent to 40 percent of the unfished spawning output ($SO_{40\%}$) calculated using the SPR target ($SPR_{50\%}$) was 728.5 meggs.

Recent removals have been close to the point estimate of potential long-term yields calculated using an $SPR_{50\%}$ reference point, though the population size has continued to increase over recent years due to several above average recruitments. The equilibrium estimates of yield relative to biomass based on a steepness value fixed at 0.72 are provided in Figure viii, where vertical dashed lines indicate the estimate of fraction unfished at the start of 2023 (current) and the estimated management targets calculated based on the relative target biomass (B target), the SPR target, and the maximum sustainable yield (MSY).

The 2023 spawning biomass relative to unfished equilibrium spawning biomass, based on the 2022 fishing year, is above (55 percent) the management target of 40 percent of unfished spawning output. The relative biomass and the ratio of the estimated SPR to the management target ($SPR_{50\%}$) across all model years are shown in Figure vii where warmer colors (red) represent early years and colder colors (blue) represent recent years. There have been periods where the stock status has decreased below the target and fishing intensity has been higher than the target fishing intensity based on $SPR_{50\%}$.




Figure vii: Phase plot of estimated 1-SPR versus fraction unfished for the base model.




Figure viii: Equilibrium yield curve for the base case model. Values are based on (the time invariant) fishery selectivities and with steepness fixed at 0.72.

Table v: Summary of reference points and management quantities, including median estimates and associated 95 percent intervals.

	Estimate	Lower Interval	Upper Interval
Unfished Spawning Output	1632.81	1577.19	1688.43
Unfished Age 0+ Biomass (mt)	11576.10	11142.20	12010.00
Unfished Recruitment (R0)	3807.86	3678.16	3937.56
Spawning Output (2023)	899.77	855.05	944.49
Fraction Unfished (2023)	0.55	0.53	0.57
Reference Points Based SB _{40%}			
Proxy Spawning Output SB _{40%}	653.12	630.88	675.37
SPR Resulting in SB _{40%}	0.46	0.46	0.46
Exploitation Rate Resulting in SB _{40%}	0.07	0.07	0.07
Reference Points Based on SPR Proxy for MSY			
Proxy Spawning Output (SPR50)	728.48	703.67	753.30
SPR50	0.50		
Exploitation Rate Corresponding to SPR50	0.06	0.06	0.07
Yield with SPR50 at SB SPR (mt)	454.96	438.30	471.63
Reference Points Based on Estimated MSY Values			
Spawning Output at MSY (SB MSY)	386.95	373.08	400.83
SPR MSY	0.31	0.31	0.31
Exploitation Rate Corresponding to SPR MSY	0.11	0.11	0.11
MSY (mt)	533.65	514.71	552.60

Management performance

Black Rockfish removals have generally been below the equivalent ABC-ACL over the recent decade, with the exception of 2017 and 2022 when removals were slightly higher (Table vi). Exploitation on Black Rockfish increased starting around 1940 and reached a high in the late 1970s. Since that time, catch has mostly fluctuated between 300 and 500 mt per year, with some years exceeding 600 mt. Removals have averaged 483 mt over the past decade. The last ten years of Black Rockfish acceptable biological catch (ABC) and annual catch limit (ACL) (which are equivalent) has been set, by definition, below the overfishing limit (OFL) (Table vi). Prior to 2017, management specifications were set for Black Rockfish in California and Oregon waters combined. The Black Rockfish OFL has not been exceeded by Oregon removals over the past decade.

Table vi: Recent trend in the overfishing limits (OFL), the acceptable biological catches (ABCs), the annual catch limits (ACLs), and the total catch (mt). Management specifications prior to 2017 were for Oregon and California combined.

Year	OFL	ABC	ACL	Catch
2013	1159	1108	1000	442.14
2014	1166	1115	1000	479.45
2015	1176	1124	1000	594.52
2016	1183	1131	1000	524.53
2017	577	527	527	541.43
2018	570	520	520	415.07
2019	565	516	516	437.55
2020	561	512	512	435.56
2021	570	512	512	435.54
2022	566	512	512	526.00

Unresolved problems and major uncertainties

The biggest uncertainty and unresolved conflict is trying to reconcile the signal in the biological data (which suggests a lower population size and status) versus the acoustic and tag surveys (which suggest high stock sizes and status). This is the major issue the current assessment is confronting. Another acoustic-visual survey data point could help resolve how much uncertainty there is in the estimate. The lack of contrast in the biological data, despite large sample sizes, is another barrier to interpreting the current conditions, though given models using only biological data, the signal seems clear that the population could be at a lower stock status.

Scientific uncertainty

The model-estimated uncertainty around the 2023 spawning biomass was $\sigma = 0.03$ and the uncertainty around the OFL was $\sigma = 0.01$. This is clearly an underestimate of overall

uncertainty because of the necessity to fix some life history parameters such as natural mortality and steepness, as well as a lack of explicit incorporation of model structural uncertainty. The alternative states of nature used to bracket uncertainty in the decision table assist with encapsulating model structure uncertainty.

Harvest Projections and Decision Table

The following text will be modified, as appropriate, after the STAR panel and SSC meeting.

The Black Rockfish assessment is being considered as a category 2 assessment with a P^* = 0.45, sigma = 0.72, and a time-varying buffer applied to the OFL. These multipliers are also combined with the rockfish MSY proxy of FSPR=50% MSY and the 40-10 harvest control rule to calculate OFLs, ABCs and ACLs. A twelve year (2023-2034) projection of the reference model using these specifications along with input removals for 2023 and 2024 provided by the Groundfish Management Team is provided in Table vii.

Table vii: Projections of potential OFLs (mt), ABCs (mt), estimated spawning output, and fraction unfished.

Year	Predicted OFL (mt)	ABC Catch (mt)	Age 0+ Biomass (mt)	Spawning Output	Fraction Unfished
2023	541.9	496.5	7981.8	899.8	0.55

Uncertainty in management quantities for the reference model was characterized by exploring various model specifications in a decision table. Initial explorations considered alternative specifications of catchability for the acoustic-visual survey. The resultant decision table will be provided in Table viii.

Further details about selecting the decision table states of nature will be added here after the STAR panel.

Table viii: Decision table summary of 10 year projections beginning in 2023 for alternative states of nature based on an axis of uncertainty related to model structure relative to the reference model (i.e., estimate catchability, Q, associated with the acoustic-visual survey and no estimation of recruitment deviations) . Columns range over low (12.5 quantile), mid (reference model), and high states (87.5 quantile) of nature and rows range over different catch level assumptions. Values in italics indicate years where the stock size prevented the full catch removals.

Year	Catch	Estimate Q		Reference Model		No Rec Devs	
		Spawning Output	Fraction Unfished	Spawning Output	Fraction Unfished	Spawning Output	Fraction Unfished
2023	-	-	-	-	-	-	-
2024	-	-	-	-	-	-	-
2025	-	-	-	-	-	-	-
2026	-	-	-	-	-	-	-
2027	-	-	-	-	-	-	-
P*=0.45	2028	-	-	-	-	-	-
<i>sigma=0.5</i>	2029	-	-	-	-	-	-
	2030	-	-	-	-	-	-
	2031	-	-	-	-	-	-
	2032	-	-	-	-	-	-
	2033	-	-	-	-	-	-
	2034	-	-	-	-	-	-
	2023	-	-	-	-	-	-
	2024	-	-	-	-	-	-
	2025	-	-	-	-	-	-
	2026	-	-	-	-	-	-
P*=0.4	2027	-	-	-	-	-	-
<i>sigma=0.5</i>	2028	-	-	-	-	-	-
	2029	-	-	-	-	-	-
	2030	-	-	-	-	-	-
	2031	-	-	-	-	-	-
	2032	-	-	-	-	-	-
	2033	-	-	-	-	-	-
	2034	-	-	-	-	-	-
	2023	-	-	-	-	-	-
	2024	-	-	-	-	-	-
	2025	-	-	-	-	-	-
	2026	-	-	-	-	-	-
Equilibrium	2027	-	-	-	-	-	-
yield from	2028	-	-	-	-	-	-
FMSY proxy	2029	-	-	-	-	-	-
of SPR=0.5	2030	-	-	-	-	-	-
	2031	-	-	-	-	-	-
	2032	-	-	-	-	-	-
	2033	-	-	-	-	-	-
	2034	-	-	-	-	-	-

Research and data needs

Recommended avenues for research to help improve future Black Rockfish stock assessments:

1. Continue work on the investigation into the movement, and behavior or mortality of older (> age 10) females to further reconcile their absence in fisheries data.
2. Conduct population genetics studies on fish observed off of the continental shelf (middle of the gyre and at sea mounts) to determine their association with the nearshore stocks.
3. Continue to build evidence for appropriate natural mortality values for females and males.
4. Improved historical catch reconstructions. Specifically, the historic trawl fishery catches (pre-1987) in particular require particular attention. Given the huge historical removals of that fleet in each state, the assessment is very sensitive to the assumed functional form of selectivity. A synoptic catch reconstruction is recommended, where states work together to resolve cross-boundary state catch issues as well as standardize the approach to catch recommendations to the extent possible.
5. Stock structure for black rockfish is a complicated topic that needs further analysis. How this is determined (e.g., exploitation history, genetics, life history variability, biogeography, etc.) and what this means for management units needs to be further refined. This is a general issue for all nearshore stocks that likely have significant and small scale stock structure among and within states, but limited data collections to support small-scale management.
6. Continue acoustic-visual fisheries independent coastwide survey to develop a time series. Further refine the survey by addressing the recommendations of the SSC methodology review from 2022. Examine the potential of using spatial modeling to reduce the uncertainty in the population estimates from the acoustic-visual fisheries independent coastwide survey.
7. Reconcile contradictory signal in the Black Rockfish biology versus the population scale.
8. Better understand the ecology and habitats of Black Rockfish from settlement to age 1. Further development of surveys aimed specifically at recruitment or settlement rates of nearshore species, such as OSU's SMURF collections, that are not frequently encountered in offshore federal age-0 surveys is needed.

1 Introduction

This assessment report describes the stock of Black Rockfish (*Sebastodes melanops*) off the Oregon coast in U.S. waters, using data through 2022. The stocks of Black Rockfish in Washington, Oregon, and California waters are each modeled as separate stocks with the population estimates reported in stock-specific assessment reports (Figure 1). This assessment does not account for populations located in other areas off the U.S. West Coast and assumes that California and Washington populations do not contribute to nor take from the Oregon population being assessed here.

1.1 Basic Information

Black Rockfish (*Sebastodes melanops*) are an important component of the recreational fisheries in the nearshore waters off central and northern California, Oregon, and Washington, as well as the non-trawl commercial fisheries in California and Oregon. They range as far north as Amchitka and Kodiak islands in Alaska and are considered uncommon south of central California (Love et al. 2002).

An early assessment of Black Rockfish considered the population off Oregon and California (Ralston and Dick 2003). This assessment reviewed the evidence supporting genetic stock structure for Black Rockfish and other rockfish off the U.S. West Coast. This work concluded that the Oregon and California populations of Black Rockfish are probably not genetically heterogeneous. That assessment treated the Black Rockfish off California and Oregon as a unit stock. Previous assessments of Black Rockfish off Washington (Wallace et al. 1999; Wallace and Tsou 2007) describe a study of coastal Black Rockfish genetic structure using 10 sampled sites collected from northern California to southern British Columbia. Results of that study supported the notion of separate genetic stocks north and south of Cape Falcon, Oregon. However, a later study (Baker 1999) using Black Rockfish collected from eight sites along the northern Oregon coast concluded that Black Rockfish from north and south of Cape Falcon were genetically very similar.

Although a stock boundary line at the Columbia River seems reasonable for Black Rockfish, both because it is a state fishery management boundary and because the Columbia River plume is likely to be somewhat of a natural barrier (impediment) to the north-south exchange of Black Rockfish adults and larvae, the 2007 assessment of Black Rockfish off Oregon and California (Sampson 2007) differed slightly from Ralston and Dick (2003) by placing the northern boundary at Cape Falcon rather than at the Columbia River. The boundary was changed to avoid overlap with the separate northern assessment (Wallace and Tsou 2007) and to simplify the process of assembling historical commercial landings data, which are largely available in terms of Pacific Marine Fisheries Commission (PMFC) statistical areas. Given the spatial resolution of the historical commercial fishery data, it is problematic to estimate the catch of Black Rockfish taken north of Cape Falcon but south of the Columbia

River. Addressing the spatial distribution of the catch around the OR-WA state boundary was a major focus of both the 2015 Black Rockfish assessment (Cope et al. 2016) and the current assessment.

The 2015 Black Rockfish assessment delineated the stock at state borders and three models were used to estimate a coastwide stock status (Cope et al. 2016). This approach was considered consistent with two ideas: (a) that nearshore species do not exhibit much adult movement and (b) exploitation and management histories have varied significantly among the three states. This stratification was maintained in the current assessment and this document addresses the portion of the Black Rockfish stock in Oregon waters.

In Oregon, accounting for location of capture is problematic for historic commercially caught fish, primarily for trawl gear types, which was the predominant gear type during the 1940s to 1970s. For state-specific regional assessments the commercially caught Black Rockfish were apportioned to assessment region based on the port of landing, with the exception of trawl caught fish landed into Astoria, OR. Most of these fish were found to have been caught off Washington and most of the trawl landings into Astoria were therefore included with the catch history for the Washington assessment region. This approach was refined in the current assessment to allow for state-specific species compositions to be applied to aggregated trawl landings. Details are provided in Section 2.1.1 Commercial Landings and Discards.

1.2 Life History

Adult Black Rockfish tend to occur in schools over rocky structure at depths less than 40 fathoms, and sometimes feed actively on or near the surface. They feed on a wide variety of prey including zooplankton, krill, mysids, sand lance, and juvenile rockfish, and are subject to predation by Lingcod and marine mammals (Love et al. 2002).

Although tagging studies have documented some individuals moving long distances (several hundreds of miles), the vast majority of recaptured individuals were found close to the areas of initial capture and tagging (Culver 1987; Ayres 1988; Starr and Green 2007; Wallace et al. 2010). Results from a 2004-05 study off Newport, OR of 42 Black Rockfish implanted with acoustic tags indicated that all but seven fish remained within range of a 3 x 5 km array of acoustic receivers during one full year of monitoring and had relatively small home ranges that did not vary seasonally (Parker et al. 1995). Green and Starr (2011) report similar findings from a study in Carmel Bay, CA of 23 acoustically tagged Black Rockfish. An extensive Washington state tagging study also supported low movements for most individuals, with some exceptional movements recorded (Wallace et al. 2010).

Like all members of the genus *Sebastodes*, Black Rockfish have internal fertilization and bear live young approximately two months after insemination. Black Rockfish are quite fecund, with a six-year-old female annually producing about 300,000 embryos and a 16-year-old producing about 950,000 embryos (Bobko and Berkeley 2004a). Recent studies have demonstrated that

the relative number and quality of larvae increase with age in female Black Rockfish (Berkeley et al. 2004; Hixon et al. 2014a). Parturition of larvae occurs during winter (Echeverria 1987) and larvae and small juveniles are pelagic for several months to a year (Boehlert and Yoklavich 1983). Settlement occurs in estuaries, tide-pools, and in the nearshore at depths less than 20 m (Stein and Hassler 1989).

Black Rockfish begin recruiting to nearshore fisheries at 3-4 years of age, corresponding to a fork length of about 25-30 cm, and 50% of females attain maturity at about 6-8 years, corresponding to a fork length of about 38-42 cm. Adult female Black Rockfish grow 3-5 cm larger than males, with a few females attaining fork lengths greater than 55 cm.

1.3 Ecosystem Considerations

The prominent position of Black Rockfish as both a major predator (adult stage) and prey (larvae to juvenile stages) in mostly nearshore areas suggest it to be a notable player in the nearshore ecosystem. The California Current is a highly variable and dynamic system, and it has been recognized for years that rockfishes are subject to large swings in recruitment that are tied to environmental conditions. It is believed to be one of the reasons rockfishes exhibit long lives and the ability to go years without significant recruitment, but can produce large recruitment events when conditions are favorable (a phenomenon known as the “storage effect”).

Black Rockfish off central Oregon have recently been shown to exhibit changes in larval and juvenile growth rates that correlate with prey abundance and water temperature, to name just a few of the factors (Fennie et al. 2023). Settlement rates exhibited a dome-shaped relationship, demonstrating a “window” of good growth conditions for successful settlement. Black Rockfish mature at a relatively young age. This assessment uses a functional maturity relationship, updated from the previous assessment that used a biological maturity relationship.

No formal (e.g., inclusion of environmental indices) ecosystem considerations have been made given the lack of data for such an undertaking. McClure et al. (2023) report the climate vulnerability for several west coast groundfishes, including Black Rockfish. Black Rockfish demonstrated both high biological sensitivity and high climate exposure risk, to give it an overall high vulnerability score to climate change. This result should also be considered with the fact that like many rockfishes, periods of low productivity is not unusual to Black Rockfish and their extended longevity has historically allowed them to wait for advantageous productivity periods. Additional stressors such as fishing and climate change that possibly truncate longevity could bring significant challenges to population sustainability.

1.4 Historical and Current Fishery Information

Black Rockfish are harvested by a wide variety of fishing methods including trawling, trolling, and hook-and-line fishing with jigs and longlines. Black Rockfish are a major component of the nearshore fixed gear groundfish fishery, known as the “live-fish” fishery, and as incidental catch in the troll fishery for salmon and groundfish. Historically, Black Rockfish was a common, though more incidental, component of the trawl fishery. With the decline of salmon fishing opportunities in the 1980s and 1990s, Black Rockfish became a vital target of marine recreational fisheries in Oregon, especially during periods of restricted or slack fishing for salmon, halibut, and tuna.

Recreational harvest of Black Rockfish has become the dominant source of removals in Oregon. Since 1990, annual recreational harvests of Black Rockfish have averaged 315.1 tons off Oregon. Commercial annual harvests by non-trawl gear types during the same period averaged 127.9 mt in Oregon. Harvests by trawl on average during this period have been less than 3 mt annually.

Removal histories have been a significant axis of uncertainty in the past assessments of Black Rockfish. A major task for the 2007 assessments of Black Rockfish was developing a plausible reconstruction of historical landings of Black Rockfish and exploring the consequences of those landings. A formal commercial catch reconstruction had been recently developed in Oregon (Karnowski et al. 2014) for the 2015 Black Rockfish assessment, but was subsequently modified to better delineate trawl landings in Oregon and Washington (Cope et al. 2016). This current assessment also utilizes a recent recreational catch reconstruction (Whitman 2023).

1.5 Summary of Management History and Performance

Prior to 2000 the Pacific Fishery Management Council (PFMC or Council) managed the federal fishery for Black Rockfish as part of the *Sebastodes* complex, with no separate Acceptable Biological Catch (ABC) or Optimum Yield (OY) for Black Rockfish. Beginning in 2000, the Council established an ABC of 1,200 mt for Black Rockfish caught north of Cape Mendocino (in the Eureka, Columbia, and Vancouver INPFC statistical areas), but left Black Rockfish south of Cape Mendocino as part of the “other rockfish” category. Oregon Black Rockfish continued to be managed as a part of a complex following the 2003 and 2007 assessments, as these assessments combined the Oregon portion of the stock with California (Ralston and Dick 2003; Sampson 2007).

The 2015 Black Rockfish assessment split the stock into three units for modeling, including California, Oregon and Washington. This was the first time the Oregon Black Rockfish stock was assessed in isolation (Cope et al. 2016). Following the implementation of harvest specifications from this assessment, Oregon Black Rockfish were moved in 2017 for the

purpose of management to the Oregon Black/Blue/Deacon complex, and management of Black Rockfish within this complex has continued since that time. The ACLs/OYs have not been exceeded since this complex was created (Table 21).

In Oregon, Black Rockfish are allocated to both the commercial and recreational fisheries. The primary commercial fishery for Black Rockfish is the nearshore live-fish fishery, which is managed as a year-round fishery with annual state harvest caps and two-month trip limits. The commercial nearshore fishery first developed as an open access fishery in the early 1990s and transitioned to a State limited-entry permit system in 2004. The commercial nearshore fleet is composed of small vessels that target shallow nearshore rocky reefs using mostly hook and line jig or bottom longline fishing gear.

As noted above, Black Rockfish is the predominant species in the recreational fishery, which is managed using annual harvest guidelines, daily bag limits, and different types of spatial and temporal closures. Oregon had no recreational bag limits for marine fishes until 1976 when the state established a 25-fish limit. In 1978 the state established a daily limit of 15 fish for each angler's combined bag of rockfish, Cabezon and greenling, which stayed in effect until 1994 when the state established a 10-fish-per-angler daily bag limit specifically for Black Rockfish. Over time, the bag limits for marine species, of which Black Rockfish is the primary species, have gradually declined from 10 in 2000 to four or five fish in recent years (2017 - 2022). The use of in-season spatial closures to regulate effort have become common as well. In-season adjustments to regulations can be made more restrictive or less restrictive, depending on circumstances and the prospects for early attainment of harvest caps. Seasonal depth restrictions (e.g., inside 30 fathoms April 1 to September 30) are one tool used regularly in recent years to control the fishery, driven largely by the need to avoid bycatch of the primary rebuilding species, Canary Rockfish and Yelloweye Rockfish.

Some recent important regulations or management events include the following:

- 2000: Black Rockfish began to be managed by the Council as a minor nearshore species. Commercial trip limits were significantly reduced, with specific restrictions applying to Black Rockfish.
- 2002: Oregon adopted an Interim Nearshore Fishery Management Plan in anticipation of increased pressure on nearshore stocks due to reduced fishing opportunities for groundfish in federal waters. Regulations included fishing-sector specific caps on retained harvests, set approximately at the levels attained in 2000.
- 2003: The Council established Rockfish Conservation Areas (RCAs) to control catches of overfished rockfish species, and large portions of the shelf were closed to fishing. Differential trip limits were applied north and south of a management boundary at 40°10' N. latitude for nearshore *Sebastodes* species. Nearshore permittees in Oregon became subject to depth restrictions consistent with the shoreward non-trawl RCA boundary.
- 2004: The Oregon nearshore commercial fishery became a limited-entry permit system. Trip limits were established. The sport fishery in Oregon closed in September 2004

due to early attainment of the state's limit for sport-caught Black Rockfish. This was the first time that the sport rockfish fishery in Oregon had not been open all year.

- 2005: The recreational fishery in Oregon closed early.
- 2007: Black/Blue rockfish harvest cap was attained early and the nearshore commercial fishery closed for remainder of year.
- 2017: The recreational fishery in Oregon closed early.

1.6 Canadian and Alaska fisheries

Black Rockfish is one of multiple Inshore Rockfish species on the West Coast of British Columbia. The most recent evaluation was completed by Yamanaka and Lacko (-Yamanaka and Lacko (2001)), which determined that there was insufficient information to recommend a Black Rockfish-specific catch quotas for the five management areas on the Pacific coast of British Columbia. Black Rockfish continues to be a “Non-Quota” species in the Department of Fisheries and Oceans Management Plan (Fisheries and Canada 2014).

Directed fisheries for Black Rockfish in Alaska are limited. In the Westward region (Kodiak area) of Alaska, an age-structured assessment for Black Rockfish is under development by the Alaska Department of Fish and Game, using an acoustic survey as a primary source of data. No assessments have been developed for Black Rockfish for the Southeast or Central regions in the Gulf of Alaska.

2 Data and Model Inputs

A description of each data source is provided below (Figure 2), including removals (Figure 3), indices, and biological compositions.

2.1 Removals

2.1.1 Commercial Landings and Discards

Commercial landings and discards by fleet are shown in Table 1.

2.1.1.1 Background The previous Black Rockfish assessment (Cope et al. (2016)) used a number of sources of information to most accurately compile commercial catches of Black Rockfish. A comprehensive historical commercial catch reconstruction (Karnowski

et al. (2014)) had just been completed prior to the 2015 Black Rockfish assessment that provided gear-specific landings starting in 1892, but ODFW staff noted high year to year variability and unreasonably high magnitude in Black Rockfish trawl landings, primarily from the port of Astoria on the northern border with Washington and during the early years of the development of the trawl fishery (approximately 1940s – 1980s). Further, the Karnowski reconstruction did not account for fish captured outside of Oregon waters but landed in Oregon, as is the case with the majority of trawl landings for Black Rockfish in Astoria. As an alternative to the Karnowski reconstruction, ODFW reconstructed rockfish trawl landings in PFMC Area 3A, which includes both Oregon and Washington fishing grounds. These rockfish category landings utilized data sources with finer spatial delineation to separate landings among PFMC areas in Oregon than were used in the comprehensive reconstruction (Karnowski et al. (2014)). These include: FSUS reports for the Columbia River district (1940 – 1955), PFMC reports for Area 3A (1956 – 1977), ODFW estimated Astoria trawl landings documented in ODFW Marine Program data series report (Douglas (1998); 1978 – 1982) and ODFW landing receipts from Astoria (1983 – 1986). To separate Black Rockfish from rockfish market category landings in 3A, an aggregated proportion of Black Rockfish were used from trawl-specific species compositions available in Douglas (1998). From 1940 to 1975, an aggregated proportion from 1963 – 1975 was applied to rockfish landings (14.1%) and for 1976 – 1986, an aggregated proportion from 1976 – 1986 was used (6.8%). Finally, Black Rockfish in 3A were apportioned north and south of the OR-WA border using ODFW species composition data that provide the state area of the catch. This resulted in 98.6% of 3A Black Rockfish apportioned to WA during 1940 – 1986 in the previous assessment.

2.1.1.2 Current Methodology: Trawl Landings (1892 - 2022) While the above methodology was acceptable for the 2015 Black Rockfish assessment, there was a desire for a more equitable approach between Oregon and Washington for the current assessment. To that end, ODFW provided reconstructed 3A rockfish category landings to WDFW staff that were apportioned north and south of the OR-WA border using the above methodology for 1940 - 1986. This allowed each state to utilize a separate approach to delineate Black Rockfish-specific landings.

For Oregon-specific trawl 3A landings, the previous aggregated proportion of Black Rockfish were applied to the apportioned 3A rockfish landings (14.1% for 1940 – 1975 and 6.8% from 1976 – 1986). There are no other sources of information on historic trawl rockfish compositions and aggregating data across years increases available samples, though it does assume that compositions do not vary widely across time. This approach best utilizes the available species composition data from this time period. These 3A Black Rockfish landings were added to the reconstructed landings using an identical methodology for the other Oregon PFMC areas (2A, 2B and 2C) developed for the previous assessment for the complete trawl landings during the historical time period.

Trawl landings for Black Rockfish from 1892 – 1939 and from 1987 forward were not included in the reconstruction for the previous assessment. These values are available from the Karnowski reconstruction (Karnowski et al. (2014)) for 1892 – 1939 and are identical to

those used in the 2015 assessment. From 1987 - 2022, landings for Black Rockfish were obtained from PacFIN (accessed March 9, 2023). In addition to PacFIN landings, ODFW also reconstructed the URCK and POP categories in PacFIN from 1987 – 1999 (“Oregon department of fish and wildlife informational report regarding speciation of unspecified rockfish landings in oregon for inclusion in stock assessmetn time series of removals” (2017)). This reconstruction was not available for the previous assessment.

2.1.1.3 Current Methodology: Non-Trawl Landings (1892 - 2022) Non-trawl landings were provided by ODFW and by PacFIN, depending on the time period. Non-trawl landings from 1892 – 1986 era available from the Karnowski reconstruction (Karnowski et al. (2014)) and are identical to the 2015 assessment. Again, a combination of ODFW’s URCK/POP reconstruction (“Oregon department of fish and wildlife informational report regarding speciation of unspecified rockfish landings in oregon for inclusion in stock assessmetn time series of removals” (2017)) and PacFIN landings were used to estimate landings from 1987 – 1999. From 2000 – 2022, PacFIN landings were used exclusively.

2.1.1.4 Commercial discards The West Coast Groundfish Observer Program (WC-GOP) has monitored discards and landings coastwide to provide total catch mortality by species or species group since 2002. WCGOP estimates of Black Rockfish discards from 2002 - 2021 were used in this assessment to account for commercial discard mortality (obtained March 30, 2023). Fleet-specific rates were obtained by averaging the annual rate of discarded fish to the total mortality and this rate was applied to annual estimates of catch for each fleet. Trawl discard rates averaged 4.1% but were highly variable during this time period. Non-trawl discard rates averaged 0.9%.

2.1.1.5 Foreign Fishery Catches of Black Rockfish Rogers (2003) developed catch reconstructions for removals by foreign trawlers operating off the U.S. West Coast during the late 1960s to mid-1970s. Although this study reports that Japanese vessels operating in the Columbia and Eureka statistical areas (Oregon and northern California) caught substantial amounts of Black Rockfish, with cumulative catches of more than 500 mt over 10 years, it seems very unlikely that foreign vessels could have operated sufficiently close to shore to catch appreciable amounts of Black Rockfish. This assessment does not include Rogers’ estimates of foreign fleet removals of Black Rockfish, nor was it included in the previous assessment (Cope et al. (2016)).

2.1.2 Recreational Landings and Discards

Recreational landings and discards by fleet are shown in Table 1. Comparisons of the catch in each recreational fishery for the current and previous assessments are in Figure ??.

2.1.2.1 Historic Ocean Boat Landings and Discards (1973 - 2000) Recently, ODFW undertook an effort to comprehensively reconstruct all marine fish recreational ocean boat landings prior to 2001 (Whitman (2023)). Reconstructed catch estimates from the Oregon Recreational Boat Survey (ORBS) improve upon estimates from the federal Marine Recreational Fisheries Statistical Survey (MRFSS), which have known biases related to effort estimation and sampling (Van Voorhees et al. (2000)) that resulted in catch estimates considered implausible by ODFW. However, the ORBS sample estimates are known to lack the comprehensive spatial and temporal coverage of MRFSS. Addressing this coverage issue is a major part of this reconstruction. In general, the base data and methodology for these reconstructed estimates are consistent with recent assessments for other nearshore species (Cope and Whitman (2021); Langseth et al. (2021); Taylor et al. (2021); Wetzel et al. (2021)).

Prior to 2001, ORBS monitored marine species in both multi-species categories, such as rockfish, flatfish, and other miscellaneous fishes, and as individual species, such as lingcod or halibut. For this comprehensive reconstruction, four species categories were selected to reconstruct, including rockfish, lingcod, flatfish and miscellaneous, which constitute the bulk of the managed marine fish species. Black Rockfish are a component of the rockfish species category.

Category-level estimates were expanded to account for gaps in sampling coverage in two separate pathways. First, estimates from five major ports were expanded to include unsampled winter months in years lacking complete coverage. Expansions were based on available year-round sampling data and excluded years where regulations may have impacted the temporal distribution of catch. Second, all other minor port estimates were expanded to include seasonal estimates in years lacking any sampling based on the amount of minor port catch as compared to all major port estimates. A subset of landings were sampled by ORBS for species compositions within these categories. Once category-level landings were comprehensive in space and time, species compositions were applied for the three multi-species categories, including rockfish, flatfish and miscellaneous fish. Borrowing rules for species compositions were specific to the category and determined based on a series of regression tree analyses that detailed the importance of each domain (year, month, port and fishing mode) to variability in compositions.

Ocean boat estimates from 1979 – 2000 in numbers of fish of Black Rockfish from the above described methods were converted to biomass using biological samples from MRFSS. MRFSS biological data are available from 1980 – 1989 and 1993 – 2000. An annual average weight was applied to the total annual number of fish to obtain an annual biomass estimate of the landings. Several years missing biological data (1979, 1990 – 1992) were filled in using neighboring years or interpolation. MRFSS biological data were filtered for direct measured samples only. These landings in biomass were provided by ODFW and do not include an estimate of discarded fish. Landings during this time period average 309.9 mt and are of similar magnitude to the previous assessment (Cope et al. (2016)). Landings from 1973 to 1978 were initiated at zero in 1973 and ramped up to the initial biomass estimate from 1979. Bag limits in the recreational fishery during this time period (prior to 2001) are generally

liberal and ODFW staff recommended that no additional mortality of discarded fish be included from 1973 - 2000.

2.1.2.2 Modern Ocean Boat Landings and Discards (2001 - 2022) Recreational landings for ocean boat modes from 2001 – 2022 are available from RecFIN (accessed March 25, 2023). Both retained and discarded (released) estimates of mortality are included, though retained mortality contributes the vast majority to total mortality. Release mortality is estimated from angler-reported release rates and the application of discard mortality rates from the PFMC. The average proportion of Black Rockfish discarded was 0.92% from 2001 – 2022. From 2001 – 2022, total landings averaged 307.6mt, ranging from 198.7 to 458.6 mt. In 2022, estimated ocean boat landings were 374.3 mt.

2.1.2.3 Shore and Estuary Boat Landings (1892 - 2022) ODFW provided reconstructed estimates of shore and estuary landings for black rockfish from 1892 – 2022, using methodology similar to recent assessments (Langseth et al. (2021); Taylor et al. (2021); Wetzel et al. (2021)). Data sources include MRFSS and the Oregon Shore and Estuary Boat Survey (SEBS). Numbers of fish were provided by MRFSS from 1980 – 1989 and 1993 – June 2003, and by SEBS from July 2003 – June 2005. An annual mode-specific average weight was applied to numbers of Black Rockfish from 1980 – 1989 and 1993 – 2005. Separate weights were calculated for shore and estuary boat modes, and excluded extreme outliers and imputed values. This reconstruction also applied two scaling factors to remove bias towards freshwater sampling and underestimation of estuary boats, as detailed in (2018). To estimate Black Rockfish landings from July – December 2005, an expansion was developed using the three year average of the ratio between the first six months of the year and the total annual landings from MRFSS and SEBS landings from 2002 - 2004. Separate expansions were developed for shore mode and estuary boat modes.

Unlike the ocean boat recreational fishery, which developed during the 1970s, it was assumed that some shore and estuary fishing for Black Rockfish has occurred for many decades. As a result, the previous Black Rockfish assessment used fishing licenses to scale shore and estuary landings (Cope et al. (2016)). However, the link of Black Rockfish landings to fishing license sales is not well supported (pers. comm. A. Whitman, ODFW) and this assessment assumes a simple linear ramp from zero catch in 1892 to the estimated shore/estuary biomass of 28.1 mt in 1980, which produces a similar trajectory in catches to the previous assessment. Additionally, the ODFW does not currently sample shore and estuary boat fishing trips, and so a ten year average landing (1996 – 2005; 13.3 mt/year) was used to estimate shore and estuary boat landings during 2006 – 2022. Again, this methodology differed from the previous assessment and did not use fishing license sales to scale the removals but still produces a similar trajectory and magnitude of removals as the 2015 assessment shore/estuary removals. Shore and estuary boat landings for Black Rockfish are minor compared to recreational ocean boat removals but consistent. Shore and estuary boat landings average 17.2 mt annually from 1980 – 2003. To obtain an estimate of discarded fish for the shore and estuary fleet, the ocean boat proportion of discarded fish from RecFIN (0.92%; 2001 – 2022) was applied

to the estimated shore and estuary landings during 2001 – 2022. As with the ocean boat fleet, discard mortality was not estimated for shore and estuary landings prior to 2001, as bag limits were relatively high and discarding was assumed to be very limited.

As described above, recreational discard mortality for the ocean boat fleet was included from RecFIN from 2001 - 2022. Based on these landings data, an average discard rate of 0.92% was applied to the shore and estuary boat fleet to account for discarded fish from 2001 - 2022. No recreational discards were accounted for prior to 2001, based on ODFW recommendation that bag limits for Black Rockfish were high and discards were likely extremely low. There are no data that can speak to discard rates in the recreational fishery prior to 2001.

2.2 Length and Age Composition Data

Fish length measurements, from both the commercial and recreational fisheries, are one of the major sources of data for this assessment. Length composition data from the commercial and recreational fisheries in Oregon were included, as were age composition data from the commercial and recreational fisheries in Oregon. ODFW Port Biologists collect biological data and samples from commercial fisheries coastwide. These data are available from PacFIN (accessed JASON - ADD DATE HERE) and were processed using the PacFIN Utilities package (ADD CITATION OR OTHER INFORMATION). The vast majority of commercial biological data are gender-specific.

For the recreational fleets, prior to 2001, the length composition data are from the MRFSS, which was a federally funded program operating beginning in 1980 that collects information on the marine sport fisheries. The MRFSS program includes an intercept survey in which sport anglers are interviewed as they return from fishing trips, and where samplers can identify and measure the retained catches. The MRFSS sampling is intended to cover all forms of marine recreational fishing, including shore-based activities from beaches, jetties, and piers. In contrast the ORBS program that operates only in Oregon interviews and samples anglers operating from boats. The MRFSS length data, which are housed in the RecFIN database, generally do not indicate the sex of individual fish that were measured. In 2001, MRFSS was discontinued and ORBS became the primary source of recreational marine biological data. Similarly, the length data collected by the ORBS program does not generally indicate gender. All recreational biological data are available on RecFIN (accessed March 25, 2023).

2.2.1 Length Compositions

The length data for the assessment model were tabulated into 2-cm length bins ranging from 10 cm to 64 cm, with accumulator bins at each end.

The length composition data indicate some general differences among the fleets, with the trawl fisheries producing the largest fish, the ocean boat recreational fishery and the non-trawl fisheries producing fish of intermediate length and the shore fleet producing the smallest fish. There is little evidence in any of the length composition data of distinct modes or successions of modes from one year to the next that might represent strong year-classes.

2.2.1.1 Commercial - JASON - double check this section These data are from trawl and non-trawl fisheries from ports south of Astoria. The PacFIN dataset contains records for Oregon landings into Astoria; however, the majority of these samples are believed to have been landed in Washington waters and are used in the Washington model. Length composition data are reported in fork length. These data are expanded to reduce the effect of non-uniform sampling effort. The expansions are by weight, catch/sampled catch; first on a per-trip level, and then on a per-year, per-fishery level. Expansion factors have a minimum value of 1, and are capped at their 90th percentile value. The final sample size is the product of the two expansion factors, which is then capped at its 90th percentile value. The data were stratified by gender and fishery (Table ??). The final sample sizes were stratified and summed by length bin (10 cm to 60 64 cm bins, 2 cm in width), and an effective sample size is computed from the number of trips and number of fish each stratum represents, according to the Stewart and Miller method for multinomial fishery data. A small number of unsexed fish were present in the data; as these did not represent a distinct length distribution, they were excluded from the model.

2.2.1.2 Recreational Recreational length samples were obtained from two primary sources: MRFSS and ORBS (available on RecFIN). ODFW special project sampling data were considered but given the high sample sizes available from standard sampling, these data were not utilized. From 1980 – 1989 and from 1993 – 2000, the MRFSS program collected samples from both ocean and inland (estuary) areas ($n = 46,656$). ODFW provided MRFSS samples with the addition of a column that flagged length values imputed from weights to allow for selection of directly measured values ($n = 32,130$). From 1980 – 1989, total lengths (mm) were collected by MRFSS, which were converted to fork length. From 1993 – 2000, fork length (mm) was collected. Length samples from 2001 – 2022 from the ORBS sampling program are available on RecFIN ($n = 197,216$). All ORBS samples are by fork length (mm). The vast majority (95.5%) of these samples are from ocean trips ($n = 188,372$). Table 3 details sample sizes by year and recreational fleet.

2.2.2 Age Compositions

The fishery data for the assessment model consisted of otolith age-readings, mostly from the recreational fishery (ADD COMM AGE TABLE; Table 5). Age composition data were a subset of the length data, 41,212 records in total. The age composition data for the assessment model were tabulated into 1-yr age bins from 1 to 40 years. For the data tabulation provided

in this document, the accumulator bins were extended to compress and simplify display of the data.

The age composition data generally do not show much evidence of distinct year-classes that can be easily tracked from one year to the next, which suggests that there is not much recruitment variability from year-to-year or that age-reading error is sufficient to mask the appearance of strong year-classes. Age-at-length compositions were not expanded; the final sample sizes were set to 1 before tallying. The ages were modeled as conditional age-at-length.

2.3 Mean weights

Mean weights were used in the previous assessment (Cope et al. (2016)) but not developed for this assessment.

2.4 Fishery-Dependent Abundance Indices

Age and length composition data by themselves do not provide sufficient information to reliably determine trends in stock abundance and biomass. Most assessments of U.S. West Coast groundfish stocks rely on estimates of stock biomass from research trawl surveys to provide information on biomass trends, but Black Rockfish are very infrequently caught in any of the bottom trawl surveys, which have a limited coverage of shallow nearshore waters (none of the surveys have ever been conducted in waters shallower than 55 m). Multiple fishery-dependent indices are available for these assessments that are based on commercial or recreational catch-per-unit-effort (CPUE). For the commercial non-trawl fleet, an index of abundance based on retained catch from logbook data from the coastwide live-fish fishery was updated. For the recreational ocean boat fleet, an index of abundance from MRFSS interview level data was developed for the previous assessment (Cope et al. (2016)), covering 1980 - 1989 and 1993 - 2000. A second index from ORBS dockside sampling was updated for this assessment (2001 - 2022). Finally, a third recreational ocean boat index based on the Oregon At-Sea charter observer program was updated for this assessment, but ultimately not used in the reference model.

2.4.1 Commercial Index of Abundance - Nearshore Fishery Logbook CPUE

Only one abundance index is available from the Oregon commercial fisheries, one derived from a nearshore logbook CPUE time-series from a fixed gear fleet that operates coastwide. The ODFW has required nearshore commercial fishers (both nearshore permitted vessels and open access vessels) to submit fishing logbooks since 2004. Compliance is generally high,

averaging around 80%, but has varied through time ranging from 65% in 2007 to greater than 90% in recent years. Although required to provide all requested information in the logbook per fishing gear set, there has been substantial variation in the quantity and quality of information reported in logbooks. The logbook database contains information on catch by species (number or pounds of retained and released fish), effort (hook hours), sample location (port), date, vessel, fishing depth, fishing gear, fishing permit, number of fishers, and harvest trip limits.

Logbook information went through multiple data quality filters to attain the best possible consistent and representative data set through time to estimate a relative abundance trend. Individual observations of catch and effort were at the trip level, where multi-set trips were aggregated to the trip level. CPUE was calculated for each trip, where total catch was defined as the total of all reported retained catch (in pounds) and total effort was defined by hook-hours (number of hooks used multiplied by the number of hours fished). Previous iterations of this index utilized both released and retained fish; however, released fish were converted to weight by applying a median catch weight. This was deemed an unnecessary uncertainty for a Black Rockfish index, as few boats release substantial amounts of black rockfish. In general, data filters that were applied included eliminating records with missing or unrealistic values, including permitted trips using only hook and line jig gear from ports with appreciable data, and using only vessels that fished in at least three years over the logbook history. Vessel operators may have changed through time as the filter was tied to the vessel name only. Gear type was restricted to hook-and-line only (excluding longline gear) because this gear type accounts for the majority of trips for Black Rockfish (approximately 90.3% of unfiltered dataset). These filters are consistent with the most recent Black Rockfish assessment (Cope et al. (2016)) and, where applicable, other recent assessments that utilize the nearshore logbook index (Cope et al. (2019); Taylor et al. (2021)). In the final, filtered dataset, Black Rockfish are present at a high frequency (Table 6).

Covariates evaluated included year, month, port, depth, number of people onboard and the two-month Black Rockfish catch limit. A negative binomial model was fit to the trip-level data (catch with a log offset for hook hours). This full model was selected as the best fit model by AIC Table 7. However, when fit via MLE using the sdmTMB R package (version 0.3.0), this model had unacceptably large standard errors for the fixed parameters. Acceptable diagnostics were achieved by excluding the two-month trip limit covariate. The final model included all of the covariates of the full model with the exception of the two-month trip limit.

2.4.2 Oregon Recreational Indices of Abundance

The three recreational fishery abundance indices available for the Oregon regional assessment are summarized in Table ?? and Figure ?? . The sections below describe the underlying data and derivations of the indices.

2.4.2.1 MRFSS Dockside CPUE for Oregon, 1980 to 2000 This index was included in the 2015 Black Rockfish assessment but not updated for this assessment. For the RecFIN data from Oregon, the logistic regression analysis to select likely Black Rockfish trips was based on data from 6,165 charter-boat trips and a suite of 23 species (excluding Black Rockfish). The analysis generally produced large positive coefficients for shallow-water species that one would expect to co-occur with Black Rockfish (e.g., copper rockfish and blue rockfish), and large negative coefficients for deep-water or pelagic species that one would not expect to co-occur with Black Rockfish (e.g., Pacific halibut and coho salmon). Those trips having an estimated probability of producing a Black Rockfish that exceeded the cut-off value of 0.758 were selected for the CPUE analysis.

This cut-off value was chosen to balance the false-positives against the false-negatives and resulted in some trips that were estimated to be false positives, where Black Rockfish were caught, but should not have been, given the other species caught during those trips. These probably represent trips that fished in multiple locations, and thus caught a mix of shallow- and deep-water species. The screening also resulted in the inclusion of trips (false negatives) that should have caught Black Rockfish (given the other species), but did not. A total of 5,261 trips were selected for the CPUE analysis.

The MRFSS dockside standardized CPUE index for Oregon (Figure ??) was developed from the selected subset of the catch and effort data using GLMs, with a binomial model to estimate the probability of catching at least one Black Rockfish and a gamma model to estimate the magnitude of the positive catches per angler-fishing-hour. In all cases, the structural models had three main effects for the factors Year, Wave (bimonthly period) and Region (southern versus northern OR), and there were no interaction terms.

The annual index values were derived as the product of two components: predicted values for the probability of catching a Black Rockfish during a trip, and predicted values for the number of Black Rockfish caught by an angler per hour of fishing, given that at least one Black Rockfish was caught. This CPUE index for Oregon has a high amount of inter-annual variation, particularly in the early part of the time-series. The index shows a fairly steady upward trend starting from 1987.

2.4.2.2 ORBS Dockside CPUE for Oregon, 2001 to 2022 Trip-level catch-per-unit-effort data from ORBS dockside sampling was obtained from ODFW. To mitigate the confounding of hourly effort associated with these trips with travel, the travel time was subtracted from the hours fished. Travel time was stratified by boat type (charter and private) and was calculated as boat type-specific speeds (13 mph for charter boat trips and 18 mph for private boat trips) multiplied by twice the distance between the port of origin and the reef that was fished. CPUE, expressed in terms of fish per angler-hour, was calculated by multiplying the number of anglers and the adjusted travel time. The database contains information on catch by species (number of retained fish), effort (angler hours), sample location (port where data were collected), date, bag limits and other relevant regulations, boat type (charter or private), and trip type (e.g., bottom associated fish).

The unfiltered data set contained 414,587 trips. Multiple standardized filters are applied to ORBS trip-level data to remove outliers and data unsuitable for an index. These filters include trips with incorrect interview times, which impact calculation of effort, unreasonably long or short trips, and retaining only bottomfish target trips. Further filters were utilized for fishing closures (i.e. temporal or spatial closures) and catches exceeding bag limits, which would presumably impact catch rates. An alternative configuration of the dataset included a binary variable to identify trips that hit the bag limit to include as a covariate, rather than a filtering criterion, and did not filter out trips that had reached the daily bag limit. Trips from several ports with extremely small sample sizes (<1% of total trips) were also excluded and finally, trips that met criteria for irrational effort reporting (i.e., implausible values) or extreme catch rates were excluded as well. The final dataset included 96,707 trips. The previous assessment only included trips onboard charter vessels (Cope et al. (2016)), but this filter was not retained for this updated index. Instead, boat type was evaluated as a covariate. In the final, filtered dataset, Black Rockfish are present at a high frequency (Table 8).

Covariates evaluated included year, month, port, the open depths to fishing (all depths or inside 20/30/40fm), boat type and the daily bag limit for Black Rockfish. A negative binomial model was fit to the trip-level data (catch with a log offset for adjusted angler hours). This full model was selected as the best fit model by AIC Table 9. However, when fit via MLE using the sdmTMB R package (version 0.3.0), this model did not converge. Acceptable diagnostics were achieved by excluding the bag limit covariate. An additional sdmTMB model was fit that used the alternative configuration of the dataset, and included all of the full model covariates but replaced the bag limit covariate with the binary variable of whether the bag limit was reached on the trip. While acceptable diagnostics were achieved with the alternative full model, there was little effect on the resultant index and for parsimony, the final model included all of the covariates of the original full model with the exception of the daily bag limit.

2.4.2.3 On-Board Observer CPUE for Oregon, 2001 and 2003 to 2022

Drift-level catch-per-unit-effort data from ORBS at-sea charter observer sampling program was obtained from ODFW. The database contains information on catch by species (number of retained and released fish), effort (angler hours), sample depth, and bag limits and other relevant regulations.

The unfiltered data set contained 12,344 drifts. Multiple standardized filters are applied to the drift level data to remove outliers and data unsuitable for an index. These filters were co-developed with SWFSC staff (Monk et al. (2013)). These filters include drifts without data needed for CPUE information, long drifts (below 95th quantile), drifts in deeper waters (less than 64fm, 99th quantile), drifts that were targeting primarily mid-water species, and drifts outside of the legal fishing depth (with a five fm buffer). Additionally, drifts with recreational longleader gear were removed, as Black Rockfish is not a legal longleader species in Oregon, and years with less than 100 samples were also removed. This included 2020 and 2021, during which the COVID-19 pandemic dramatically limited the ability to sample

onboard charter vessels. Finally, drifts onboard charters from Port Orford were removed due to small sample sizes.

Covariates evaluated included year, month (or two-month wave), port, the open depths to fishing (all depths or inside 20/30/40fm), and a 10fm-binned depth of drift covariate. A negative binomial model was fit to the drift-level data (catch with a log offset for adjusted angler hours). A model without the open fishing depths was selected as the best fit model by AIC Table 10. However, when fit using the sdmTMB R package (version 0.3.0), the QQ plot indicated heavily skewed data and acceptable diagnostics were not achieved for any model explored. As a result, and given an alternative recreational ocean boat index with much larger sample sizes overall (ORBS), this index was not recommended for use in this assessment and subsequently not included in the reference model.

2.5 Fishery-Independent Data

2.5.1 Abundance Indices

2.5.1.1 Tagging Study Estimates of Abundance off Newport, OR, 2002 to 2014

Details of the tagging study are available in Krutzikowsky et al. (2019) and the implementation in the last stock assessment in Cope et al. (2016). Briefly, in a study that started in 2002 and concluded in 2014, the ODFW used Passive Integrated Transponder (PIT) tags to mark 2,500 to 4,000 black rockfish annually off Newport, OR. Marked fish were recovered from recreational fishery landings, with sampling focused on the charter vessel fleet. Approximately 80% of the annual landings were sampled for marked fish, resulting in the recovery of 3,263 marked fish to date.

The multi-stage mark-recovery model used to estimate annual survival and recovery rates for the black rockfish population off Newport was similar to “Model 0”, as described in Brownie et al. (1985), except that the recovery rates after the initial year at liberty were held constant. This particular tagging model configuration was selected because it provided a better AIC score than other models that were evaluated. It allows direct (first-year) recovery rates to differ from recovery rates of previously marked cohorts, which appeared to be the case in the black rockfish mark-recovery data. Model 0 parameters were then used to calculate annual exploitation rates, which were then applied to the annual landings to estimate annual abundance.

The basic approach is to estimate the numbers of fish from the equation $N_y = C_y/u_y$, where C_y is the catch (in numbers of fish) in year y , N_y is the population abundance at the start of the year, and u_y is the exploitation rate. As in Cope et al. (2016), u_y can be estimated from the ratio of the estimated recovery rate (\hat{f}_y) times C_y divided by the number of fish sampled for marks (cs_y). The C_y appearing in the numerator of the equation for N_y cancels with the C_y in the numerator of the equation for \hat{u}_y , leaving as the following estimator for

$\hat{N}_y = cs_y/\hat{u}_y$. Note that cs_y is the number of fish checked for marks, which is known without error in this study. Approximate estimates of variance for the \hat{N}_y values were derived from the delta method.

$$var[\hat{N}_y] \approx [(cs_y)^2/(\hat{f}_y)^4] * var[\hat{f}_y]$$

2.5.1.1.1 Spatial Coverage of the Oregon Tagging Study off Newport

One feature of the 2015 Oregon assessment model that is somewhat unique is the use of a prior probability distribution for the catchability parameter associated with the tagging study estimates of abundance of exploitable Black Rockfish off Newport. The analysis of habitat area coupled with CPUE fish densities indicates that on average 12.7% of the exploitable portion of the Black Rockfish population off Oregon reside in the waters off Newport. The lognormal prior distribution was assumed to have a standard deviation of 0.5, which is more than double the between-port variability calculated from the available CPUE data (CV=0.157). Although trip-level variability in CPUE is typically much larger than 50%, most of the variation in CPUE is due to variability in catchability rather variability in the abundance of the fish. A history of how this prior was established is available in Cope et al. (2016).

2.5.1.2 Nearshore Acoustic-Visual Survey

Oregon conducted the first statewide combined acoustic-visual survey of nearshore semi-pelagic rockfish in 2021 (Rasmussen et al. 2022). Background details on survey design, data processing, and analyses can be found Rasmussen et al. (2022a) and Rasmussen et al. (2022b), including an independent validation of survey methodology (i.e., test area population estimates from the combined acoustic-visual survey were similar to estimates from the overlapping Black Rockfish pit tagging experiment (Krutzikowsky et al. 2019)). A brief overview of the survey design and the analysis of acoustic data are provided below.

Survey sampling was designed using a systematic sampling layout with parallel transects. Parallel (perpendicular to the north south axis) transects were evenly spaced in Oregon, starting from the Washington border in the north extending to the California border in the south, after random selection of the location for the most northerly transect. Transects were identified as “full” (spaced every 15 km and extending from the 80 m contour shoreward to the surf zone, ~0 m contour) and “rock” transects (spaced every 1 km where “hard” bottom habitat had previously been identified; see Rasmussen et al. (2022)). The length of each rock transect was equal to the east to west length of each hard habitat cluster plus a 0.5 km buffer applied around the cluster.

The primary survey vessel was the RV Pacific Surveyor, a 17 m retired commercial fishing vessel converted for research purposes. The secondary vessel, used for shallow water applications, was the RV Arima, a 4.9 m recreational fishing vessel converted for nearshore research in depths shallower than the primary vessel could achieve. Acoustic data were collected

using two downward-facing, pole-mounted transducers (BioSonics DT-X split beam Scientific Echosounders: 38 and 201 kHz). Acoustic data were informed using video collected with the Benthically Anchored Suspended Stereo Camera (BASSCam). Video deployment locations were selected *in situ* by the scientist monitoring the acoustic data in real-time. Twice daily hook and line sampling was also conducted.

A linked Online Map of the survey is available showing locations of all sampling events. A total of 595 acoustic transects were conducted and paired with 496 video drops and 48 fishing stations from 1 August to 9 September 2001. A total of 35,839 fish were identified in the video and 869 fish captured using hook and line. Acoustic target-strength to length conversions were conducted using an aggregate of all video data from the survey. Video drops conducted >50 m from acoustically observed fish were used to predict how many fish were missed by the acoustics. Fish counts were converted to biomass using the length-weight relationship from the hook and line data.

Acoustic data were analyzed using both echo integration and echo counting. Echo integration conversions were done using a combination of target-strength length relationships from the literature (following recommendations from the PFMC Science and Statistical Committee, SSC, methodology review). Acoustic data within 1 m of the bottom (acoustic exclusion zone) were corrected for using data from above the deadzone. Schools of fish were observed over rocky habitats at all of the major nearshore reefs. Additional small patch reefs were also observed interspersed throughout the survey area.

A simple design-based population abundance estimate was generated by averaging the echo integration density of the acoustics in each habitat type (rock, gravel and sand) and multiplying it by the total area of each of those habitat types in three area-based strata (North, Central, and South) spanning the Oregon coast from the Washington border to the California border. Biomass in each area was then calculated by multiplying the estimated abundance by the proportion of fish in each 1 cm bin length class and then converting those length bins to weight using the length to weight relationship obtained from the hook and line data. After summing by region, the result is an independent estimate of coastwide abundance (and biomass) in Oregon waters in 2021. The coefficient of variation of the biomass estimate was also calculated, and it was used to generate the standard error for the assumed lognormal error structure of this data when integrated into candidate stock assessment models.

2.5.1.3 Marine Reserves Hook and Line Survey The Marine Reserves program at ODFW monitors Oregon's marine reserves. A standardized hook and line survey, modeled after the California marine reserves survey, collects drift-level catch and biological data in both the marine reserves and nearby comparison areas. There are four marine reserve sites (Redfish Rocks, Cape Perpetua, Cascade Head, and Cape Falcon) that span Oregon's coastline where the hook and line survey is a suitable tool for monitoring. These comparison areas are pre-selected areas with similar bathymetry and habitat characteristics as the reserve but are open to fishing. Each reserve has at least one comparison area, but several have more than one that are monitored regularly along with the reserve. A 500-meter square

grid overlaid on the area defines the sampling units or cells. Cells are randomly selected within a marine reserve or comparison area for each sampling event. Three replicate drifts are executed in each cell. The specific location of the drifts within the cell is selected by the captain. Data are aggregated to the cell-day level and subsequently aggregated to counts of fish per cell-day as the metric for CPUE. Sampling occurs in the spring and late summer/fall seasons. Over time, cells without appropriate habitat for the focus species, mainly groundfish, have been removed from the selection procedures, and those presented in this dataset include only those that are currently “active”. The number of cells sampled in a day can vary slightly and range from three to five. Sampling was not conducted in 2020 or 2021 due to the COVID-19 pandemic and this survey’s reliance on paid volunteers.

The dataset contains information on catch by species (number of fish), effort (angler hours), reserve site, area of sampling, treatment (reserve or comparison area), and date of sampling. Additional information that are available include proportion of rock habitat at the cell-level, water temperature, and the average drift depth, though these data are not available for the entire dataset. Average depth (fm) was binned to 10-fm bins (0 – 50fm), and those shallower than 10fm were excluded to a small sample size. April – June was aggregated to a spring season and August - October to a fall season. Black Rockfish are commonly encountered in this survey (Table 11).

Covariates evaluated included year, month (or season), site, treatment (reserve or comparison area), and depth bin. A negative binomial model was fit to the cell-day-level data (catch in numbers of Black Rockfish with a log offset for angler hours). Two full model series were considered, one with all the potential covariates with month and one with season. For the series with month, the full model was selected as the best fit model by AIC, but month was almost non-significant. For the series with season, the full model was again selected as the best fit by AIC (Table 12). When fit via MLE using the sdmTMB R package (version 0.3.0), acceptable diagnostics were achieved for both full models. However, the error around the point estimates was reduced with the season covariate and ultimately, this was the model selected to be used in the reference model. Previous assessments for nearshore species have explored the use of this survey as a raw index of abundance (Taylor et al. (2021), Wetzel et al. (2021), Langseth et al. (2021), Cope et al. (2019)) but to date, this index has not been a part of a reference model. Length compositions from this survey are also included.

2.6 Biological Data and Parameters

The major biological inputs to the models are natural mortality, age and growth parameters, weight-length, maturity and stock-recruitment parameters. The following sections outline the treatment of each section.

2.6.1 Natural Mortality

Natural mortality is a critical parameter that drives much of the outcome of stock assessments. This value is not directly measured for black rockfish, so it either needs to be estimated or fixed in the model. Prior treatments have either used fixed ramps from lower to higher female natural mortality values (0.16 to 0.24 for females (2007 assessment); 0.17 to 0.2 (2015 assessment)) to constant male natural mortality value (0.16 in 2007; 0.17 in 2015). Females rapidly disappear from the population after 20 years of age, whereas males can still be found in their 30 and 40s, with the oldest individuals along the coast aged at 56 years (Love 1957). Females are rarely found in their 30s and males in their 40s in Oregon.

The reason for the lack of females has been debated for many years. The “hide them” (using age-based selectivity curves to hide older females) or “kill them” (using the above mentioned ramps of death to account for no older females in samples) was specifically considered since the last assessment among researchers from California to Alaska, and it was agreed that the “hide them” hypothesis is the least feasible situation (see Rasmussen et al. (2023) for a specific study that went looking for old females). It was also agreed a constant natural mortality rate should be used for this assessment.

Determining reasonable natural mortality values is also challenging as the quick disappearance of females from the population after 20 years old challenges typical biological assumptions, especially since black rockfishes have been the focus species when developing the theory of big old fat fecund female contributions to spawning output (Bobko and Berkeley 2004b; Hixon et al. 2014b). In a study confirming the advanced capacity for output of older females (Berkeley et al. 2008) the oldest aged females in the study were under 20 years, so the enhanced reproductive capacity, despite the loss of females after 20 years of age, is still intact.

Using the Hamel and Cope (2022) longevity-based estimator of natural mortality as implemented in the natural mortality tool (2022), the following M values correspond to the longevity estimates:

- 0.108 at 50 years
- 0.135 at 40 years
- 0.180 at 30 years
- 0.216 at 25 years
- 0.270 at 20 years

These provide reasonable bookends for likely natural mortality values for black rockfish. For females, estimates based on the von Bertalanffy growth function range from 0.27-0.32 and for males, 0.34 to 0.38. Those estimates are on the very high side, and thus are not considered further.

Exploratory runs first attempted to estimate natural mortality with a high bound of 0.25, but this bound was always achieved. This consistent attribute of unconstrained models wanting

high natural mortality rates argued for the need to fix natural mortality, as was done in the 2015 assessment. For this reason, the constant natural mortality values of 0.19 for females (within the ramp of 0.17 and 0.2 used in that assessment) and 0.17 for males (same as the last assessment) were used. A likelihood profile across the above mentioned range of natural mortality values, but maintaining the above ratio of female to male natural mortality, is included to explore model sensitivity and demonstrate the models desire to push the upper bounds of realistic values.

2.6.2 Growth (Length-at-Age)

2.6.2.1 Age validation Fish ages estimated for age-based stock assessment models need to be accurate and precise because ageing error can affect all model input data. Current protocol for ageing groundfish is to count growth marks on break-and-burn preparations of sagittal otoliths (*Manual on generalized age determination procedures for groundfish* 2006) but interpreting the growth marks can be challenging and lead to errors in ageing. To validate the break-and-burn ageing method for Black Rockfish off Oregon, researchers at ODFW and the University of Alberta used secondary ion mass spectroscopy (SIMS) to determine oxygen isotope ratios (^{18}O) in Black Rockfish sagittal otoliths. Because an otolith is acellular, metabolically inert, and grows throughout the life of the fish, any elements accreted onto its surface are permanently retained (Campana 1999). Otolith oxygen isotopes exist in equilibrium with ambient seawater (Iacumin et al. 1992) and otolith ^{18}O is inversely related to water temperature and is a function of salinity (Høie et al. 2004); therefore, sequential ^{18}O measurements across the otolith are a proxy for seasonal temperature cycles experienced by the fish throughout its lifespan.

SIMS proved to be effective for producing detailed otolith ^{18}O chronologies with adequate temporal resolution for identification of seasonal trends in water temperatures. Locations of annuli counted during break-and-burn ageing of Black Rockfish sagittae generally corresponded to maxima in the ^{18}O chronologies, inferring formation during periods when water temperatures were relatively cold. Generally, chronologies were characterized as having one strong signal peak and trough between annuli; however, ^{18}O chronologies could be highly irregular and chaotic between maxima, inferring vertical and lateral movements known to occur in this species, as well as prolonged larval duration and settlement in shallow nearshore areas with short-term variation in water temperatures and salinities . The break-and-burn method of ageing was validated for Oregon Black Rockfish up to 12 years old (Terwilliger et al. 2023).

2.6.2.2 Age and growth relationship The length-at-age was estimated for female and male Black Rockfish using data from collections sampling the commercial (trawl and non-trawl) and recreational fisheries off the coast of Oregon (Figure 4 and Figure 5), with all lengths in fork length and all ages in years. Figure 6 shows the predicted von Bertalanffy

growth function (VBGF) fits to the data. Females grow larger than males and sex-specific growth parameters were estimated at the following values:

$$\text{Females } L_{\infty} = 48.82 \text{ cm; } k = 0.18 \text{ per year; } t_0 = -2.00$$

$$\text{Males } L_{\infty} = 44.88 \text{ cm; } k = 0.21 \text{ per year; } t_0 = -2.21$$

The coefficient of variation of length by age fluctuated around 0.06 to 0.1 for the most well sampled ages and was similar for each sex (Figure 7). When estimated in the models, these same values would often be produced, but it was ultimately determined it is more parsimonious to fix to 0.1 for both sexes.

The estimated VBGF parameters provided initial values for the estimation of growth in the model, as all age and length data are included in the model. The t_0 value was fit to zero in the reference model in order to achieve more realistic L_{∞} and k values. The resultant growth curves estimated by the model are presented in Figure 39. Sensitivity to fixing the growth parameters to the external values or estimating t_0 are explored through sensitivity analyses.

2.6.3 Ageing Bias and Precision

Counting ages from ageing structures in long-lived, temperate fishes is challenging. Ages derived from these structures can be hard to reproduce within and between readers (i.e., imprecision), and may not contain the true age (i.e., bias). Stock assessment outputs can be affected by bias and imprecision in ageing, thus it is important to quantify and integrate this source of variability when fitting age data in assessments. In Stock Synthesis, this is done by including ageing error matrices that include the mean age (row 1) and standard deviation in age (row 2). Ageing bias is implemented when the inputted mean age deviates from the expected middle age for any given age bin (e.g., 1.75 inputted versus 1.5 being the true age); ageing imprecision is given as the standard deviation for each age bin (row 2).

The Oregon Department of Fish and Wildlife has a long history of ageing black rockfishes. Because of this long tenure, the main age reader has changed several times over the years. There are six distinct age readers, but the most current age reader has concluded a comparison among all of the other age readers. The most recent reader is considered the least biased. This assumption enables the construction of ageing error matrices relative to the most recent reader, thus resulting in 6 different ageing error matrices for each period:

- up to 2000: Reader 1 vs Readers 5 & 6
- 2001: Reader 1 vs Reader 5
- 2002: Reader 1 vs Reader 4
- 2003 to 2009: Reader 1 vs Reader 3

- 2003 to 2016 (recreational only in 2016): Reader 1 vs Reader 2
- 2016 (commercial) to 2022: Reader 1 vs Reader 1

Estimation of ageing error matrices used the approach of Punt et al. (2008) and release 1.1.0 of the R package nwfscAgeingError (Thorson et al. 2012). The ageing error matrix offers a way to calculate both bias and imprecision in age reads. Reader 1 is always considered unbiased, but may be imprecise. Bias relative to the primary reader is given for the second or additional readers. Several model configurations are available for exploration based on either the functional form (e.g., constant CV, curvilinear standard deviation, or curvilinear CV) of the bias in the second read or reader or in the precision of the readers. Model selection uses AIC corrected for small sample size (AICc), which converges to AIC when sample sizes are large. Bayesian Information Criterion (BIC) was also considered when selecting a final model. Table 13 through Table 15 provide model selection results.

Age bias plots are provided in Figure 9. The calculated bias relationships from the best fit model are shown in Figure 11 and show only small biases between readers. Figure 10 shows the imprecision estimates of the best fit models. In the cases where two models were similarly selected, the least extreme form (i.e., drastic drops or increases in imprecision) were chosen for the final matrix. Each ageing error matrix was then applied to the appropriate time and fleet combination.

2.6.4 Length-Weight Relationship

The length(cm)-weight(kg) relationship for Black Rockfish was estimated outside the model using biological data available from the Oregon commercial and recreational fisheries (Figure 16). The resultant relationship is very similar for both males and females, and is very close also to what is seen in the state of Washington (Figure 17). The estimated length-weight relationship for female fish was $W=2.54537e-05L^{2.9}$ and males at $W=2.56318e-05L^{2.89}$.

2.6.5 Maturation and Fecundity

Black Rockfish maturity was assumed to be based on length, as in past assessments. This assessment used functional maturity instead of biological maturity to describe the maturity schedule. Functional maturity was classified by a more stringent definition of maturity that considered abortive maturation (delayed participation in reproductive event), skipped spawning (mature individuals forgo spawning), and level of follicular atresia as opposed to biological maturity that only considers physiological development. Functional maturity included the biologically mature individuals that were actually expected to contribute to spawning in a given year.

Claire Rosemond (Oregon State University) and Melissa Head (Northwest Fisheries Science Center) provided estimates of both biological and functional maturity for Black Rockfish sampled in 2014 to 2021 from September through April (the time period that includes yolk development and spawning). Samples were collected for fish caught in Oregon and Washington waters by biologists at Oregon State University, Oregon Department of Fish and Wildlife, and Washington Department of Fish and Wildlife. Biological maturity and functional maturity observations were fitted in separate models. Biological maturity and functional maturity status observations (0 = immature and 1 = mature, n = 644) were fitted in a logistic regression model (`glm` function, `family = binomial`, `link = "logit"`) and flexible spline model (Head et al. 2020). The estimated model parameters were used to calculate length at 50% maturity (L50%; Table 16) and maturity ogives (Figure 12). The delta method was used to calculate 95% confidence intervals for estimated L50% in the logistic regression and a bootstrapping method was used to calculate 95% confidence intervals for L50% in the flexible spline model.

A flexible spline model can capture skipped spawning in the maturity ogive by allowing a decreased asymptote from 1.0. There was evidence of skipped spawning in larger size classes and so the flexible spline model was determined to be the best representation of the reproductive biology of Black Rockfish for the sampled time period (Figure 13). The maturity vector input into the model as a fixed relationship is shown in Figure 14. Sensitivity of model output to the use of the biological and functional logistic relationships are explored.

The Black Rockfish fecundity-at-length relationship was provided by E.J. Dick (SWFSC) and based on the work from Dick (2009). The fecundity relationship was estimated equal to $Fec=1.41e-08L^{4.68}$ in millions of eggs where L is length in cm. Fecundity-at-length is shown in Figure 15.

2.6.6 Stock-recruitment function and compensation

The Beverton-Holt stock-recruit model (Beverton and Holt 1957) has been the traditional recruitment function for rockfishes and is assumed for black rockfish. Specifically, the re-parameterized Beverton-Holt that uses a steepness parameter defined as the proportion of average recruitment for an unfished population expected for a population at 20% of unfished spawning output (Mace and Doonan) was used in these assessments. This is a notoriously difficult parameter to estimate, thus several attempts to derive a prior of steepness have been attempted (Myers et al. 1995; Dorn 2002). The Thorson-Dorn rockfish prior (developed for use West Coast rockfish assessments) was reviewed and endorsed by the Scientific and Statistical Committee (SSC) in 2017, and is the primary source of information on steepness for west coast rockfishes. The prior (h ; beta distribution with $\mu=0.72$ and $\sigma=0.15$) is used in this assessment, but attempts to estimate steepness were not successful, so it is fixed and its influence is explored via a likelihood profile.

2.6.7 Sex Ratio

No information on the sex ratio at birth was available so it was assumed to be 50:50.

2.7 Environmental and Ecosystem Data

This stock assessment does not explicitly incorporate trophic interactions, habitat factors or environmental factors into the assessment model. More predation, diet and habitat work, and mechanistic linkages to environmental conditions would be needed to incorporate these elements into the stock assessment and should remain a priority. McClure et al. (2023) report the climate vulnerability for several west coast groundfishes, including Black Rockfish. Black Rockfish demonstrated both high biological sensitivity and high climate exposure risk, to give it an overall high vulnerability score to climate change. This result should also be considered with the fact that, like many rockfishes, periods of low productivity is not unusual to Black Rockfish and their extended longevity (though admittedly this seems shorter than previously believed and should be reconsidered) has historically allowed them to wait for advantageous productivity periods. Additional stressors such as fishing and climate change that possibly truncate longevity could bring significant challenges to population sustainability.

2.8 Data Considered but Not Included

Special projects data for Black Rockfish were reviewed but ultimately, with the large number of commercial and recreational samples available under standard sampling protocols, these data were not used in the reference model. There were also multiple sources of fishery-independent information that ODFW has provided to assessments in recent years as ancillary data that were considered. These included ROV survey data, video lander survey data and the Marine Reserves longline survey data. ODFW ROV data were reviewed by an SSC methodology review in 2020 (see PFMC September 2020 briefing book; Agenda Item D.4 Attachment 1) and used to independently estimate population size for Quillback Rockfish in Oregon (Langseth et al. (2021)). Video lander data from multiple projects was recently summarized by Marine Resources Program staff (Rasmussen et al. (2020)). These data were used to develop density estimates and independent abundance estimates for the most recent assessments for both Quillback Rockfish (Langseth et al. (2021)) and Lingcod (Taylor et al. (2021)). These two datasets were not included, as they were not designed specifically to survey Black Rockfish and the ODFW Nearshore Acoustic-Visual survey was regarded as a stronger estimate of the scale of the Black Rockfish population scale without the major caveats that come with using the ROV or lander data. Finally, the Marine Reserves longline survey data collected in Oregon's marine reserves were available from 2015 - 2019. While Black Rockfish are encountered in this survey, sample sizes are lower than the Hook and Line survey and the gear type is not ideal for Black Rockfish.

3 Assessment Model

3.1 Summary of Previous Assessments and Reviews

3.1.1 History of Modeling Approaches Used for this Stock

3.1.1.1 Black Rockfish South of Cape Falcon The first stock assessment of black rockfish off Oregon (Stewart 1993), which was limited in geographic scope to the northern portion of Oregon, was a Cohort Analysis based on age composition data collected from fish landed at Garibaldi. The first comprehensive analysis of the black rockfish stock off Oregon and California was by Ralston and Dick (2003), who developed a statistical catch-at-age model using Stock Synthesis. Sampson (2007) used a similar model configuration and approach.

In the 2007 assessment model the data were organized into three basic gear-types (Hook-and-Line, Trawl, and Recreational), the data from Oregon and California were kept separate, and the tuning indices were recreational angler CPUE series based on the same or similar data sources (MRFSS for both states, ORBS for Oregon, and CPFV surveys for California). Fishing effort was measured in terms of angler-days rather than the angler-hours metric used in the current California and Oregon regional assessment models. The 2007 assessment used the ODFW tagging study estimates of black rockfish abundance off Newport as a relative abundance index. Those data were unavailable for the 2003 assessment. The 2007 assessment also used a juvenile rockfish pre-recruit index, which was unavailable for the previous assessment.

The landings data series in the 2007 assessment differed quite substantially from the series developed by Ralston and Dick for the 2003 assessment. Neither of those assessments attempted to account for discards, instead assuming that discards were negligible.

3.1.1.2 Black Rockfish North of Cape Falcon Three full assessments for black rockfish, conducted in 1994, 1999, and 2007, modeled the black rockfish population found in coastal waters between Cape Falcon, Oregon and north to the U.S./Canadian border (Wallace and Tagart 1994; Wallace et al. 1999, 1999; Wallace and Tsou 2007). There have been no update assessments for black rockfish resources.

The 1994 assessment utilized a Stock Synthesis model configuration, with two auxiliary data sets as black rockfish abundance indicators, one based on tagging CPUE and one on based coastal recreational bottomfish directed effort (Wallace and Tagart 1994).

Wallace et al (1999) constructed an assessment model by using the AD Model Builder software (ADMB; (Fournier 1997)) to assess black rockfish abundance. Three key features of the 1999

model were (1) the parameterization of the expected catches at age, (2) the definitions of the sampling units for the different types of data inputs, and (3) the integration of tagging data explicitly. The parameterization chosen mostly affected parameter bias whereas the sampling unit designation mostly affected estimator variance. Both bias and variance were components of overall parameter uncertainty. The parameterization and the sampling unit definitions were both designed to conform to the actual sampling protocol used, thereby propagating sampling uncertainty through to the final biomass estimates.

The 2007 assessment (Wallace and Tsou 2007) employed Stock Synthesis 2. Unlike the 1999 assessment, CPUE from the tag release trips and Petersen tagging study abundance estimates were included as relative abundance indices.

3.1.1.3 California, Oregon, and Washington Assessments in 2015 The 2015 assessment defined three distinct stocks for assessment. Each stock matched the state boundaries of California, Oregon and Washington. All assessments used the Stock Synthesis 3 version 3.24V. The Oregon model had five fisheries (three commercial and two recreational) and considered five surveys. A research fleet was also designated in order to use ages for growth estimation. There were four primary data likelihood components for survey indices, mean weights, lengths and ages. Fits to catches also contribute to the total likelihood, but is typically very small. The models was tuned using the Francis (2011) method for biological compositions and added variance for survey indices. Mean weights were not tuned. Recruitment deviations were not estimated. Natural mortality was treated as constant for males and used a step function (with a breakpoint at age 10) for females that was also constant over years. Age-based dome-shaped selectivity in the ocean boat-based recreational fishery was also implemented for females.

Results for the Oregon assessment of black rockfish in 2015 estimated stock status in that year at 60.4%, and never showed a decline below the target biomass. This assessment was highly constrained by the choice to treat the catchability of the tagging index as known, thus the uncertainty in the assessment is very low.

3.1.2 Most Recent STAR Panel and SSC Recommendations

The STAR panel identified the following issues as sources of major uncertainty:

- Natural mortality, especially in females. There is no data to differentiate whether the missing older females are dying or are avoiding capture. The choice between using a constant (as used in the California and Washington assessments) or step function (used in Oregon) is also an point of uncertainty.
- The level of cryptic biomass. This is a result of using dome-shaped selectivity to explain the absence of old females.

- Uncertainty in historical catch, especially in the historical trawl fishery.
- Acknowledging that there remains uncertainty in the stock-recruit relationship parameters (particularly the Beverton-Holt steepness parameter).

The Oregon model faced many challenges and ultimately was re-evaluated in a secondary review panel. That panel noted the following issues and recommendations (in addition to those above).

- The lack of recruitment estimation should be re-evaluated despite the last assessment indicating no apparent information contained in the biological data as to notable recruitment deviations. There was concern recruitment was being driven by catch rather than biological signals.
- The tagging data q was fixed and strongly dictated scaling in the assessment. The tagging study data and subsequent constraint in q values needs further consideration.
- Continued ageing and age validation of black rockfish samples
- A fishery-independent nearshore survey was recommended

Most of the above recommendations were included in the 2015 assessment research recommendations. Additionally, stock structure for black rockfish was highlighted as a topic for further consideration.

3.2 Model description

3.2.1 Modelling Platform

Stock Synthesis version 3.30.21 was used as the statistical catch-at-age modelling framework. This framework allows the integration of a variety of data types and model specifications. The SS-DL tool (<https://github.com/shcaba/SS-DL-tool>) was used for model exploration, likelihood profiling, and sensitivity analyses. The companion R package r4ss (version 1.38.0) along with R version 4.2.2 were used to investigate and plot model fits.

3.2.2 Bridging the assessment model from Stock Synthesis 3.24 to 3.30

Since several years have passed from the last assessment model, the Stock Synthesis (SS) modelling framework has undergone many changes. While the specific changes in the model can be found in the model change log, here we simply update the model from the older 3.24V version to the newer 3.30.20 version. The point here is to present any differences in the model outputs when using the same information. This was first done by migrating the

data and parameter specifications from the former files to the newer files. This migration was assisted using the SS-DL tool. Once the old data was transferred to the SS 3.30.20 file, two versions of the model were run.

- 1) Fixing all parameter values to the values found in the 2015 model.
- 2) Allowing the same parameters estimation specification as in the 2015 model

Results are similar between models when all parameters are fixed from the 2015 model in the updated SS files, while there are slight scale differences (Figure 18), but little relative stock status differences (Figure 19), when the new SS version is allowed to estimate the same parameters as estimated in the 2015 version. These model comparisons are adequate to move ahead using the newest version of SS 3.30.20 without expecting large differences in reference models being due to versions of SS.

3.3 Model Structure and Assumptions

3.4 Model Structure, Evaluation, and Specification

3.4.1 Fleet and survey designations

The Oregon model is structured to track several fleets and include data from several surveys:

- Fleet 1: Commercial trawlfishery
- Fleet 2: Commercial non-trawl fishery
- Fleet 3: Recreational ocean boat fishery
- Fleet 4: Recreational shore fishery
- Survey 1: Commercial non-trawl CPUE

- Survey 2: ORBS CPUE survey
- Survey 3: MRFSS CPUE survey
- Survey 4: MPA CPUE survey
- Survey 5: Acoustic visual survey
- Survey 6: Tagging abundance survey

The specifications of the assessment are listed in Table 17.

3.5 Model likelihood components

There are five primary likelihood components for each assessment model:

1. Fit to survey indices of abundance.
2. Fit to length composition samples.
3. Fit to age composition samples (all fit as conditional age-at-length).
4. Penalties on recruitment deviations (specified differently for each model).
5. Prior distribution penalties

3.6 Reference model exploration, key assumptions and specification

The base assessment model for Oregon Black Rockfish was developed to balance parsimony and realism, and the goal was to estimate a spawning output trajectory and relative stock status for the population of Black Rockfish in state and federal waters off Oregon. The model contains many assumptions to achieve parsimony and uses different data types and sources to estimate reality. A series of investigative model runs were done to achieve the final base model. Constructing integrated models (i.e., those fitting many data types) takes considerable model exploration using different configurations of the following treatments:

- Data types
- Parameter treatments: which parameter can, cannot and do not need to be estimated
- Phasing of parameter estimation
- Data weighting
- Exploration of local vs global minima (see model convergence section below)

The different biological data with and without the catch time series were first included to obtain an understanding of the signal of stock status coming from the data (Figure 20). Both the lengths and ages, with and without the removal time series, demonstrated a signal for a highly depleted stock. Stock scale was comparable once removal history was included, and showed lower estimates of scale. This departure from the ultimate reference model highlights the major uncertainty in the treatment of the biological data in when characterizing model uncertainty. Further sensitivities to the inclusion or exclusion of different indices of abundance are provided in the model sensitivity section.

Numerous exploratory models that included all data types and a variety of model specifications were subsequently explored and too numerous to fully report. In summary, the estimation of which life history parameters to estimate and fix was liberally explored. The following is a list of things that were explored, typically in combination with one another

- Estimate or fix M
- Estimate or fix any of the three growth parameter for each sex
- Estimate or fix the stock-recruit relationship
- Estimate or assume constant recruitment. If estimating recruitment, for what years?
- Estimate or fix survey catchability for each survey
- Estimate additional survey variance for which survey
- Estimate or fix selectivity parameters
- Logistic or dome-shaped selectivity

After much consideration, it was determined that some parameters were inestimable (M, L_{min} for both sexes), some did not move much for initial values and could be fixed (e.g., CV at length values, some selectivity parameters), and others could be estimated (e.g., $L_\infty, k, \ln R_0$). Estimation of L_{min} or fixing it to the externally estimated values consistently gave very high estimates of L_∞ for both sexes, thus the L_{min} value for both females and males was fixed so $t_0 = 0$, as this gave the most realistic L_∞ values. No priors were used on any of the estimated parameters except female L_∞ which used a normal prior and a standard deviation taken from the external fit to the growth curve (0.17). Length at maturity, fecundity-weight, and length-weight relationship, steepness (h) and recruitment variance were all fixed.

Examination of the catches in the trawl fishery indicated much larger fish being caught versus those caught in the commercial non-trawl and recreational ocean boat fisheries (both of which have very similar mean lengths and aggregate catch compositions) in the same years. Dome-shaped selectivity was therefore allowed in those fisheries as well as the shore-based fleet (which catches much smaller individuals) and the MPA survey. The ORBS and MRFSS surveys were assumed to have the same selectivity as the recreational boat fishery fleet. While constant selectivity was assumed across all fleets, it was also noted that the earlier time series (prior to 2003) of the recreational boat fleet, as sampled by the MRFSS program, had consistently smaller mean lengths than the ORBS sampled years. The MRFSS years also had smaller samples with more uncertainty. Time varying selectivity was explored in the two periods to see if this improved the fit to the data, but no perceivable improvement was made and the resultant difference was very small in the derived model outputs, so a constant selectivity was assumed for the whole time period.

The full list of estimate and fixed parameters are found in Table 18.

The biggest uncertainty revealed in the model exploration was conflict in the signal of the biological data and that in the acoustic visual survey catchability. If the inherent uncertainty in the survey ($CV = 0.45$) was used, additional variance estimated on a lower CV, or the catchability was estimated, the acoustic visual survey was essentially ignored (catchability was between 3 and 4, saying the survey overestimates abundance by 3 to 4 times), the scale and stock status dropped significantly. If the acoustic survey catchability, reviewed and deemed informative of overall black rockfish biomass in Oregon (Rasmussen et al. 2022), was assumed to be equal to 1, the biomass and stock status was much higher. Given there are serious issues with assuming either of the previous hypotheses are acceptable, the STAT

determined a realistic upper bound (based on a calculated catchability from the uncertainty in the acoustic visual survey) for catchability of 1.822 (or $\ln(q)$ of 0.6). This served as fixed value of catchability (which also required a small CV of 0.0001 on that index) to build the final reference model around.

General attributes of the reference model are that indices of abundance are assumed to have lognormal measurement errors. Length compositions and conditional age at length samples are all assumed to follow a multinomial sampling distribution, where the sample size is fixed at the input sample size calculated during compositional example, and where this input sample size is subsequently reweighted to account for additional sources of overdispersion (see below). Recruitment deviations were also estimated are assumed to follow a lognormal distribution, where the standard deviation of this distribution is tuned as explained below.

Sensitivity scenarios and likelihood profiles (on the catchability of the acoustic visual survey, steepness, and natural mortality) were used to explore uncertainty in the above model specifications and are reported below.

3.6.1 Data Weighting

The reference model allowed for the estimation of additional variance on all surveys except the acoustic visual survey. This additional variance allows the model to balance model fit to that data while acknowledging that variances may be underestimated in the index standardization. A sensitivity was run with no extra variance estimated, as well as removal of the index data were explored.

Initial sample sizes for the commercial and recreational length and conditional age-at-length compositions were based on the number of input effective samples sizes. The method of Francis (2011, equation TA1.8) was then used to balance the length and conditional age-at-length composition data among other inputs and likelihood components. The Francis method treats mean length and age as indices, with effective sample size defining the variance around the mean. If the variability around the mean does not encompass model predictions, the data should be down-weighted until predictions fit within the intervals. This method accounts for correlation in the data (i.e., the multinomial distribution), but can be sensitive to years that are outliers, as the amount of down-weighting is applied to all years within a data source, and are not year-specific. Sensitivities were performed examining different data-weighting treatments: 1) the Dirichlet-Multinomial approach (Thorson et al. 2017), 2) the McAllister-Ianelli Harmonic Mean approach (McAllister and Ianelli 1997), or 3) no data-weighting of lengths.

3.6.2 Model Changes from the Last Assessment

Besides the addition of eight years of data, changes to some historical data series, and some changes in the estimation of some parameters, the biggest changes to the past assessment are:

- Addition of and fixing to the acoustic visual survey. This is different than the last model that fixed the stock scale to the tag survey catchability
- Estimation of recruitment
- Female natural mortality is no longer a step, but a constant value
- Length-based selectivity only (i.e., no age-based selectivity for females), and dome-shaped selectivity for the ocean boat fisheries (assumed logistic last time)
- Change in the removal history

3.7 Reference Model Diagnostics and Results

3.7.1 Model Convergence and Acceptability

While there is no definitive measure of model convergence, several measures are routinely applied. These criteria include a low maximum gradient (4.69204×10^{-4}), inversion of the Hessian (passed), acceptable fits to data (passed), and reasonable parameter values (passed).

Model efficiency was explored by doing a short run Bayesian analysis using the Random Walk Metropolis with 2000 draws, keeping all the draws and examining the fast mixing parameters. Those estimated parameters that do not move much from the initial values slow the model down and are recommended to be fixed at the starting value (Monnahan et al. 2019). This resulted in the fixing of 3 selectivity parameters (Figure 23).

An extra effort was given to ensure the model did not rest on a local likelihood minimum. This was done by starting the minimization process from dispersed parameter values away from the maximum likelihood estimates to determine if the approach found a better model fit (i.e., minimum negative log-likelihood value). Starting parameters used a jitter shift value of 0.01. This was repeated 100 times with 17 out of 100 runs returned to the reference model likelihood (Figure 22). A better fit, lower negative log-likelihood model was not found in any of remaining runs. The model did not experience convergence issues when provided reasonable starting values. Through the jittering and likelihood profiles, the present reference model represents the best fit to the data given the assumptions made.

3.8 Base Model Results

3.8.0.1 Fits to the Data

3.8.0.1.1 Lengths Fits to the length data are examined based on the Pearson residuals-at-length, the annual mean lengths, and aggregated length composition data for the commercial and recreational fleets. Fits to the annual length composition are shown in Appendix A.

Pearson residuals of fits to the fishery and MPA survey length data are reasonably small with no distinct patterns (Figure 24), though the shore-based fishery lengths, which are small samples, show a possible recruitment in 1985 that is not seen in the recruitment residuals.

Model fits to the mean lengths, assuming Francis data-weighting, do not follow the dynamics in means lengths in the relatively small trawl fishery (Figure 25). The mean lengths are better fit in the larger and better sampled non-trawl (Figure 26) and ocean boat (Figure 27) fisheries. Note the poor fit to the MRFSS-era (pre-2003) mean lengths in the ocean boat fishery. Those length are consistently fit higher than the mean, but the samples are much lower than in the ORBS era sampling. A time block in selectivity was attempted when exploring potential model specifications, but the fits were not improved. The shore-based, also poorly sampled, are not fit well and miss what could be a potential recruitment in the mid-1980s (Figure 28). Fits to the mean length in the MPA length data are generally good (Figure 29). All length weighting are near the value of 1 as is expected when using the Francis weighting method. One thing to note is the general lack of contrast in the data over the past two decades.

Aggregate fits over year by fleet are shown in Figure 30. The two main fisheries (non-trawl and ocean) and the MPA survey show good fits to the aggregated data. The trawl fishery, which requires the asymptotic size of the populations to be much higher than is expected given the age and growth data, is fit less well. But the trawl data does demonstrate that mostly large individuals are present in the deeper trawl fishery, suggesting less availability of those sizes in the shallower non-trawl and recreational fisheries. This is instructive to the fact that the trawl fishery is catching only the largest individuals in the population, thus possibly over representing those length classes in the population, and indicating that the non-trawl and ocean fishery is missing some of those bigger fish (i.e., a dome-shaped selectivity). The shore-based fishery is also being fit poorly, but the fishery is so small it makes little difference to model performance.

3.8.0.1.2 Conditional Age at Length Fits to the mostly sex-specific conditional age at length data are examined based on the age-at-length Pearson residuals, the annual mean ages, and mean age at length by year for the trawl, non-trawl and ocean boat fishery samples. The maximum size of the Pearson residuals for each fleets was fairly small (maximum = 13.44, 12.48, and 13.12 for the trawl, non-trawl and ocean boat samples, respectively; Appendix B). The biggest residuals were limited, as most of the residuals were small and unnoteworthy and demonstrate the expected shape of the growth curve. There is more contrast in the age data, as the mean age by year increased for all fleets through the most recent years (Figures 31, 32, and 33). The mean age for the trawl fishery was between 10-15 years, whereas the mean ages for the non-trawl and ocean boat fishery was between 8 and

10 years. Fits to the mean ages by length bins show acceptable fits consistent with model expectations Appendix C.

3.8.0.1.3 Marginal Age compositions Marginal age compositions are not fit in the model, but they are included in order to see how well they fit the reference model without influencing the likelihood Appendix D. Marginal length and age composition cannot be used in the same model given the overlap of the same fish in both samples. This is why ages conditioned on lengths are often used with the length compositions. But it still stands that age compositions, instead of lengths, could be used. So adding the marginal age compositions passively (i.e., not contributing to the overall likelihood of the model) can offer insight into how consistent they are with the current model fit. Overall the fits are very good, and in an exploratory model run the length compositions were replaced with the age compositions and a very similar model result was obtained. Thus the conditional age at length compositions coupled with either length or age compositions provide similar results. This is likely due to the fact that the conditional ages capture the underlying age structure sufficiently to follow the age structure over time.

3.8.0.1.4 Fits to Indices of Abundance The fits to the five available indices of abundance (not including the acoustic-visual survey which the model was predefined to fit with a $\ln q=0.6$) are generally weak. The best fit index is found for the commercial non-trawl fishery (Figure 34), with a very good fit of an upward trend in the contemporary time series. In an interesting contrast is the ocean boast survey that covers the same years and shows a very similar selectivity, but shows an opposing downward trend (Figure 34). The earlier time series for the recreational fishery is also not well fit to the dynamics of the model, and requires high additional variance to fit (Figure 35).

The two fishery independent surveys, the MPA (Figure 38) and the tagging survey (Figure 37) are also not well fit. Despite this, the estimated catchability for the tagging survey is 0.3, not far from the assumed value of 0.25 in the 2015 assessment.

3.8.1 Reference Model Outputs

3.8.1.1 Parameter Estimates and Derived Outputs A total of five population parameters, five survey variances and sixteen selectivity parameters were estimated, along with thirty-eight recruitment deviations. The reference model parameter estimates along with asymptotic standard errors are shown in Table 18 and the likelihood components are shown in Table 19. Estimates of derived outputs and reference points and approximate 95 percent asymptotic confidence intervals are provided in Table v.

The estimates of sex-specific growth parameters showed some differences from the externally estimated starting values (Table 18 and Figure 39). The estimated L_∞ and k for females

and males were slightly greater than the values estimated externally. The majority of female and male Black Rockfish growth occurs at younger ages, reaching near maximum length by age 20-25, depending upon sex, with female Black Rockfish reaching larger maximum lengths (Figure 39).

The time series of estimated recruitment deviations and annual recruitment deviations are shown in Figures 40 and 41. Years with major recruitment deviations were estimated to have occurred around 1994, 1999-2000, 2008, and 2016. These years are very similar to recruitment deviations seen in other rockfishes. One curious signal is the string of positive deviations from 2010-2014, though this is consistently apparent in most model runs. The variance check on the recruitment deviations indicates well informed recruitments from 1985 to 2017, providing justification for the estimation of recruitment (Figure 45). Recruitment deviations after 2017 are relatively uninformed and are not estimated. The recruitment bias adjustment applied within the model across years is shown in Figure 44. The Beverton-Holt stock-recruitment relationship compared to the recruitment deviations are given in Figure 43.

Estimated selectivity curves for each fleet and survey (Figure 46) look plausible (i.e., as a model convergence check for realism, the selectivity curves must look plausible). The trawl fishery is highly right shifted indicating only the largest individuals are taken by that fishery. Both the non-trawl and the recreational ocean-based fleet (the MRFSS fleet is mirrored to this fleet) demonstrate dome-shaped selectivity and are very similar though the recreational fishery is expected to take more larger individuals. The estimate of the peak size of selectivity for the two biggest removal sources (non-trawl = 42.63 cm; recreational ocean boat = 41.84 cm) are above the size of 50% functional maturity (40 cm).

3.8.1.2 Population Trajectory The predicted spawning output (in millions of eggs) is provided in Table 22 and plotted in Figure 47. Estimated spawning output shows a large decline starting in the late 1970s when the recreational fishery begins in earnest. This decline is consistent until the late 1990s when it then rebounds due to several strong recruitments consistent with other rockfish stock assessments. The estimate of total biomass over time, which tracks that of spawning output, is shown in Figure 48.

Relative spawning output declined below the management target ($SB_{40\%}$) by the 1990s, but rebounded to above the target in the 2010s (Figure 49). The relative stock status at the start of 2023 (0.53) is estimated to be above the rockfish relative biomass target of 0.4. Uncertainty intervals are extremely small and unbelievable, thus emphasizing alternative model specifications (mainly in the treatment of catchability) to capture uncertainty. The strong recruitment events are responsible for the dramatic increase and elevated stock status. Numbers of age-0 individuals indicate those years of particularly strong recruitment (Figure 40).

This current estimate shows a higher overall biomass (Figure 50), but similar relative stock status, as the 2015 stock assessment (Figure 51).

3.9 Characterizing uncertainty

3.9.1 Sensitivity Analyses

Sensitivity analyses were conducted to evaluate model sensitivity to alternative data treatment and model specifications.

3.9.1.1 Data treatment sensitivities Data treatments explored included the treatment of catchability for the acoustic-visual survey and the tagging data, the removal of any one data source, and alternative approaches to data weighting. All scenarios match the reference model specifications in all other aspects unless otherwise stated.

- Treatment of absolute abundance catchability (q)
 1. Acoustic visual survey $q = 1$, CV = 0.0001
 2. Acoustic visual survey q Estimated, CV = 0.0001
 3. Acoustic visual survey $q = 1$, CV = 0.45
 4. Acoustic visual survey q estimated, CV = 0.45
 5. Tag survey q = to 2015 value (0.25) with #2 above
 6. Tag survey q = to 2015 value (0.25), but extra variance estimated, with #2 above
- Remove abundance index series
 7. No Acoustic visual
 8. No tagging index
 9. No non-trawl index
 10. No ORBS index
 11. No MRFSS index
 12. No MPA index
 13. Add SMURFs index
- Data weighting
 14. No data-weighting
 15. Dirichlet data-weighting
 16. McAllister-Ianelli data weighting

Likelihood values and estimates of key parameters and derived quantities from each data treatment sensitivity are available in Tables 23 and 24. Derived quantities relative to the reference model are provided in Figure 52. Time series of spawning output and relative spawning output are shown in Figures 53 and 54.

From a data standpoint, the model was the most sensitive to how information on absolute abundance, or population scale, from the acoustic-visual survey and the tagging data was

treated in the model. In particular, the parameterization of catchability (model estimated or fixed) and the level of error (input CVs and model estimated extra variance) associated with these two data sources provided considerably discrepancies in key management quantities (e.g., stock size and status; Figures 52, 53, and 54). The model was comparatively robust to the removal of any given single data source, with the exception of the acoustic-visual survey, and the relative weighting of length and age composition data. The harmonic mean data weighting approach (based on the McAllister-Ianelli method) resulted in negligible change from the reference model that used the Francis weighting method. The Dirichlet method only minimally adjusted the input sample sizes, nearly mimicking the sensitivity model where no weighting was conducted, which had the effect of increasing current estimates of stock size (20%) but no appreciably change in stock status.

Alternative treatment of the acoustic-visual survey data largely resulted in two alternative states of nature in terms of stock size and status (high and low) relative to the base model. Fixing catchability to 1.0 and reducing the input CV to near zero essentially forced the model to fit the 2021 acoustic-survey data point perfectly (high state of nature). However, when doing so the model estimated biological parameters (e.g., natural mortality and growth) that were extreme, nearing implausibility. Alternatively, estimating catchability, or similarly inputting the externally estimated CV of 0.45, resulted in a low state of nature where estimates of spawning output over recent years were below OFLs and stock status was in the precautionary zone. However, the interpretation of resulting acoustic-survey catchability was impractically high (3.0 or higher) given the comprehensive study design of the survey. Fixing the tagging catchability to the value assumed in the last assessment largely counteracts the change to the reference model after fixing the acoustic-visual catchability to 1.0 and reducing its input CV to near zero.

3.9.1.2 Model specification sensitivities Model specifications looked at the estimation of individual and combinations of life history parameters, including the treatment of natural mortality, growth, maturity, fecundity and recruitment, and fishery selectivity. All scenarios match the reference model specifications in all other aspects unless otherwise stated.

- Life history estimation
 - Natural mortality (M)
 1. Use 2015 M values
 2. Lorenzen age varying M
 3. Estimate M
 - Growth parameters
 4. Fix all growth parameters to external values, estimate M
 5. Fix all growth parameters to external values
 6. Estimate $L_{m\text{in}}$
 7. Estimate CV_{young} and CV_{old}

- Reproductive Biology
 - 8. Use biological maturity ogive (logistic model)
 - 9. Use functional maturity ogive (logistic model)
 - 10. Fecundity proportional to weight
- Recruitment estimation
 - 11. No recruitment estimation
 - 12. Estimate recruitment for all years in the model
- Selectivity 13. Estimate logistic selectivity for all fleets

Likelihood values and estimates of key parameters and derived quantities from each model specification sensitivity are available in Table 25. Derived quantities relative to the reference model are provided in Figure 55. Time series of spawning output and relative spawning output are shown in Figures 56 and 57. In general, model specifications for natural mortality, selectivity, and growth caused the population to drop below the reference model estimate of stock size, while alternative reproductive biology models increased stock size. Despite considerable differences in the historic trajectory of relative stock status across sensitivity models, there was considerably less variability in the estimate of current (2023) stock status (ranging from 0.49 to 0.65, where 0.55 is the reference model, with the exception of the estimate L_{min} growth model that had convergence issues when estimating the length at minimum age).

Estimating female and male M led to a considerably higher mortality rate for females (0.25 compared to 0.19 used in the reference model) and slightly higher for males (0.18 compared to 0.17). This had the effect of lowering both the beginning and ending stock scale, but had only a marginal influence on current stock status. Similarly, fixing growth but estimating M also dropped stock scale but preserved stock status relative to the sensitivity that only fixed the growth parameters to external estimates. Due to the similarity between M in the reference model and what was used in the previous 2015 assessment, there was little change in overall management quantities associated with this sensitivity model.

Fixing growth parameters in the model to external-to-the-assessment estimates resulted in a considerable change in the stock trajectory, both in terms of stock size and status, yet end up in about the same place by 2023. For example, over the most recent 20 years, this sensitivity model estimates the stock declining from a nearly unfished state in the early 2000s. In contrast, the reference model (and nearly all other sensitivity model) estimate the stock has been increasing from a precautionary state in the early 2000s. Estimating the CV for growth of young and old fish had negligible impact on results. The model that estimated L_{min} did not pass key model diagnostic tests (e.g., stability and convergence) so was not considered further.

Using a logistic function to externally estimate maturity ogives, based on either biological or functional maturity measurements, increased scale and overall spawning output relative to the reference model. This result was most prominent when biological maturity was used to

describe the maturity schedule due to the appreciable reduction in the length at 50 percent maturity indicating that females mature at a younger age on average than when using functional maturity (Figure 12). The minor differences between using a logistic function (sensitivity) versus a spline (reference model) was due to the latter allowing for skip spawning of older, mature females. Assuming that fecundity is proportional to female body weight, instead of an exponential relationship with length as used in the reference model, resulted in an increase in beginning and ending scale but little different in overall stock status.

Recruitment estimation (assuming $\sigma_R = 0.6$) was sensitive to the estimation of different ranges of years of recruitment deviations. While the truncated time series of recruitment used in the reference model can be considered parsimonious, the addition of estimating all recruitment years adds additional uncertainty to the stock status measure earlier in the time series. Estimating recruitment deviations for all years in the model dropped the scale of the population, altered the historic stock status trajectory, but had little effect on current stock status. Estimating the full time series of recruitments resulted in a significant negative trend in deviates early in the time period that was a concern. Removing the estimation of any recruitment deviates from the stock-recruitment curve resulted in reduced stock dynamics, where the decline in stock status from an unfished state was gradual and largely driven by the catch history.

Estimating all fleet selectivity patterns as logistic effectively removed the potential for cryptic biomass resulting from fleets not encountering and catching fish of older sizes or ages. This model resulted in a decrease in beginning and ending scale and reduced stock status since 1990 (even to the point where the stock would have been considered overfished from the 1990s to the early 2010s). The stock status discrepancy is reduced the closer the model gets to the current year, where the fraction of unfished is estimated to be 53 percent in 2023 compared to the reference model of 55 percent. The fit to the logistic selectivity model is not as good in terms of residual patterns and overall likelihood (based on AIC).

3.9.2 Likelihood Profiles

Likelihood profiles were conducted for the acoustic-visual survey catchability (q) in natural log-space ($\ln(q)$), steepness (h), and female and male natural mortality (M) varying together with a constant ratio matching the reference model (0.19/0.17, female and male values respectively). Likelihood profiles were conducted by fixing the featured parameter(s) at specific values across a range of values and estimating all remaining parameters. A likelihood profile offers insight into model sensitivity to changing model parameter values, while providing an additional way to describe uncertainty in the parameter by identifying the range of parameters within 1.96 likelihood units of the reference model.

The profile on the assumption of the acoustic-visual survey catchability demonstrates the expected scaling up and down of the population with the decrease and increase of the catchability, respectively (Figure 58). The change in the current spawning output is steeper

than the initial spawning output, leading to a steep decline in relative biomass as $\ln(q)$ gets larger. As the $\ln(q)$ approaches 1.5, the stock approaches the overfished limit reference point. The most supported value of the $\ln(q)$ is found near 1.17 with a very tight support interval. The most prominent length data (in samples and associated fisheries) support larger $\ln(q)$ values, while age compositions support smaller $\ln(q)$ values near the value used in the reference model (Figure 59). The less informed survey indices support larger $\ln(q)$ values on the strength of the best fit index (non-trawl CPUE index).

The steepness profile showed little information content for this parameter (Figure 60) as the best fit value went towards the bound of 1. Changing steepness mostly changed the estimate of initial stock size, with a higher initial stock size needed to make up for the lower recruitment compensation. Overall, the relative stock size changed little across a large range of steepness values. The biological compositions pushed the steepness to the highest value, whereas the index data indicated a mixed signal for the lowest and highest steepness values (Figure 61).

The combined profile that varies female and male M while maintaining the reference model offset demonstrated what had been seen in the model development, that when freely estimated, the model wants natural mortality values to be unreasonably high (Figure 62). Given the highly constrained model specification (i.e., a fix q on a survey), the range of reasonable natural mortality values adjust the scale of the population in order to match the value fixed in 2021. Similar to the steepness profile, the lower natural mortality values lead to lower productivity, and thus the need for a higher initial biomass, which changes quicker than the current, more pinned down, current biomass value. While the scale does change substantially, the current relative stock status is fairly stable and always above the target level.

3.9.3 Retrospective Analysis

A ten-year retrospective analysis was conducted by running the model and sequentially removing one year of data up through minus 10 years. Retrospective spawning output (Figure 63) and relatives stock status (Figure 64) estimates show a large shift away from a growing population once the acoustic-visual survey is removed from the population. The scale shifts downward and the population status show a declining population at or below the unfished level. This shows the weight of the treatment choice in the acoustic survey, as well as the more recent data that confirm a recent recruitment. The Mohn's rho evaluation of the degree of retrospective pattern is given in Table 26.

3.9.4 Unresolved Problems and Major Uncertainties

The biggest uncertainty and unresolved conflict is trying to reconcile the signal in the biological data (which want a lower population size and status) versus the acoustic and

tag surveys (which want high stock sizes and status). This is the major issue the current assessment is confronting. Another acoustic-visual survey data point could help resolve how much uncertainty there is in the estimate. The lack of contrast in the biological data, despite large sample sizes, is another barrier to interpreting the current conditions, though given models using only biological data, the signal seems clear that the population could be at a lower stock status.

4 Management

4.1 Reference Points

Reference points were based on the rockfish FMSY proxy ($SPR_{50\%}$), target relative biomass (40%), and estimated selectivity and catch for each fleet (Table 20). The Black Rockfish population in Oregon at the start of 2023 is estimated to be 1.38 times (above) the target biomass, and fishing intensity during 2022 is estimated to be 0.96 times (below) the fishing intensity target (Figure 66). The yield values are lower than the previous assessment for similar reference points due to updated life history estimates. The proxy MSY values of management quantities are the most conservative compared to the estimated MSY and MSY relative to 40% biomass. Sustainable total yield, removals, using the proxy $SPR_{50\%}$ is 455 mt. The spawning output equivalent to 40 percent of the unfished spawning output ($SO_{40\%}$) calculated using the SPR target ($SPR_{50\%}$) was 728.5 meggs.

Recent removals have been close to the point estimate of potential long-term yields calculated using an $SPR_{50\%}$ reference point, though the population size has continued to increase over recent years due to several above average recruitments. The equilibrium estimates of yield relative to biomass based on a steepness value fixed at 0.72 are provided in Figure 67, where vertical dashed lines indicate the estimate of fraction unfished at the start of 2023 (current) and the estimated management targets calculated based on the relative target biomass (B target), the SPR target, and the maximum sustainable yield (MSY).

The 2023 spawning biomass relative to unfished equilibrium spawning biomass, based on the 2022 fishing year, is above (55 percent) the management target of 40 percent of unfished spawning output. The relative biomass and the ratio of the estimated SPR to the management target ($SPR_{50\%}$) across all model years are shown in Figure 66 where warmer colors (red) represent early years and colder colors (blue) represent recent years. There have been periods where the stock status has decreased below the target and fishing intensity has been higher than the target fishing intensity based on $SPR_{50\%}$.

4.2 Management performance

Black Rockfish removals have generally been below the equivalent ABC-ACL over the recent decade, with the exception of 2017 and 2022 when removals were slightly higher (Table 21). Exploitation on Black Rockfish increased starting around 1940 and reached a high in the late 1970s. Since that time, catch has mostly fluctuated between 300 and 500 mt per year, with some years exceeding 600 mt. Removals have averaged 483 mt over the past decade. The last ten years of Black Rockfish acceptable biological catch (ABC) and annual catch limit (ACL) (which are equivalent) has been set, by definition, below the overfishing limit (OFL) (Table 21). Prior to 2017, management specifications were set for Black Rockfish in California and Oregon waters combined. The Black Rockfish OFL has not been exceeded by Oregon removals over the past decade.

4.3 Unresolved Problems and Major Uncertainties

The biggest uncertainty and unresolved conflict is trying to reconcile the signal in the biological data (which suggests a lower population size and status) versus the acoustic and tag surveys (which suggest high stock sizes and status). This is the major issue the current assessment is confronting. Another acoustic-visual survey data point could help resolve how much uncertainty there is in the estimate. The lack of contrast in the biological data, despite large sample sizes, is another barrier to interpreting the current conditions, though given models using only biological data, the signal seems clear that the population could be at a lower stock status.

4.4 Harvest Projections and Decision Tables

The following text will be modified, as appropriate, after the STAR panel and SSC meeting.

The Black Rockfish assessment is being considered as a category 2 assessment with a P^* = 0.45, sigma = 0.72, and a time-varying buffer applied to the OFL. These multipliers are also combined with the rockfish MSY proxy of FSPR=50% MSY and the 40-10 harvest control rule to calculate OFLs, ABCs and ACLs. A twelve year (2023-2034) projection of the reference model using these specifications along with input removals for 2023 and 2024 provided by the Groundfish Management Team is provided in Table 27.

Uncertainty in management quantities for the reference model was characterized by exploring various model specifications in a decision table. Initial explorations considered alternative specifications of catchability for the acoustic-visual survey. The resultant decision table will be provided in Table 28.

Further details about selecting the decision table states of nature will be added here after the STAR panel.

Further details summarizing the results presented in the decision table will be added here after the STAR panel.

4.5 Evaluation of Scientific Uncertainty

The model-estimated uncertainty around the 2023 spawning biomass was $\sigma = 0.03$ and the uncertainty around the OFL was $\sigma = 0.01$. This is clearly an underestimate of overall uncertainty because of the necessity to fix some life history parameters such as natural mortality and steepness, as well as a lack of explicit incorporation of model structural uncertainty. The alternative states of nature used to bracket uncertainty in the decision table assist with encapsulating model structure uncertainty.

4.6 Research and Data Needs

This section briefly highlights progress on research and data needs identified in the most recent (2015) Black Rockfish assessment, and then provides recommendations for future research.

Research and data needs identified in the last assessment are listed here followed by a brief response.

Further investigation into the movement and behavior of older (> age 10) females to reconcile their absence in fisheries data. If the females are currently inaccessible to fishing gear, can we find where they are? This information is essential before another black rockfish assessment is undertaken. Response: Work by ODFW, Rasmuson et al. (2023), specifically looked at finding older female Black Rockfish. While this work was a step in the right direction, more hypothesis-drive work is needed to reconcile the fate of older females (thus this recommendation remains moving forward, see item 1 below).

\textit{Appropriate natural mortality values for females and males. This will help resolve the extent to which dome-shaped age-based selectivity may be occurring for each. This is a larger question on how to empirically estimate \$M% when direct measures are not available (which is usually the case).} Response: While natural mortality remains an unknown, this assessment does explore the estimation of M using informed priors based on the Hamel and Cope (2022) method. Research related to male and female M remains a topic for further research (see item 3 below).

All states needed improved historical catch reconstructions. The trawl fishery catches in particular need particular attention. Given the huge historical removals of that fleet in each

state, the assessment is very sensitive to the assumed functional form of selectivity. A synoptic catch reconstruction is recommended, where states work together to resolve cross-state catch issues as well as standardize the approach to catch recommendations. Response: While work has been conducted to improve historical catch reconstructions for various fleets (much of which went into the updated catch time-series used in this assessment), there still is more work to do (see item 4 below).

Identifying stanzas or periods of uncertainty in the historical catch series will aid in the exploration of catch uncertainty in future assessment sensitivity runs. Response: This has largely been addressed with recent historic catch reconstruction updates.

The ODFW tagging study off Newport should be continued and expanded to other areas. To provide better prior information on the spatial distribution of the black rockfish stock, further work should be conducted to map the extent of black rockfish habitat and the densities of black rockfish residing there. Response: The coastwide Black Rockfish acoustic-visual survey conducted in 2021 used contemporary habitat maps during study design development and has a goal of estimating coastwide abundance as well as habitat-specific densities.

Interpreting the ODFW tagging study needs further work. The Brownie model is a closed-model estimate of abundance, an assumption that could have meaningful influence on the interpretation of the survey catchability. Given that catchability parameter is currently driving the Oregon, realistic values for it need further consideration. Response: The coastwide acoustic-visual survey was used to pin the absolute estimate of abundance in 2021, thereby providing an anchor for the tagging data catchability.

Age validation and verification is still needed, particularly in Oregon. A number of historical ages were excluded from the Oregon model due to concerns over differences among age readers. These age structures need to be re-read to recover the information in those samples. Historical structures from trawl and recreational fisheries in particular should be re-aged by reliable readers and included in future assessments. Oregon ageing error was also quite different from the other two states, so further investigation on the inter-reader error and how to decrease it need reconsideration. Response: Considerable effort went into not only reading new Black Rockfish ages but also double reading historical age structures (see the section on Ageing Bias and Precision for specific details).

An independent nearshore survey should be supported in all states to avoid the reliance on fishery-based CPUE indices. Response: A coastwide acoustic-visual survey was conducted in Oregon waters during 2021.

Stock structure for black rockfish is a complicated topic that needs further analysis. How this is determined (e.g., exploitation history, genetics, life history variability, biogeography, dispersal and movement, etc.) and what this means for management units needs to be further refined. This is a general issue for all nearshore stocks that likely have significant and small scale stock structure among and within states, but limited data collections to support small-scale

management. Response: This recommendation still stands and is included in future research recommendations (see item 5 below).

Recommended avenues for research to help improve future Black Rockfish stock assessments include (but are not limited to):

1. Continue work on the investigation into the movement, and behavior or mortality of older (> age 10) females to further reconcile their absence in fisheries data.
2. Conduct population genetics studies on fish observed off of the continental shelf (middle of the gyre and at sea mounts) to determine their association with the nearshore stocks.
3. Continue to build evidence for appropriate natural mortality values for females and males.
4. Improved historical catch reconstructions. Specifically, the historic trawl fishery catches (pre-1987) in particular require particular attention. Given the huge historical removals of that fleet in each state, the assessment is very sensitive to the assumed functional form of selectivity. A synoptic catch reconstruction is recommended, where states work together to resolve cross-boundary state catch issues as well as standardize the approach to catch recommendations to the extent possible.
5. Stock structure for black rockfish is a complicated topic that needs further analysis. How this is determined (e.g., exploitation history, genetics, life history variability, biogeography, etc.) and what this means for management units needs to be further refined. This is a general issue for all nearshore stocks that likely have significant and small scale stock structure among and within states, but limited data collections to support small-scale management.
6. Continue acoustic-visual fisheries independent coastwide survey to develop a time series. Further refine the survey by addressing the recommendations of the SSC methodology review from 2022. Examine the potential of using spatial modeling to reduce the uncertainty in the population estimates from the acoustic-visual fisheries independent coastwide survey.
7. Reconcile contradictory signal in the Black Rockfish biology versus the population scale.
8. Better understand the ecology and habitats of Black Rockfish from settlement to age 1. Further development of surveys aimed specifically at recruitment or settlement rates of nearshore species, such as OSU's SMURF collections, that are not frequently encountered in offshore federal age-0 surveys is needed.

5 Acknowledgments

Here are all the mad props! We thank...

6 References

- Ayres, D.L. 1988. Black rockfish investigations: A summary of 1986 and 1987 black rockfish tagging studies. Washington Department of Fish; Wildlife.
- Baker, B.M. 1999. Genetic analysis of eight black rockfish collections from northern oregon. Washington Department of Fish; Wildlife.
- Berkeley, S.A., Chaoman, C., and Sogard, S.M. 2004. Maternal age as a determinant of larval growth and survival in a marine fish, *sebastes melanops*. Ecology **85**: 1258--1264.
- Berkeley, S.A., Chapman, C., and Sogard, S.M. 2008. Maternal age as a determinant of larval growth and survival in a marine fish, *sebastes melanops*.
- Beverton, R.J.H., and Holt, S.J. 1957. On the dynamics of exploited fish populations. Ministry of Agriculture, Fisheries; Food, Berkeley.
- Bobko, S.J., and Berkeley, S.A. 2004a. Maturity, ovarian, cycle, fecundity, and age-specific parturition of black rockfish (*sebastes melanops*). NOAA Fishery Bulletin **102**: 418--429.
- Bobko, S.J., and Berkeley, S.A. 2004b. Maturity, ovarian cycle, fecundity, and age-specific parturition of black rockfish (*sebastes melanops*). Fishery Bulletin **102**(3): 418–429. Available from <http://aquaticcommons.org/15063/> [accessed 23 October 2015].
- Boehlert, G.W., and Yoklavich, M.M. 1983. Effects of temperature ration and fish size on growth of juvenile black rockfish, *sebastes melanops*. Environmental Biology of Fishes **8**(1): 17--28.
- Campana, S.E. 1999. Chemistry and composition of fish otoliths: Pathways, mechanisms and applications. Marine Ecology Progress Series **188**. Available from <https://doi.org/10.3354/meps188263> [accessed].
- Cope, J.M., Berger, A.M., Whitman, A.D., Budrick, J.E., Bosley, K.M., Tsou, S., Niles, C.B., Privitera-Johnson, K., Hillier, L.K., and Hinton, K.E. 2019. Assessing cabezon (*scorpaenichthys marmoratus*) stocks in waters off of California and Oregon, with catch limit estimation for Washington state. : 439. Available from <https://www.pcouncil.org/documents/2019/10/assessing-cabezon-scorpaenichthys-marmoratus-stocks-in-waters-off-of-california-and-oregon-with-catch-limit-estimation-for-washington-state-10-18-2019.pdf/>.
- Cope, J.M., and Hamel, O.S. 2022. Upgrading from m version 0.2: An application-based method for practical estimation, evaluation and uncertainty characterization of natural mortality. Fisheries Research **256**: 106493. doi:10.1016/j.fishres.2022.106493.

Cope, J.M., Sampson, D.B., Stephens, A., Key, M., Mirick, P.P., Stachura, M.M., Tsou, T.-S., Weyland, P., Berger, A.M., Buell, T., Councill, E., Dick, E.J., Fenske, K.H., Monk, M., and Rodomsky, B.T. 2016. Assessments of California, Oregon, and Washington stocks of black rockfish (*Sebastodes melanops*) in 2015. Pacific Fishery Management Council, Pacific Fishery Management Council, 7700 Ambassador Place NE, Suite 200, Portland, OR 97220.

Cope, J.M., and Whitman, A.D. 2021. Status of vermillion rockfish (*sebastes miniatus*) along the US west - oregon coast in 2021. Pacific Fisheries Management Council, Portland, OR.

Council, P.F.M. 2023. Report of the pre-assessment workshop for 2023 groundfish stock assessments of copper rockfish, canary rockfish and black rockfish. Available from <https://www.pcouncil.org/documents/2023/04/report-of-the-pre-assessment-workshop-for-2023-groundfish-stock-assessments-of-copper-rockfish-canary-rockfish-and-black-rockfish.pdf/>.

Culver, B.N. 1987. Results of tagging black rockfish (*sebastes melanops*) off the washington and northern oregon coast. University of Alaska Sea Grant.

Dick, E.J. 2009. Modeling the Reproductive Potential of Rockfishes (*sebastes* Spp.). PhD Thesis, University of California at Santa Cruz.

Dick, E.J., Berger, A., Bizzarro, J., Bosley, K., Cope, J., Field, J., Gilbert-Horvath, L., Grunloh, N., Ivens-Duran, M., Miller, R., Privitera-Johnson, K., and Rodomsky, B.T. 2018. The combined status of blue and deacon rockfishes in u.s. Waters off california and oregon in 2017. Pacific Fisheries Management Council, Portland, OR.

Dorn, M.W. 2002. Advice on West Coast rockfish harvest rates from Bayesian meta-analysis of stock-recruit relationships. North American Journal of Fisheries Management **22**(1): 280–300.

Douglas, D.A. 1998. Species composition of rockfish in catches by oregon trawlers, 1963-93. Oregon Department of Fish; Wildlife.

Echeverria, T.W. 1987. Thirty-four species of califonia rockfishes: Maturity and seasonality of reproduction. Fishery Bulletin **85**(2): 229–250.

Fennie, H.W., Grorud-Colvert, K., and Sponaugle, S. 2023. Larval rockfish growth and survival in response to anomalous ocean conditions. Sci Rep **13**(1): 4089. doi:10.1038/s41598-023-30726-5.

Fisheries, D. of, and Canada, O. 2014. Pacific region integrated fisheries management plan: groundfish. Available from <http://www.dfo-mpo.gc.ca/fm-gp/peches-fisheries/ifmp-gmp/index-eng.htm>.

- Fournier, D. 1997. An introduction to AD model builder for use in nonlinear modeling and statistics. Otter Research Ltd. PO Box 2040, Sidney, B.C. V8L 3S3 Canada. Available from <http://www.dfo-mpo.gc.ca/fm-gp/peches-fisheries/ifmp-gmp/index-eng.htm>.
- Francis, R.I.C.C., and Hilborn, R. 2011. Data weighting in statistical fisheries stock assessment models. Canadian Journal of Fisheries and Aquatic Sciences **68**(6): 1124–1138. doi:10.1139/f2011-025.
- Green, K.M., and Starr, R.M. 2011. Movements of small adult black rockfish: Implications for the design of MPAs. Marine Ecology Progress Series **414**: 219–230.
- Hamel, O.S., and Cope, J.M. 2022. Development and considerations for application of a longevity-based prior for the natural mortality rate. Fisheries Research **256**: 106477. doi:10.1016/j.fishres.2022.106477.
- Hess, J.E., Hyde, J.R., and Moran, P. 2023. Comparative phylogeography of a bathymetrically segregated pair of sister taxa of rockfishes (genus *sebastes*): Black rockfish, *sebastes melanops*, and yellowtail rockfish, *sebastes flavidus*. Mar Biol **170**(5): 62. doi:10.1007/s00227-023-04207-2.
- Hixon, M.A., Johnson, D.W., and Sogard, S.M. 2014a. BOFFFFs: On the importance of conserving old-growth age structure in fishery populations. ICES Journal of Marine Science **71**(8): 2171–2185.
- Hixon, M.A., Johnson, D.W., and Sogard, S.M. 2014b. BOFFFFs: On the importance of conserving old-growth age structure in fishery populations. ICES Journal of Marine Science **71**(8): 2171–2185. doi:10.1093/icesjms/fst200.
- Høie, H. E. Otterlei, and Folkvord, A. 2004. Temperature-dependent fractionation of stable oxygen isotopes in otoliths of juvenile cod (*gadus morhua l.*). ICES Journal of Marine Science **61**. Available from <https://doi.org/10.1016/j.icesjms.2003.11.006> [accessed].
- Iacumin, P, G. Bianucci, and Longinelli, A. 1992. Oxygen and carbon isotopic composition of fish otoliths. Marine Biology **113**. Available from <https://doi.org/10.1007/BF00349696> [accessed].
- Karnowski, M., Gertseva, V.V., and Stephens, A. 2014. Historical Reconstruction of Oregon's Commercial Fisheries Landings. Oregon Department of Fish; Wildlife, Salem, OR.
- Krutzikowksy, G., D. W. Wagman, and Davis, R. 2019. Population status of black rockfish in oregon coastal waters. Annual Progress Report, Oregon Department of Fish; Wildlife.
- Langseth, B.J., Wetzel, C.R., Cope, J.M., and Whitman, A.D. 2021. Status of quillback rockfish (*sebastes maliger*) in u.s. Waters off the coast of oregon in 2021 using catch and length data. Pacific Fisheries Management Council, Portland, OR.

Love, M.S. 1957. Certainly more than you want to know about the fisheries of the pacific coast: A postmodern experience. Really Big Press, Santa Barbara, CA.

Love, M.S., Yoklavich, M., and Thorsteinson, L. 2002. The rockfishes of the northeast pacific. In 1st Edition. University of California Press, Berkeley.

Manual on generalized age determination procedures for groundfish. 2006. Committee of Age Reading Experts. Available from <https://care.psmfc.org/about/ageing-manual/>.

McAllister, M.K., and Ianelli, J.N. 1997. Bayesian stock assessment using catch-age data and the sampling - importance resampling algorithm. Canadian Journal of Fisheries and Aquatic Sciences **54**: 284–300.

McClure, M.M., Haltuch, M.A., Willis-Norton, E., Huff, D.D., Hazen, E.L., Crozier, L.G., Jacox, M.G., Nelson, M.W., Andrews, K.S., Barnett, L.A.K., Berger, A.M., Beyer, S., Bizzarro, J., Boughton, D., Cope, J.M., Carr, M., Dewar, H., Dick, E., Dorval, E., Dunham, J., Gertseva, V., Greene, C.M., Gustafson, R.G., Hamel, O.S., Harvey, C.J., Henderson, M.J., Jordan, C.E., Kaplan, I.C., Lindley, S.T., Mantua, N.J., Matson, S.E., Monk, M.H., Moyle, P., Nicol, C., Pohl, J., Rykaczewski, R.R., Samhouri, J.F., Sogard, S., Tolimieri, N., Wallace, J., Wetzel, C., and Bograd, S.J. 2023. Vulnerability to climate change of managed stocks in the california current large marine ecosystem. Frontiers in Marine Science **10**. Available from <https://www.frontiersin.org/articles/10.3389/fmars.2023.1103767> [accessed 28 February 2023].

Monk, M., Dick, E.J., Buell, T., ZumBrunnen, L., Dauble, A.D., and Pearson, D. 2013. Documentation of a relational database for the oregon sport groundfish onboard sampling program. U.S. Department of Commerce. Available from <https://users.soe.ucsc.edu/~msmangel/Monk%20et%20al%20Sept%202013.pdf> [accessed 9 April 2019].

Monnahan, C.C., Branch, T.A., Thorson, J.T., Stewart, I.J., and Szuwalski, C.S. 2019. Overcoming long bayesian run times in integrated fisheries stock assessments. ICES Journal of Marine Science **76**(6): 1477–1488. doi:10.1093/icesjms/fsz059.

Myers, R.A., Bridson, J., and Barrowman, N.J. 1995. Summary of worldwide spawner and recruitment data. Canadian Technical Report of Fisheries and Aquatic Sciences, Department of Fisheries; Oceans Canada, Northwest Atlantic Fisheries Centre, St. Johns, Newfoundland.

Oregon department of fish and wildlife informational report regarding speciation of unspecified rockfish landings in oregon for inclusion in stock assessmetn time series of removals. 2017, March. Pacific Fishery Management Council. Available from https://www.pcouncil.org/documents/2017/03/i2a_odfw_rpt_speciation_of_oregon_urck_and_pop1_mar2017bb.pdf [accessed 30 June 2021].

- Parker, S.J., Rankin, P.S., Olson, J.M., and Hannah, R.W. 1995. Movement patterns of black rockfish *Sebastodes melanops* in oregon coastal waters. In Biology, assessment, and management of north pacific rockfishes. Alaska Sea Grant, University of Alaska Fairbanks. pp. 51–72.
- Punt, A.E., Smith, D.C., KrusicGolub, K., and Robertson, S. 2008. Quantifying age-reading error for use in fisheries stock assessments, with application to species in Australia's southern and eastern scalefish and shark fishery. Canadian Journal of Fisheries and Aquatic Sciences **65**(9): 1991–2005. doi:10.1139/F08-111.
- Ralston, S., and Dick, E.J. 2003. The status of black rockfish *Sebastes melanops* off oregon and northern california in 2003. Pacific Fishery Management Council, 7700 Ambassador Place NE, Suite 200, Portland, OR.
- Rasmussen, L., Blume, M.T.O., Lawrence, K.A., Bailey, E.J., Terwilliger, M.R., Fields, S.A., Rankin, P.S., and Phillips, A.J. 2022. Oregon nearshore semi-pelagic rockfish survey. Report to SSC methodology review September 2022, Oregon Department of Fish; Wildlife.
- Rasmussen, L.K., Lawrence, K.A., Krutzikowsky, G.K., Watson, J.L., Aylesworth, L., Hannah, R.W., Rodomsky, B.T., Huntington, B.E., Matteson, K.M., and Easton, R.R. 2020. Nine years of video landers at the oregon department of fish & wildlife's marine resources program. Informational Report, Oregon Department of Fish; Wildlife, Salem, OR.
- Rasmussen, L., Terwilliger, M.R., Bailey, E.J., Blume, M.T.O., and Lawrence, M.T.O. 2023. Finding oregon's old female rockfish. Science Bulletin, Oregon Department of Fish; Wildlife.
- Rasmussen, S.A.F., L. K., and Rankin, P.S. 2022a. Combined video-hydroacoustic survey of nearshore semi-pelagic rockfish in untrawlable habitats. ICES Journal of Marine Science **79**: 100–116.
- Rasmussen, S.R.M., L. K., and Rankin, P.S. 2022b. Influence of near bottom fish distribution on the efficacy of a combined hydroacoustic video survey. ICES Journal of Marine Science **79**: 2069–2083.
- Rogers, J.B. 2003. Species allocation of *Sebastes* and *sebastolobus* species caught by foreign countries off Washington, Oregon, and California, U.S.A. In 1965-1976. Unpublished document.
- Sampson, D.B. 2007. The status of black rockfish off oregon and california in 2007. Pacific Fishery Management Council, 7700 Ambassador Place NE, Suite 200, Portland, OR.
- Starr, R.M., and Green, K. 2007. Species composition, relative abundance, and movements of important nearshore fish species along the north central california coast. Final report to the Pacific States Marine Fisheries Commission.

Stein, D., and Hassler, T.J. 1989. Species profiles: Life histories and environmental requirements of coastal fishes and invertebrates (pacific southwest): Brown rockfish, copper rockfish, black rockfish. U.S. Fish and Wildlife Service Biological Report.

Stewart, E.M. 1993. Status of black rockfish off the northern oregon coast in 1993. In Appendices to the status of the pacific coast groundfish fishery through 1993 and recommended acceptable biological catches for 1994. Stock Assessment and Fishery Evaluation, Pacific Fishery Management Council, 2130 SW Fifth Ave., Suite 224, Portland, OR.

Taylor, I.G., Johnson, K.F., Langseth, B.J., Stephens, A., Lam, L.S., Monk, M.H., Whitman, A.D., and Haltuch, M.A. 2021. Status of lingcod (*ophiodon elongatus*) along the northern u.s. West coast in 2021. Pacific Fisheries Management Council, Portland, OR.

Terwilliger, M, L. Rasmuson, and Stern, R. 2023. Age validation of black rockfish, copper rockfish, and cabezon using secondary ion mass spectrometry (SIMS) to elucidate seasonal patterns in otolith stable oxygen isotopes. Environmental Biology of Fishes **106**. Available from <https://doi.org/10.1007/s10641-023-01392-3> [accessed].

Thorson, J.T., Johnson, K.F., Methot, R.D., and Taylor, I.G. 2017. Model-based estimates of effective sample size in stock assessment models using the Dirichlet-multinomial distribution. Fisheries Research **192**: 84–93. doi:10.1016/j.fishres.2016.06.005.

Thorson, J.T., Stewart, I.J., and Punt, A.E. 2012. nwfscAgeingError: A user interface in R for the Punt *et al.* (2008) method for calculating ageing error and imprecision. Available from: <http://github.com/pfmc-assessments/nwfscAgeingError/>.

Van Voorhees, D., Hoffman, A., Lowther, A., Buskirk, W.V., Weinstein, J., and White, J. 2000. An evaluation of alternative estimators of ocean boat fishing effort and catch in oregon. RecFIN. Available from http://old.recfin.org/lib/RecFIN_ORBS_MRFSS_Comparison.PDF.

Wallace, F.R., Hoffman, A., and Tagart, J.V. 1999. Status of the black rockfish resource in 1999. Pacific Fishery Management Council, 7700 Ambassador Place NE, Suite 200, Portland, OR.

Wallace, F.R., and Tagart, J.V. 1994. Status of the coastal black rockfish stocks in washington and northern oregon in 1994. Pacific Fishery Management Council, 7700 Ambassador Place NE, Suite 200, Portland, OR.

Wallace, F.R., Tsou, T.S., Cheng, Y.W., and Wargo, L. 2010. Summary of the coastal black rockfish tagging program 1981-2008. Technical report No. FPT 11-02, State of Washington Department of Fish; Wildlife.

Wallace, Y.W., F. R. aand Cheng, and Tsou, T.S. 2007. Status of the black rockfish resource north of cape falcon, oregon to the u.s.-canadian border in 2006. Pacific Fishery Management Council, 7700 Ambassador Place NE, Suite 200, Portland, OR.

Wetzel, C.R., Langseth, B.J., Cope, J.M., and Whitman, A.D. 2021. The status of copper rockfish (*sebastes caurinus*) in u.s. Waters off the coast of oregon in 2021 using catch and length data. Pacific Fisheries Management Council, Portland, OR.

Whitman, A.D. 2023. Oregon historical marine recreational catch reconstruction (1979 - 2000). Oregon Department of Fish; Wildlife, Salem, OR.

Yamanaka, K.L., and Lacko, L.C. 2001. Inshore rockfish (*sebastes ruberrimus*, s. Malinger, s. Caurinus, s. Melanops, s. Nigrocinctus, and s. Nebulosus) stock assessment for the west coast of canada and recommendations for management. Research Document, Fisheries; Oceans Canada, Nanaimo, BC. Available from https://www.dfo-mpo.gc.ca/csas-sccs/publications/resdocs-docrech/2001/2001_139-eng.htm.

7 Tables

Table 1: Landings and discards by year and fleet

Year	Trawl	Non-Trawl	Ocean Boat	Shore	Total
1892	0.00	0.25	0.00	0.25	0.50
1893	0.00	0.25	0.00	0.25	0.50
1894	0.00	0.25	0.00	0.25	0.50
1895	0.00	0.06	0.00	0.06	0.12
1896	0.00	0.02	0.00	0.02	0.04
1897	0.00	0.02	0.00	0.02	0.04
1898	0.00	0.01	0.00	0.01	0.02
1899	0.00	0.02	0.00	0.02	0.04
1900	0.00	0.02	0.00	0.02	0.04
1901	0.00	0.03	0.00	0.03	0.06
1902	0.00	0.03	0.00	0.03	0.06
1903	0.00	0.04	0.00	0.04	0.08
1904	0.00	0.05	0.00	0.05	0.10
1905	0.00	0.05	0.00	0.05	0.10
1906	0.00	0.06	0.00	0.06	0.12
1907	0.00	0.06	0.00	0.06	0.12
1908	0.00	0.07	0.00	0.07	0.14
1909	0.00	0.08	0.00	0.08	0.16
1910	0.00	0.08	0.00	0.08	0.16
1911	0.00	0.09	0.00	0.09	0.18
1912	0.00	0.10	0.00	0.10	0.20
1913	0.00	0.10	0.00	0.10	0.20
1914	0.00	0.11	0.00	0.11	0.22
1915	0.00	0.11	0.00	0.11	0.22
1916	0.00	0.12	0.00	0.12	0.24
1917	0.00	0.13	0.00	0.13	0.26
1918	0.00	0.13	0.00	0.13	0.26
1919	0.00	0.14	0.00	0.14	0.28
1920	0.00	0.15	0.00	0.15	0.30
1921	0.00	0.15	0.00	0.15	0.30

Table 1: Landings and discards by year and fleet (*continued*)

Year	Trawl	Non-Trawl	Ocean Boat	Shore	Total
1922	0.00	0.16	0.00	0.16	0.32
1923	0.00	0.16	0.00	0.16	0.32
1924	0.00	0.17	0.00	0.17	0.34
1925	0.00	0.18	0.00	0.18	0.36
1926	0.00	0.18	0.00	0.18	0.36
1927	0.00	0.19	0.00	0.19	0.38
1928	0.00	0.33	0.00	0.33	0.66
1929	0.00	0.89	0.00	0.88	1.77
1930	0.00	1.13	0.00	1.12	2.25
1931	0.00	0.68	0.00	0.67	1.34
1932	0.28	0.14	0.00	0.14	0.56
1933	0.17	0.26	0.00	0.26	0.69
1934	0.01	0.32	0.00	0.32	0.65
1935	0.14	0.21	0.00	0.21	0.56
1936	0.50	0.85	0.00	0.84	2.19
1937	0.97	1.74	0.00	1.73	4.44
1938	0.00	1.78	0.00	1.77	3.55
1939	1.37	1.97	0.00	1.95	5.29
1940	1.32	2.61	0.00	2.59	6.52
1941	3.43	2.14	0.00	2.12	7.69
1942	5.28	2.94	0.00	2.92	11.15
1943	32.62	3.90	0.00	3.87	40.40
1944	75.31	4.17	0.00	4.14	83.62
1945	125.30	4.63	0.00	4.59	134.52
1946	88.26	4.25	0.00	4.22	96.74
1947	29.37	1.62	0.00	1.61	32.60
1948	17.74	2.75	0.00	2.73	23.22
1949	5.92	1.65	0.00	1.64	9.21
1950	56.44	1.94	0.00	1.92	60.30
1951	15.02	1.18	0.00	1.17	17.37
1952	5.66	1.69	0.00	1.68	9.03

Table 1: Landings and discards by year and fleet (*continued*)

Year	Trawl	Non-Trawl	Ocean Boat	Shore	Total
1953	19.22	0.89	0.00	0.88	20.99
1954	24.92	0.83	0.00	0.82	26.57
1955	33.37	1.57	0.00	1.56	36.51
1956	23.05	0.90	0.00	0.89	24.83
1957	22.48	1.53	0.00	1.52	25.54
1958	33.53	0.64	0.00	0.64	34.81
1959	41.18	1.02	0.00	1.01	43.21
1960	73.43	1.55	0.00	1.54	76.53
1961	60.25	1.81	0.00	1.80	63.86
1962	58.02	1.44	0.00	1.43	60.89
1963	45.43	1.52	0.00	1.51	48.46
1964	28.08	0.69	0.00	0.68	29.44
1965	48.29	2.94	0.00	2.92	54.15
1966	43.47	2.06	0.00	2.04	47.57
1967	26.46	4.39	0.00	4.36	35.21
1968	23.67	4.14	0.00	4.11	31.92
1969	49.51	8.27	0.00	8.20	65.98
1970	29.71	3.89	0.00	3.86	37.46
1971	32.32	8.19	0.00	8.12	48.63
1972	43.24	10.61	0.00	10.53	64.39
1973	38.43	11.32	49.59	11.23	110.57
1974	24.17	13.94	98.34	13.83	150.28
1975	26.52	7.49	147.10	7.43	188.54
1976	5.37	9.62	195.86	9.54	220.38
1977	7.18	11.65	244.61	11.56	275.00
1978	10.11	16.35	293.37	16.22	336.05
1979	27.87	39.97	341.29	39.65	448.78
1980	14.25	25.15	237.96	28.09	305.45
1981	21.77	22.26	419.97	12.57	476.56
1982	39.27	45.44	556.80	9.85	651.36
1983	2.98	95.32	386.14	14.59	499.03

Table 1: Landings and discards by year and fleet (*continued*)

Year	Trawl	Non-Trawl	Ocean Boat	Shore	Total
1984	16.54	63.96	444.91	7.97	533.38
1985	14.14	61.03	258.11	8.68	341.97
1986	7.57	45.62	242.37	13.15	308.71
1987	0.51	116.43	214.23	26.62	357.80
1988	2.03	90.03	239.73	62.48	394.27
1989	1.60	121.95	322.84	11.37	457.76
1990	0.40	171.42	316.20	13.84	501.85
1991	0.04	179.63	145.93	16.41	342.01
1992	10.44	343.27	267.26	19.08	640.06
1993	4.92	118.90	397.50	21.86	543.18
1994	39.55	149.79	279.91	16.30	485.55
1995	2.08	94.78	336.26	12.36	445.48
1996	0.04	144.76	362.78	11.73	519.31
1997	1.89	172.99	362.21	13.28	550.38
1998	11.14	132.21	318.08	5.67	467.09
1999	0.32	125.70	155.04	6.54	287.60
2000	0.64	107.14	212.69	15.75	336.23
2001	0.18	147.85	290.72	18.67	457.42
2002	1.23	126.32	284.61	27.34	439.51
2003	0.36	116.62	321.18	19.53	457.69
2004	1.47	117.43	324.27	7.00	450.17
2005	0.52	100.29	303.88	6.81	411.49
2006	2.29	93.67	259.11	13.28	368.35
2007	0.32	101.85	247.43	13.28	362.88
2008	0.19	99.19	227.67	13.28	340.32
2009	0.22	134.48	277.62	13.28	425.59
2010	0.34	101.12	292.14	13.28	406.89
2011	0.76	97.54	198.72	13.28	310.30
2012	0.76	96.72	210.14	13.28	320.90
2013	0.08	107.47	321.31	13.28	442.14
2014	0.71	122.92	342.55	13.28	479.45

Table 1: Landings and discards by year and fleet (*continued*)

Year	Trawl	Non-Trawl	Ocean Boat	Shore	Total
2015	0.51	122.12	458.61	13.28	594.52
2016	0.64	105.95	404.65	13.28	524.53
2017	0.26	125.21	402.67	13.28	541.43
2018	0.03	122.98	278.78	13.28	415.07
2019	0.01	119.19	305.07	13.28	437.55
2020	0.04	101.79	320.45	13.28	435.56
2021	0.00	101.53	320.73	13.28	435.54
2022	0.00	118.11	394.61	13.28	526.00

Table 2: Sampled commercial length composition by year, fleet and sex.

Year	Trawl Female	Trawl Male	Trawl Total	Non- trawl Female	Non- trawl Male	Non- trawl Un- sexed	Non- trawl Total
1985	283	321	604	0	0	0	0
1992	0	0	0	79	124	0	203
1994	19	22	41	0	0	0	
1995	0	0	0	214	220	5	439
1996	0	0	0	75	153	0	228
1997	62	105	167	107	167	0	274
1998	25	43	68	100	213	0	313
1999	0	0	0	58	94	0	152
2000	0	0	0	280	323	0	603
2001	17	3	20	511	518	0	1029
2002	0	0	0	670	569	93	1332
2003	18	25	43	631	694	46	1371
2004	38	30	68	1811	1687	0	3498
2005	28	8	36	1017	1208	0	2225
2006	83	89	172	2097	2254	49	4400
2007	0	0	0	2033	1781	24	3838
2008	22	32	54	1495	1360	5	2860
2009	9	4	13	1438	1417	76	2931
2010	14	15	29	1895	2109	224	4228
2011	35	77	112	2368	1991	0	4359
2012	33	48	81	1802	1642	53	3497
2013	3	2	5	2249	1967	60	4276
2014	68	55	123	3446	3238	57	6741
2015	42	28	70	3001	3334	18	6353
2016	56	70	126	3266	2890	11	6167
2017	24	20	44	2787	2461	4	5252
2018	5	7	12	2825	2235	22	5082
2019	4	1	5	2106	2506	31	4643
2020	5	4	9	834	822	13	1669
2021	0	0	0	1703	1494	2	3199
2022	0	0	0	1996	2010	0	4006

Table 3: Available recreational fishery length samples by year, sex and fleet

Year	Ocean Female	Ocean Male	Ocean Un- known	Ocean Total	Shore Female	Shore Male	Shore Un- known	Shore Total
1980			653	653			324	324
1981			475	475			149	149
1982			950	950			110	110
1983			314	314			129	129
1984			1384	1384			178	178
1985			1826	1826			298	298
1986			1336	1336			309	309
1987			913	913			168	168
1988			1409	1409			267	267
1989			1041	1041			113	113
1993			1893	1893			343	343
1994			2233	2233			263	263
1995			2172	2172			200	200
1996			2228	2228			261	261
1997			3080	3080			279	279
1998			3916	3916			84	84
1999	1803	1805	5251	8859			199	199
2000	2318	2515	4259	9092			163	163
2001	1656	1445	5671	8772			0	0
2002	1966	1791	3712	7469			0	0
2003	1747	1714	3637	7098			0	0
2004	1652	1609	2754	6015			0	0
2005	1538	1587	3954	7079			0	0
2006	1050	1123	7295	9468			0	0
2007	1020	1032	10741	12793			0	0
2008		1	11764	11765			0	0
2009	565	528	11344	12437			0	0
2010	585	576	11901	13062			0	0

Table 3: Available recreational fishery length samples by year, sex and fleet (*continued*)

Year	Ocean Female	Ocean Male	Ocean Un- known	Ocean Total	Shore Female	Shore Male	Shore Un- known	Shore Total
2011	540	567	11750	12857				0
2012	523	544	12394	13461				0
2013	516	522	12652	13690				0
2014	553	530	11709	12792				0
2015	530	548	12784	13862				0
2016	537	561	10596	11694				0
2017	442	549	10233	11224				0
2018	584	619	11825	13028				0
2019	578	576	11727	12881				0
2020	550	560	989	2099				0
2021	533	562	9328	10423				0
2022	557	592	7857	9006				0

Table 4: Sampled commercial ages by year, fleet and sex

1985	283	321	604				
1992				52	91		143
1994	19	22	41				
1995							
1996							
1997							
1998	16	20	36	44	114		158
1999							
2000				134	153		287
2001				99	106		205
2002				172	144		316
2003	18	25	43	218	225		443
2004	38	30	68	206	179		385
2005				132	178		310
2006	83	88	171	286	281	34	601
2007				311	324		635
2008	22	32	54	281	284		565
2009	9	4	13	401	394	31	826
2010	14	14	28	402	369	65	836
2011	31	71	102	440	338		778
2012	15	26	41	376	367	4	747
2013	3	2	5	223	203	5	431
2014	53	14	67	558	299	11	868
2015	14	10	24	492	429	6	927
2016	29	21	50	473	426	1	900
2017	18	6	24	524	456	2	982
2018	2	4	6	570	441	10	1021
2019	3	1	4	406	463	8	877
2020	5	4	9	489	450	13	952
2021				467	440	1	908

Table 4: Sampled commercial ages by year, fleet and sex (*continued*)

Year	Trawl	Trawl	Total	Non-	Non-	Non-	Total
	Female	Male		trawl	trawl	trawl	
			Female	Male	Unsexed		
2022				479	422		901

Table 5: Age samples by year and sex for the Ocean Boat fleet. There are no ages for the recreational Shore fleet

Year	Female	Male	Unknown	Total
1999	1804	1806	160	3770
2000	2318	2515	24	4857
2001	1656	1445	57	3158
2002	1966	1791	5	3762
2003	1747	1714	6	3467
2004	1688	1649	0	3337
2005	1538	1587	9	3134
2006	1049	1122	15	2186
2007	1020	1032	9	2061
2009	565	527	0	1092
2010	585	576	17	1178
2011	540	566	22	1128
2012	523	543	25	1091
2013	516	522	10	1048
2014	553	530	10	1093
2015	530	548	11	1089
2016	537	561	3	1101
2017	442	549	2	993
2018	584	619	0	1203
2019	578	576	5	1159
2020	550	560	8	1118
2021	533	562	7	1102
2022	557	592	4	1153

Table 6: Number of commercial logbook samples (trips) with and without detecting the target species, Black Rockfish.

Year	Samples with Target	Samples without Target	Total Samples	Percent Positive
2004	1143	68	1211	0.94
2005	1164	91	1255	0.93
2006	1159	65	1224	0.95
2007	1028	24	1052	0.98
2008	1112	68	1180	0.94
2009	998	43	1041	0.96
2010	1320	33	1353	0.98
2011	1427	68	1495	0.95
2012	1387	51	1438	0.96
2013	1621	101	1722	0.94
2014	1368	40	1408	0.97
2015	1503	63	1566	0.96
2016	1167	53	1220	0.96
2017	1080	39	1119	0.97
2018	1311	52	1363	0.96
2019	1280	44	1324	0.97
2020	1085	53	1138	0.95
2021	1001	21	1022	0.98
2022	1117	83	1200	0.93

Table 7: Model construction (variables included) and selection for the nearshore logbook index.

Black RF Limit	Depth (binned)	Month	Num. People	Port	Year	Effort Offset	DF	Neg. Log- Likelihood	AICc	Delta AICc
yes	yes	yes	yes	yes	yes	yes	77	138210.0	276574.5	0.0
yes	no	yes	yes	yes	yes	yes	75	138278.8	276708.0	133.5
yes	yes	no	yes	yes	yes	yes	66	138306.2	276744.7	170.2
yes	no	no	yes	yes	yes	yes	64	138378.0	276884.4	309.9
no	yes	yes	yes	yes	yes	yes	56	138400.3	276912.8	338.3
yes	yes	yes	no	yes	yes	yes	62	138428.8	276981.9	407.4
no	no	yes	yes	yes	yes	yes	54	138468.9	277046.0	471.5
yes	yes	no	no	yes	yes	yes	51	138519.0	277140.2	565.7
yes	no	yes	no	yes	yes	yes	60	138523.6	277167.5	593.0
no	yes	no	yes	yes	yes	yes	45	138548.8	277187.8	613.3
no	yes	yes	no	yes	yes	yes	41	138614.8	277311.7	737.2
no	no	no	yes	yes	yes	yes	43	138616.6	277319.3	744.8
yes	no	no	no	yes	yes	yes	49	138617.2	277332.5	758.0
no	no	yes	no	yes	yes	yes	39	138709.4	277496.9	922.4
no	yes	no	no	yes	yes	yes	30	138750.5	277561.2	986.7
yes	no	yes	yes	no	yes	yes	67	138724.7	277583.8	1009.3
yes	yes	yes	yes	no	yes	yes	69	138723.7	277585.8	1011.3
no	no	no	no	yes	yes	yes	28	138844.3	277744.7	1170.2
yes	no	no	yes	no	yes	yes	56	138878.3	277868.9	1294.4
yes	yes	no	yes	no	yes	yes	58	138876.7	277869.7	1295.2
no	no	yes	yes	no	yes	yes	46	138905.2	277902.6	1328.1
no	yes	yes	yes	no	yes	yes	48	138904.5	277905.2	1330.7
no	yes	no	yes	no	yes	yes	37	139148.5	278371.1	1796.6
no	no	no	yes	no	yes	yes	35	139150.5	278371.2	1796.7
yes	no	yes	no	no	yes	yes	52	139145.8	278395.8	1821.3
yes	yes	yes	no	no	yes	yes	54	139144.5	278397.3	1822.8
no	no	yes	no	no	yes	yes	31	139317.8	278697.6	2123.1
no	yes	yes	no	no	yes	yes	33	139316.1	278698.2	2123.7

Table 7: Model construction (variables included) and selection for the nearshore logbook index (*continued*)

Black RF Limit	Depth (binned)	Month	Num. People	Port	Year	Effort Offset	DF	Neg. Log- Likelihood	AICc	Delta AICc
yes	no	no	no	no	yes	yes	41	139311.1	278704.3	2129.8
yes	yes	no	no	no	yes	yes	43	139310.4	278707.0	2132.5
no	no	no	no	no	yes	yes	20	139582.5	279205.0	2630.5
no	yes	no	no	no	yes	yes	22	139581.9	279207.9	2633.4

Table 8: Number of ORBS samples (trips) with and without detecting the target species, Black Rockfish.

Year	Samples with Target	Samples without Target	Total Samples	Percent Positive
2001	3539	304	3843	0.92
2002	3519	430	3949	0.89
2003	3911	346	4257	0.92
2004	3158	261	3419	0.92
2005	6197	436	6633	0.93
2006	6247	516	6763	0.92
2007	4374	354	4728	0.93
2008	5062	400	5462	0.93
2009	5122	336	5458	0.94
2010	5679	349	6028	0.94
2011	4794	569	5363	0.89
2012	4707	604	5311	0.89
2013	6119	931	7050	0.87
2014	5190	516	5706	0.91
2015	7552	643	8195	0.92
2016	6170	520	6690	0.92
2017	6664	471	7135	0.93
2018	5866	1075	6941	0.85
2019	5375	736	6111	0.88
2020	5869	1063	6932	0.85
2021	4964	688	5652	0.88
2022	5591	748	6339	0.88

Table 9: Model construction (variables included) and selection for the ORBS dockside survey.

Boat Type	Open Depths	Month	Port	Bag Limit	Year	Effort Offset	DF	Neg. Log-Likelihood	AICc	Delta AICc
yes	yes	yes	yes	yes	yes	yes	49	330152.7	660403.5	0.0
yes	no	yes	yes	yes	yes	yes	46	330188.0	660468.0	64.5
yes	yes	yes	yes	no	yes	yes	45	330210.9	660511.8	108.3
yes	no	yes	yes	no	yes	yes	42	330243.5	660570.9	167.4
yes	yes	no	yes	yes	yes	yes	38	330341.5	660759.1	355.6
yes	yes	no	yes	no	yes	yes	34	330434.4	660936.8	533.3
yes	no	no	yes	yes	yes	yes	35	330507.3	661084.7	681.2
no	yes	yes	yes	yes	yes	yes	48	330546.4	661188.9	785.4
yes	no	no	yes	no	yes	yes	31	330593.7	661249.4	845.9
no	no	yes	yes	yes	yes	yes	45	330580.0	661250.0	846.5
no	yes	yes	yes	no	yes	yes	44	330606.2	661300.5	897.0
no	no	yes	yes	no	yes	yes	41	330636.4	661354.9	951.4
no	yes	no	yes	yes	yes	yes	37	330732.1	661538.3	1134.8
no	yes	no	yes	no	yes	yes	33	330824.8	661715.6	1312.1
no	no	no	yes	yes	yes	yes	34	330893.1	661854.1	1450.6
no	no	no	yes	no	yes	yes	30	330978.6	662017.2	1613.7
yes	yes	yes	no	yes	yes	yes	42	331937.7	663959.4	3555.9
no	yes	yes	no	yes	yes	yes	41	331939.6	663961.2	3557.7
yes	no	yes	no	yes	yes	yes	39	331965.9	664009.7	3606.2
no	no	yes	no	yes	yes	yes	38	331967.8	664011.6	3608.1
yes	yes	yes	no	no	yes	yes	38	331997.7	664071.5	3668.0
no	yes	yes	no	no	yes	yes	37	331999.8	664073.6	3670.1
yes	no	yes	no	no	yes	yes	35	332021.8	664113.7	3710.2
no	no	yes	no	no	yes	yes	34	332023.9	664115.7	3712.2
yes	yes	no	no	yes	yes	yes	31	332113.5	664289.0	3885.5
no	yes	no	no	yes	yes	yes	30	332115.3	664290.6	3887.1
yes	yes	no	no	no	yes	yes	27	332201.9	664457.8	4054.3
no	yes	no	no	no	yes	yes	26	332203.8	664459.6	4056.1

Table 9: Model construction (variables included) and selection for the ORBS dockside survey (*continued*)

Boat Type	Open Depths	Month	Port	Bag Limit	Year	Effort Offset	DF	Neg. Log-Likelihood	AICc	Delta AICc
no	no	no	no	yes	yes	yes	27	332317.6	664689.1	4285.6
yes	no	no	no	v	yes	yes	28	332316.7	664689.4	4285.9
no	no	no	no	no	yes	yes	23	332394.2	664834.3	4430.8
yes	no	no	no	no	yes	yes	24	332393.2	664834.4	4430.9

Table 10: Model construction (variables included) and selection for the Oregon onboard charter observer survey.

Depth (binned)	Open Depths	Month	Port	Year	Effort Offset	DF	Neg. Log-	AICc	Delta
							Likelihood	AICc	AICc
yes	yes	yes	yes	yes	yes	40	23266.1	46612.5	0.0
yes	no	yes	yes	yes	yes	37	23269.3	46612.8	0.4
no	no	yes	yes	yes	yes	32	23289.1	46642.5	30.0
no	yes	yes	yes	yes	yes	35	23286.1	46642.5	30.1
yes	yes	no	yes	yes	yes	33	23299.0	46664.3	51.8
yes	no	no	yes	yes	yes	30	23306.2	46672.7	60.2
no	yes	no	yes	yes	yes	28	23315.8	46687.7	75.3
no	no	no	yes	yes	yes	25	23321.9	46694.0	81.5
yes	yes	yes	yes	no	yes	23	23422.7	46891.6	279.1
no	yes	yes	yes	no	yes	18	23439.3	46914.6	302.1
yes	yes	no	yes	no	yes	16	23451.7	46935.5	323.1
no	yes	no	yes	no	yes	11	23466.4	46954.8	342.3
yes	no	yes	yes	no	yes	20	23458.9	46957.9	345.4
no	no	yes	yes	no	yes	15	23479.3	46988.6	376.1
yes	no	no	yes	no	yes	13	23493.3	47012.7	400.2
no	yes	yes	no	yes	yes	29	23489.0	47036.2	423.7
no	no	no	yes	no	yes	8	23510.2	47036.3	423.9
yes	yes	yes	no	yes	yes	34	23484.3	47036.9	424.4
no	no	yes	no	yes	yes	26	23492.8	47037.8	425.3
yes	no	yes	no	yes	yes	31	23487.9	47037.9	425.4
no	yes	no	no	yes	yes	22	23510.1	47064.3	451.8
no	no	no	no	yes	yes	19	23513.1	47064.3	451.8
yes	no	no	no	yes	yes	24	23508.5	47065.1	452.6
yes	yes	no	no	yes	yes	27	23505.9	47066.0	453.5
yes	yes	yes	no	no	yes	17	23636.7	47307.4	694.9
no	yes	yes	no	no	yes	12	23644.8	47313.6	701.2
yes	yes	no	no	no	yes	10	23661.4	47342.9	730.4
no	yes	no	no	no	yes	5	23669.0	47348.1	735.6

Table 10: Model construction (variables included) and selection for the Oregon onboard charter observer survey (*continued*)

Depth (binned)	Open Depths	Month	Port	Year	Effort Offset	DF	Neg.	AICc	Delta
							Log-	AICc	AICc
Likelihood									
yes	no	yes	no	no	yes	14	23681.4	47390.8	778.3
no	no	yes	no	no	yes	9	23689.5	47396.9	784.5
yes	no	no	no	no	yes	7	23703.2	47420.5	808.0
no	no	no	no	no	yes	2	23710.7	47425.5	813.0

Table 11: Number of Oregon marine reserves samples (drifts) with and without detecting the target species, Black Rockfish.

Year	Samples with Target	Samples without Target	Total Samples	Percent Positive
2011	19	5	24	0.79
2012	20	3	23	0.87
2013	68	3	71	0.96
2014	109	10	119	0.92
2015	134	28	162	0.83
2016	103	13	116	0.89
2017	70	25	95	0.74
2018	78	30	108	0.72
2019	63	32	95	0.66
2022	40	7	47	0.85

Table 12: Model construction (variables included) and selection for the Oregon marine reserves index.

Depth (binned)	Season	Site	Treat- ment	Year	Effort Offset	DF	Neg. Log- Likelihood	AICc	Delta AICc
yes	yes	yes	yes	yes	yes	19	2887.1	5813.2	0.0
yes	no	yes	yes	yes	yes	18	2891.0	5818.7	5.6
no	yes	yes	yes	yes	yes	16	2898.2	5829.1	15.9
no	no	yes	yes	yes	yes	15	2901.7	5833.9	20.7
yes	yes	no	yes	yes	yes	16	2912.4	5857.5	44.4
no	yes	no	yes	yes	yes	13	2916.9	5860.1	47.0
yes	no	no	yes	yes	yes	15	2916.6	5863.7	50.6
no	no	no	yes	yes	yes	12	2920.5	5865.4	52.2
yes	yes	yes	no	yes	yes	18	2935.1	5906.9	93.8
yes	no	yes	no	yes	yes	17	2937.0	5908.7	95.5
yes	yes	no	no	yes	yes	15	2948.2	5926.9	113.7
yes	no	no	no	yes	yes	14	2950.2	5929.0	115.8
no	yes	yes	no	yes	yes	15	2951.3	5933.1	120.0
no	no	yes	no	yes	yes	14	2952.8	5934.1	121.0
no	yes	no	no	yes	yes	12	2957.8	5940.0	126.9
no	no	no	no	yes	yes	11	2959.5	5941.3	128.2

Table 13: Ageing error models and resultant model selection (AICc) values for 9 models of bias and precision explored for each age reader used in the black rockfish assessment. Bolded text indicates indicate the chosen model. Model codes: 0= unbiased; 1 = Constant CV; 2 = Curvilinear (SD for precision); 3= Curvilinear CV

	Reader 1		Second Reader		Model selection			
Primary reader vs Reader 1								
Model	Bias	Precision	Bias	Precision	AICc	Δ AICc	BIC	Δ BIC
1	0	1	0	1	18480	61	18616	50
2	0	2	0	2	18424	5	18565	0
3	0	3	0	3	18485	66	18626	61
4	0	1	1	1	18466	46	18610	44
5	0	2	1	2	18428	9	18578	12
6	0	3	1	3	18419	0	18569	4
7	0	1	2	1	18463	44	18613	47
8	0	2	2	2	18430	11	18586	21
9	0	3	2	3	18419	0	18575	10
Primary reader vs Reader 2								
Model	Bias	Precision	Bias	Precision	AICc	Δ AICc	BIC	Δ BIC
1	0	1	0	1	1189	34	1222	38
2	0	2	0	2	1201	46	1234	49
3	0	3	0	3	1195	40	1228	43
4	0	1	1	1	1165	11	1197	13
5	0	2	1	2	1179	25	1209	25
6	0	3	1	3	1171	16	1201	16
7	0	1	2	1	1155	0	1185	0
8	0	2	2	2	1169	15	1196	12
9	0	3	2	3	1179	24	1205	21

Table 14: Ageing error models and resultant model selection (AICc) values for 9 models of bias and precision explored for each age reader used in the black rockfish assessment. Bolded text indicates indicate the chosen model. Model codes: 0= unbiased; 1 = Constant CV; 2 = Curvilinear (SD for precision); 3= Curvilinear CV

	Reader 1		Second Reader		Model selection			
Primary reader vs Reader 3								
Model	Bias	Precision	Bias	Precision	AICc	Δ AICc	BIC	Δ BIC
1	0	1	0	1	2591	12	2693	8
2	0	2	0	2	2579	0	2685	0
3	0	3	0	3	2590	11	2696	11
4	0	1	1	1	2593	14	2701	16
5	0	2	1	2	2589	10	2700	15
6	0	3	1	3	2592	13	2703	18
7	0	1	2	1	2596	17	2707	22
8	0	2	2	2	2588	9	2702	17
9	0	3	2	3	2598	18	2711	26
Primary reader vs Reader 4								
Model	Bias	Precision	Bias	Precision	AICc	Δ AICc	BIC	Δ BIC
1	0	1	0	1	559	0	592	0
2	0	2	0	2	563	5	596	4
3	0	3	0	3	567	8	600	7
4	0	1	1	1	567	8	599	6
5	0	2	1	2	574	15	604	11
6	0	3	1	3	564	5	594	1
7	0	1	2	1	571	13	601	9
8	0	2	2	2	576	18	603	11
9	0	3	2	3	577	18	603	11

Table 15: Ageing error models and resultant model selection (AICc) values for 9 models of bias and precision explored for each age reader used in the black rockfish assessment. Bolded text indicates indicate the chosen model. Model codes: 0= unbiased; 1 = Constant CV; 2 = Curvilinear (SD for precision); 3= Curvilinear CV

		Reader 1		Second Reader		Model selection			
Primary reader vs Reader 5									
Model		Bias	Precision	Bias	Precision	AICc	Δ AICc	BIC	Δ BIC
1		0	1	0	1	1521	0	1559	0
2		0	2	0	2	1551	29	1587	29
3		0	3	0	3	1528	7	1565	6
4		0	1	1	1	1529	8	1565	7
5		0	2	1	2	1560	39	1595	37
6		0	3	1	3	1537	16	1572	14
7		0	1	2	1	1530	9	1565	7
8		0	2	2	2	1533	11	1565	7
9		0	3	2	3	1552	31	1585	27
Primary reader vs Readers 5 & 6									
Model		Bias	Precision	Bias	Precision	AICc	Δ AICc	BIC	Δ BIC
1		0	1	0	1	1064	0	1128	0
2		0	2	0	2	1072	8	1140	12
3		0	3	0	3	1065	1	1133	5
4		0	1	1	1	1070	6	1139	11
5		0	2	1	2	1077	13	1150	22
6		0	3	1	3	1074	10	1147	19
7		0	1	2	1	1072	8	1145	17
8		0	2	2	2	1078	14	1154	26
9		0	3	2	3	1078	14	1154	26

Table 16: Estimated biological and functional length at 50% maturity, the 95% confidence interval (CI), and slope values with logistic regression and flexible spline models

Model	Length at 50% Maturity (95% CI)	Slope
Biological logistic	34.38 (33.90-34.86)	-0.31
Functionl logistic	40.36 (39.89-40.83)	-0.38
Biological splie	34.01 (32.10-35.41)	NA
Functional spline	40.29 (39.88-40.77)	NA

Table 17: Specifications and structure of the base model.

Model Setup	Base Model
Starting year	1892
<u>Population characteristics</u>	
Maximum age	40
Sex	2
Population lengths	4-64 cm by 2 cm bins
Summary biomass (mt)	Age 0+
<u>Data characteristics</u>	
Data lengths	10-64 cm by 2 cm bins
Data ages	0-40 ages
Minimum age for growth calculations	0
Maximum age for growth calculations	40
First mature age	0
Starting year of estimated recruitment in main period	1980
<u>Fishery characteristics</u>	
Fishing mortality method	Hybrid F
Maximum F	4
Catchability	
Non-trawl Index	Analytical estimate
Ocean Index	Analytical estimate
MPA Index	Analytical estimate
Acoustic Index	Parameter
Tagging Index	Analytical estimate
MRFSS Index	Analytical estimate
Commercial Trawl Selectivity	Length-Based Double Normal
Commercial Non-trawl Selectivity	Length-Based Double Normal
Recreational Ocean Selectivity	Length-Based Double Normal
Recreational Shore Selectivity	Length-Based Double Normal
MPA Survey Selectivity	Length-Based Double Normal
Acoustic Survey Selectivity	Length-Based Double Normal
Tagging Selectivity	Length-Based Logistic
Recreation MRFSS Selectivity	Mirrored to Ocean Fleet

Table 18: List of parameters used in the base model, including estimated values and standard deviations (SD), bounds (minimum and maximum), estimation phase (negative values not estimated), status (indicates if parameters are near bounds), and prior type information (mean and SD).

Parameter	Value	Phase	Bounds	Status	Prior (Exp.Val, SD)
NatM uniform Fem GP 1	0.19	-5	-	-	Log Norm (-1.89, 0.075)
L at Amin Fem GP 1	0	-5	-	-	None
L at Amax Fem GP 1	50.846	3	OK	0.263657	Normal (48.82, 0.3)
VonBert K Fem GP 1	0.182	3	OK	0.00186914	None
CV young Fem GP 1	0.1	-5	-	-	None
CV old Fem GP 1	0.1	-5	-	-	None
Wtlen 1 Fem GP 1	2.55e-05	-99	-	-	None
Wtlen 2 Fem GP 1	2.895	-99	-	-	None
Mat50% Fem GP 1	40.36	-99	-	-	None
Mat slope Fem GP 1	-0.38	-99	-	-	None
Eggs scalar Fem GP 1	1.41e-08	-3	-	-	None
Eggs exp len Fem GP 1	4.685	-3	-	-	None
NatM uniform Mal GP 1	0.17	-5	-	-	Log Norm (-1.89, 0.2)
L at Amin Mal GP 1	0	-3	-	-	None
L at Amax Mal GP 1	44.876	3	OK	0.26159	None
VonBert K Mal GP 1	0.224	3	OK	0.00297517	None
CV young Mal GP 1	0.1	-5	-	-	None
CV old Mal GP 1	0.1	-5	-	-	None
Wtlen 1 Mal GP 1	2.56e-05	-99	-	-	None
Wtlen 2 Mal GP 1	2.894	-99	-	-	None
CohortGrowDev	1	-1	-	-	None
FracFemale GP 1	0.5	-99	-	-	None
SR LN(R0)	8.245	1	OK	0.0173786	None
SR BH steep	0.72	-1	-	-	Log Norm (0.72, 0.24)
SR sigmaR	0.6	-6	-	-	None
SR regime	0	-99	-	-	None
SR autocorr	0	-99	-	-	None
Main RecrDev 1980	-1.08	3	act	0.392268	dev (NA, NA)

Table 18: List of parameters used in the base model, including estimated values and standard deviations (SD), bounds (minimum and maximum), estimation phase (negative values not estimated), status (indicates if parameters are near bounds), and prior type information (mean and SD). (*continued*)

Parameter	Value	Phase	Bounds	Status	Prior (Exp.Val, SD)
Main RecrDev 1981	-0.863	3	act	0.37042	dev (NA, NA)
Main RecrDev 1982	-0.317	3	act	0.27984	dev (NA, NA)
Main RecrDev 1983	-0.441	3	act	0.277759	dev (NA, NA)
Main RecrDev 1984	-0.277	3	act	0.218195	dev (NA, NA)
Main RecrDev 1985	-0.147	3	act	0.180332	dev (NA, NA)
Main RecrDev 1986	-0.333	3	act	0.176129	dev (NA, NA)
Main RecrDev 1987	-0.372	3	act	0.158349	dev (NA, NA)
Main RecrDev 1988	-0.297	3	act	0.132277	dev (NA, NA)
Main RecrDev 1989	-0.014	3	act	0.0972235	dev (NA, NA)
Main RecrDev 1990	-0.059	3	act	0.0850273	dev (NA, NA)
Main RecrDev 1991	0.078	3	act	0.0700834	dev (NA, NA)
Main RecrDev 1992	0.008	3	act	0.064519	dev (NA, NA)
Main RecrDev 1993	0.023	3	act	0.0599154	dev (NA, NA)
Main RecrDev 1994	0.344	3	act	0.0531871	dev (NA, NA)
Main RecrDev 1995	0.304	3	act	0.0566414	dev (NA, NA)
Main RecrDev 1996	0.024	3	act	0.0733322	dev (NA, NA)
Main RecrDev 1997	-0.023	3	act	0.0923232	dev (NA, NA)
Main RecrDev 1998	-0.299	3	act	0.144421	dev (NA, NA)
Main RecrDev 1999	0.631	3	act	0.0984838	dev (NA, NA)
Main RecrDev 2000	0.732	3	act	0.113712	dev (NA, NA)
Main RecrDev 2001	0.309	3	act	0.177524	dev (NA, NA)
Main RecrDev 2002	-0.447	3	act	0.332529	dev (NA, NA)
Main RecrDev 2003	0.542	3	act	0.126095	dev (NA, NA)
Main RecrDev 2004	-1.015	3	act	0.422157	dev (NA, NA)
Main RecrDev 2005	0.542	3	act	0.108269	dev (NA, NA)
Main RecrDev 2006	0.042	3	act	0.171393	dev (NA, NA)
Main RecrDev 2007	-0.278	3	act	0.219061	dev (NA, NA)
Main RecrDev 2008	0.981	3	act	0.0702548	dev (NA, NA)
Main RecrDev 2009	-0.362	3	act	0.194985	dev (NA, NA)

Table 18: List of parameters used in the base model, including estimated values and standard deviations (SD), bounds (minimum and maximum), estimation phase (negative values not estimated), status (indicates if parameters are near bounds), and prior type information (mean and SD). (*continued*)

Parameter	Value	Phase	Bounds	Status	Prior (Exp.Val, SD)
Main RecrDev 2010	0.498	3	act	0.0817707	dev (NA, NA)
Main RecrDev 2011	0.42	3	act	0.0810448	dev (NA, NA)
Main RecrDev 2012	0.212	3	act	0.090591	dev (NA, NA)
Main RecrDev 2013	0.564	3	act	0.0719959	dev (NA, NA)
Main RecrDev 2014	0.481	3	act	0.0802748	dev (NA, NA)
Main RecrDev 2015	-0.243	3	act	0.13553	dev (NA, NA)
Main RecrDev 2016	0.77	3	act	0.0927345	dev (NA, NA)
Main RecrDev 2017	-0.637	3	act	0.247401	dev (NA, NA)
Late RecrDev 2018	0	-	-	-	dev (NA, NA)
Late RecrDev 2019	0	-	-	-	dev (NA, NA)
Late RecrDev 2020	0	-	-	-	dev (NA, NA)
Late RecrDev 2021	0	-	-	-	dev (NA, NA)
Late RecrDev 2022	0	-	-	-	dev (NA, NA)
ForeRecr 2023	0	-	-	-	dev (NA, NA)
InitF seas 1 ft 1Trawl wdis	0	-1	-	-	None
InitF seas 1 ft 2Non-Trawl wdis	0	-1	-	-	None
InitF seas 1 ft 3Ocean	0	-1	-	-	None
InitF seas 1 ft 4Shore	0	-1	-	-	None
LnQ base Non-Trawl wdis(2)	-5.817	-1	-	-	None
Q extraSD Non-Trawl wdis(2)	0.028	3	LO	0.0147028	None
LnQ base Ocean(3)	-9.234	-1	-	-	None
Q extraSD Ocean(3)	0.284	3	OK	0.0516932	None
LnQ base MPA(5)	-7.391	-1	-	-	None
Q extraSD MPA(5)	0.167	3	OK	0.0841336	None
LnQ base Acoustic Visual(6)	0.6	-5	-	-	None
Q extraSD Acoustic Visual(6)	0	-3	-	-	None
LnQ base Tag(7)	-1.177	-1	-	-	None
Q extraSD Tag(7)	0.158	3	OK	0.0464933	None
LnQ base MRFSS(8)	-7.691	-1	-	-	None

Table 18: List of parameters used in the base model, including estimated values and standard deviations (SD), bounds (minimum and maximum), estimation phase (negative values not estimated), status (indicates if parameters are near bounds), and prior type information (mean and SD). (*continued*)

Parameter	Value	Phase	Bounds	Status	Prior (Exp.Val, SD)
Q extraSD MRFSS(8)	0.397	3	OK	0.0927459	None
Size DblN peak Trawl wdis(1)	52.532	4	OK	0.584074	None
Size DblN top logit Trawl wdis(1)	0.77	-4	-	-	None
Size DblN ascend se Trawl wdis(1)	3.585	-4	-	-	None
Size DblN descend se Trawl wdis(1)	-9.21	-4	-	-	None
Size DblN start logit Trawl wdis(1)	-10	-2	-	-	None
Size DblN end logit Trawl wdis(1)	11.513	-4	-	-	None
Size DblN peak Non-Trawl wdis(2)	42.629	4	OK	0.130805	None
Size DblN top logit Non-Trawl wdis(2)	-12	-4	-	-	None
Size DblN ascend se Non-Trawl wdis(2)	4	-4	-	-	None
Size DblN descend se Non-Trawl wdis(2)	2.16	4	OK	0.191113	None
Size DblN start logit Non-Trawl wdis(2)	-15	-2	-	-	None
Size DblN end logit Non-Trawl wdis(2)	-1.579	4	OK	0.21853	None
Size DblN peak Ocean(3)	41.838	4	OK	0.122527	None
Size DblN top logit Ocean(3)	-12	-4	-	-	None
Size DblN ascend se Ocean(3)	4	-4	-	-	None
Size DblN descend se Ocean(3)	2.69	4	OK	0.219738	None
Size DblN start logit Ocean(3)	-15	-2	-	-	None
Size DblN end logit Ocean(3)	-0.98	4	OK	0.231653	None
Size DblN peak Shore(4)	22.075	4	OK	1.58355	None
Size DblN top logit Shore(4)	4	-4	-	-	None
Size DblN ascend se Shore(4)	3	-4	-	-	None
Size DblN descend se Shore(4)	0	-4	-	-	None
Size DblN start logit Shore(4)	-15	-2	-	-	None
Size DblN end logit Shore(4)	4.342	-4	-	-	None
Size DblN peak MPA(5)	43.333	4	OK	0.52157	None
Size DblN top logit MPA(5)	-12	-4	-	-	None
Size DblN ascend se MPA(5)	4	-4	-	-	None
Size DblN descend se MPA(5)	1.228	4	OK	1.35908	None

Table 18: List of parameters used in the base model, including estimated values and standard deviations (SD), bounds (minimum and maximum), estimation phase (negative values not estimated), status (indicates if parameters are near bounds), and prior type information (mean and SD). (*continued*)

Parameter	Value	Phase	Bounds	Status	Prior (Exp.Val, SD)
Size DblN start logit MPA(5)	-15	-2	-	-	None
Size DblN end logit MPA(5)	-1.282	4	OK	0.63341	None
Size DblN peak Acoustic Visual(6)	25	-4	-	-	None
Size DblN top logit Acoustic Visual(6)	3.269	-4	-	-	None
Size DblN ascend se Acoustic Visual(6)	3.585	-4	-	-	None
Size DblN descend se Acoustic Visual(6)	-9.21	-4	-	-	None
Size DblN start logit Acoustic Visual(6)	-15	-2	-	-	None
Size DblN end logit Acoustic Visual(6)	11.513	-4	-	-	None
Size inflection Tag(7)	32	-3	-	-	None
Size 95%width Tag(7)	0	-4	-	-	None
SizeSel P1 MRFSS(8)	-1	-99	-	-	None
SizeSel P2 MRFSS(8)	-1	-99	-	-	None

Table 19: Likelihood components by source.

Label	Total
TOTAL	3687.49
Catch	0.00
Equil catch	0.00
Survey	-76.09
Length comp	492.51
Age comp	3253.30
Recruitment	-5.04
InitEQ Regime	0.00
Forecast Recruitment	0.00
Parm priors	22.81
Parm softbounds	0.00
Parm devs	0.00
Crash Pen	0.00

Table 20: Summary of reference points and management quantities, including estimates of the 95 percent intervals.

	Estimate	Lower Interval	Upper Interval
Unfished Spawning Output	1632.81	1577.19	1688.43
Unfished Age 0+ Biomass (mt)	11576.10	11142.20	12010.00
Unfished Recruitment (R0)	3807.86	3678.16	3937.56
Spawning Output (2023)	899.77	855.05	944.49
Fraction Unfished (2023)	0.55	0.53	0.57
Reference Points Based SB _{40%}			
Proxy Spawning Output SB40\%	653.12	630.88	675.37
SPR Resulting in SB40\%	0.46	0.46	0.46
Exploitation Rate Resulting in SB40\%	0.07	0.07	0.07
Reference Points Based on SPR Proxy for MSY			
Proxy Spawning Output (SPR50)	728.48	703.67	753.30
SPR50	0.50		
Exploitation Rate Corresponding to SPR50	0.06	0.06	0.07
Yield with SPR50 at SB SPR (mt)	454.96	438.30	471.63
Reference Points Based on Estimated MSY Values			
Spawning Output at MSY (SB MSY)	386.95	373.08	400.83
SPR MSY	0.31	0.31	0.31
Exploitation Rate Corresponding to SPR MSY	0.11	0.11	0.11
MSY (mt)	533.65	514.71	552.60

Table 21: Recent trend in the overfishing limits (OFL), the acceptable biological catches (ABCs), the annual catch limits (ACLs), and the total catch (mt). Management specifications prior to 2017 were for Oregon and California combined.

Year	OFL	ABC	ACL	Catch
2013	1159	1108	1000	442.14
2014	1166	1115	1000	479.45
2015	1176	1124	1000	594.52
2016	1183	1131	1000	524.53
2017	577	527	527	541.43
2018	570	520	520	415.07
2019	565	516	516	437.55
2020	561	512	512	435.56
2021	570	512	512	435.54
2022	566	512	512	526.00

Table 22: Time series of population estimates from the base model.

Year	Total Biomass (mt)	Spawning Output	Total Biomass 0+ (mt)	Frac-tion Un-fished	Age-0 Recruits	Total Mortal-ity (mt)	1-SPR	Ex-ploita-tion Rate
1892	11576.10	1632.81	11576.10	1.00	3807.88	0.50	0.00	0.00
1893	11575.60	1632.73	11575.60	1.00	3807.86	0.50	0.00	0.00
1894	11575.10	1632.66	11575.10	1.00	3807.85	0.50	0.00	0.00
1895	11574.70	1632.58	11574.70	1.00	3807.83	0.12	0.00	0.00
1896	11574.70	1632.57	11574.70	1.00	3807.83	0.04	0.00	0.00
1897	11574.80	1632.57	11574.80	1.00	3807.83	0.04	0.00	0.00
1898	11574.90	1632.58	11574.90	1.00	3807.83	0.02	0.00	0.00
1899	11575.00	1632.59	11575.00	1.00	3807.83	0.04	0.00	0.00
1900	11575.10	1632.60	11575.10	1.00	3807.83	0.04	0.00	0.00
1901	11575.20	1632.61	11575.20	1.00	3807.84	0.06	0.00	0.00
1902	11575.20	1632.62	11575.20	1.00	3807.84	0.06	0.00	0.00
1903	11575.30	1632.63	11575.30	1.00	3807.84	0.08	0.00	0.00
1904	11575.30	1632.64	11575.30	1.00	3807.84	0.10	0.00	0.00
1905	11575.30	1632.64	11575.30	1.00	3807.84	0.10	0.00	0.00
1906	11575.30	1632.64	11575.30	1.00	3807.84	0.12	0.00	0.00
1907	11575.20	1632.64	11575.20	1.00	3807.84	0.12	0.00	0.00
1908	11575.20	1632.64	11575.20	1.00	3807.84	0.14	0.00	0.00
1909	11575.20	1632.63	11575.20	1.00	3807.84	0.16	0.00	0.00
1910	11575.10	1632.62	11575.10	1.00	3807.84	0.16	0.00	0.00
1911	11575.10	1632.61	11575.10	1.00	3807.84	0.18	0.00	0.00
1912	11575.00	1632.60	11575.00	1.00	3807.83	0.20	0.00	0.00
1913	11574.90	1632.58	11574.90	1.00	3807.83	0.20	0.00	0.00
1914	11574.90	1632.57	11574.90	1.00	3807.83	0.22	0.00	0.00
1915	11574.80	1632.55	11574.80	1.00	3807.82	0.22	0.00	0.00
1916	11574.70	1632.54	11574.70	1.00	3807.82	0.24	0.00	0.00
1917	11574.60	1632.52	11574.60	1.00	3807.82	0.26	0.00	0.00
1918	11574.50	1632.50	11574.50	1.00	3807.81	0.26	0.00	0.00
1919	11574.40	1632.49	11574.40	1.00	3807.81	0.28	0.00	0.00
1920	11574.30	1632.47	11574.30	1.00	3807.80	0.30	0.00	0.00
1921	11574.20	1632.45	11574.20	1.00	3807.80	0.30	0.00	0.00
1922	11574.10	1632.43	11574.10	1.00	3807.79	0.32	0.00	0.00
1923	11574.00	1632.40	11574.00	1.00	3807.79	0.32	0.00	0.00
1924	11573.90	1632.39	11573.90	1.00	3807.79	0.34	0.00	0.00
1925	11573.80	1632.36	11573.80	1.00	3807.78	0.36	0.00	0.00
1926	11573.70	1632.34	11573.70	1.00	3807.78	0.36	0.00	0.00
1927	11573.60	1632.32	11573.60	1.00	3807.77	0.38	0.00	0.00
1928	11573.50	1632.30	11573.50	1.00	3807.77	0.66	0.00	0.00
1929	11573.10	1632.24	11573.10	1.00	3807.75	1.77	0.00	0.00

Table 22: Time series of population estimates from the base model. (*continued*)

Year	Total Biomass (mt)	Spawning Output	Total Biomass 0+ (mt)	Frac-tion Un-fished	Age-0 Recruits	Total Mortal-ity (mt)	1-SPR	Ex-ploita-tion Rate
1930	11571.70	1632.01	11571.70	1.00	3807.70	2.25	0.00	0.00
1931	11570.00	1631.71	11570.00	1.00	3807.63	1.34	0.00	0.00
1932	11569.20	1631.56	11569.20	1.00	3807.60	0.56	0.00	0.00
1933	11569.40	1631.51	11569.40	1.00	3807.59	0.69	0.00	0.00
1934	11569.50	1631.46	11569.50	1.00	3807.58	0.65	0.00	0.00
1935	11569.50	1631.46	11569.50	1.00	3807.58	0.56	0.00	0.00
1936	11569.70	1631.47	11569.70	1.00	3807.58	2.19	0.00	0.00
1937	11568.40	1631.20	11568.40	1.00	3807.52	4.44	0.00	0.00
1938	11565.20	1630.56	11565.20	1.00	3807.37	3.55	0.00	0.00
1939	11562.90	1630.19	11562.90	1.00	3807.29	5.29	0.01	0.00
1940	11559.40	1629.42	11559.40	1.00	3807.11	6.52	0.01	0.00
1941	11555.00	1628.52	11555.00	1.00	3806.91	7.69	0.01	0.00
1942	11550.50	1627.29	11550.50	1.00	3806.63	11.15	0.01	0.00
1943	11543.60	1625.48	11543.60	1.00	3806.21	40.40	0.04	0.00
1944	11515.40	1616.66	11515.40	0.99	3804.19	83.62	0.08	0.01
1945	11458.80	1598.03	11458.80	0.98	3799.84	134.52	0.13	0.01
1946	11372.00	1569.04	11372.00	0.96	3792.90	96.74	0.10	0.01
1947	11325.30	1553.04	11325.30	0.95	3788.96	32.60	0.03	0.00
1948	11333.30	1554.68	11333.30	0.95	3789.37	23.22	0.02	0.00
1949	11346.30	1559.18	11346.30	0.95	3790.48	9.21	0.01	0.00
1950	11367.70	1566.71	11367.70	0.96	3792.33	60.30	0.06	0.01
1951	11347.50	1560.91	11347.50	0.96	3790.91	17.37	0.02	0.00
1952	11361.70	1566.25	11361.70	0.96	3792.22	9.03	0.01	0.00
1953	11379.60	1573.14	11379.60	0.96	3793.89	20.99	0.02	0.00
1954	11386.30	1575.96	11386.30	0.97	3794.58	26.57	0.03	0.00
1955	11387.90	1576.88	11387.90	0.97	3794.80	36.51	0.04	0.00
1956	11381.60	1575.22	11381.60	0.96	3794.40	24.83	0.03	0.00
1957	11385.20	1576.39	11385.20	0.97	3794.68	25.54	0.03	0.00
1958	11387.60	1577.33	11387.60	0.97	3794.91	34.81	0.04	0.00
1959	11383.30	1575.64	11383.30	0.96	3794.50	43.21	0.05	0.00
1960	11373.40	1572.13	11373.40	0.96	3793.65	76.53	0.08	0.01
1961	11340.20	1560.89	11340.20	0.96	3790.90	63.86	0.07	0.01
1962	11320.80	1554.21	11320.80	0.95	3789.25	60.89	0.06	0.01
1963	11306.40	1549.10	11306.40	0.95	3787.98	48.46	0.05	0.00
1964	11303.10	1547.84	11303.10	0.95	3787.67	29.44	0.03	0.00
1965	11314.40	1551.43	11314.40	0.95	3788.56	54.15	0.06	0.00
1966	11304.70	1549.07	11304.70	0.95	3787.98	47.57	0.05	0.00
1967	11301.10	1548.42	11301.10	0.95	3787.81	35.21	0.04	0.00
1968	11305.80	1551.35	11305.80	0.95	3788.54	31.92	0.03	0.00

Table 22: Time series of population estimates from the base model. (*continued*)

Year	Total Biomass (mt)	Spawning Output	Total Biomass 0+ (mt)	Frac-tion Un-fished	Age-0 Re-cruits	Total Mortal-ity (mt)	1-SPR	Ex-ploita-tion Rate
1969	11311.90	1554.59	11311.90	0.95	3789.35	65.98	0.07	0.01
1970	11289.70	1549.69	11289.70	0.95	3788.13	37.46	0.04	0.00
1971	11292.90	1551.25	11292.90	0.95	3788.52	48.63	0.05	0.00
1972	11285.50	1550.65	11285.50	0.95	3788.37	64.39	0.07	0.01
1973	11265.90	1546.49	11265.90	0.95	3787.33	110.57	0.11	0.01
1974	11203.90	1535.73	11203.90	0.94	3784.62	150.28	0.14	0.01
1975	11107.70	1520.25	11107.70	0.93	3780.67	188.54	0.17	0.02
1976	10985.80	1498.52	10985.80	0.92	3775.00	220.38	0.20	0.02
1977	10842.90	1474.44	10842.90	0.90	3768.53	275.00	0.24	0.03
1978	10664.00	1442.31	10664.00	0.88	3759.61	336.05	0.29	0.03
1979	10447.50	1401.85	10447.50	0.86	3747.86	448.78	0.37	0.04
1980	10144.00	1344.83	10144.00	0.82	1136.21	305.45	0.28	0.03
1981	9980.97	1314.68	9980.97	0.81	1391.61	476.56	0.40	0.05
1982	9569.45	1260.50	9569.45	0.77	2360.32	651.36	0.50	0.07
1983	8897.34	1181.56	8897.34	0.72	2042.78	499.03	0.43	0.06
1984	8319.12	1134.26	8319.12	0.69	2366.24	533.38	0.47	0.06
1985	7695.70	1081.97	7695.70	0.66	2644.81	341.97	0.36	0.04
1986	7277.77	1052.78	7277.77	0.64	2187.26	308.71	0.35	0.04
1987	6931.25	1014.09	6931.25	0.62	2092.26	357.80	0.41	0.05
1988	6575.01	952.93	6575.01	0.58	2234.28	394.27	0.47	0.06
1989	6220.77	879.58	6220.77	0.54	2926.60	457.76	0.53	0.07
1990	5850.59	797.26	5850.59	0.49	2750.62	501.85	0.59	0.09
1991	5513.12	716.94	5513.12	0.44	3091.13	342.01	0.49	0.06
1992	5406.08	671.89	5406.08	0.41	2844.93	640.06	0.70	0.12
1993	5090.17	588.17	5090.17	0.36	2806.16	543.18	0.68	0.11
1994	4942.87	526.66	4942.87	0.32	3766.99	485.55	0.66	0.10
1995	4912.90	479.44	4912.90	0.29	3530.84	445.48	0.64	0.09
1996	4978.85	455.70	4978.85	0.28	2633.33	519.31	0.68	0.10
1997	5026.01	434.24	5026.01	0.27	2476.34	550.38	0.70	0.11
1998	5050.84	417.34	5050.84	0.26	1858.17	467.09	0.65	0.09
1999	5129.77	415.69	5129.77	0.25	4705.78	287.60	0.48	0.06
2000	5326.30	445.02	5326.30	0.27	5308.72	336.23	0.50	0.06
2001	5509.06	477.63	5509.06	0.29	3546.34	457.42	0.58	0.08
2002	5671.01	498.00	5671.01	0.30	1682.87	439.51	0.57	0.08
2003	5899.69	513.74	5899.69	0.31	4562.39	457.69	0.58	0.08
2004	6062.13	517.19	6062.13	0.32	963.31	450.17	0.57	0.07
2005	6212.95	520.93	6212.95	0.32	4576.62	411.49	0.53	0.07
2006	6310.86	544.30	6310.86	0.33	2807.51	368.35	0.47	0.06
2007	6419.57	587.37	6419.57	0.36	2075.32	362.88	0.45	0.06

Table 22: Time series of population estimates from the base model. (*continued*)

Year	Total Biomass (mt)	Spawning Output	Total Biomass 0+ (mt)	Frac-tion Un-fished	Age-0 Recruits	Total Mortal-ity (mt)	1-SPR	Ex-ploita-tion Rate
2008	6510.66	631.84	6510.66	0.39	7429.24	340.32	0.42	0.05
2009	6589.38	669.47	6589.38	0.41	1964.00	425.59	0.49	0.06
2010	6698.85	682.82	6698.85	0.42	4659.64	406.89	0.48	0.06
2011	6861.40	688.64	6861.40	0.42	4316.21	310.30	0.40	0.05
2012	7150.45	704.49	7150.45	0.43	3523.19	320.90	0.40	0.04
2013	7452.95	719.26	7452.95	0.44	5030.30	442.14	0.49	0.06
2014	7637.87	726.10	7637.87	0.44	4638.54	479.45	0.50	0.06
2015	7808.00	740.31	7808.00	0.45	2255.93	594.52	0.56	0.08
2016	7891.11	744.99	7891.11	0.46	6224.71	524.53	0.52	0.07
2017	7988.87	759.97	7988.87	0.47	1595.64	541.43	0.52	0.07
2018	8080.23	773.49	8080.23	0.47	3436.75	415.07	0.43	0.05
2019	8210.70	805.69	8210.70	0.49	3462.37	437.55	0.44	0.05
2020	8236.88	838.15	8236.88	0.51	3486.56	435.56	0.43	0.05
2021	8206.35	868.88	8206.35	0.53	3508.07	435.54	0.42	0.05
2022	8142.78	894.62	8142.78	0.55	3525.14	526.00	0.48	0.06
2023	7981.84	899.77	7981.84	0.55	3528.45	496.49	0.47	0.06

Table 23: Likelihood, parameter and derived quantities from data (catchability) treatment sensitivities.

	Reference Model	AV q=1, small CV	Est AV q, small CV	AV q=1, CV =0.45	Est AV q, CV=0.45	Tag q 2015 value	Tag q 2015 value, xvar
AIC	7493	7601	7443	7460	7456	7664	7466
deltaAIC	0	108	-50	-33	-37	171	-27
<u>Survey likelihood</u>							
Total	-76	-64	-81	-72	-75	-28	-70
Non-trawl	-38	-32	-36	-37	-36	-30	-37
Ocean	-14	-10	-19	-18	-19	-23	-18
AV	-7	-7	-7	2	-1	-7	-7
Tag	-10	-9	-11	-11	-11	44	0
MRFSS	-2	-3	-2	-2	-2	-7	-2
MPA	-5	-4	-6	-6	-6	-6	-6
<u>Length likelihood</u>							
Total	493	515	471	472	471	521	473
Trawl	72	70	73	73	73	69	73
Non-trawl	286	297	272	272	272	306	273
Ocean	92	98	87	87	87	102	87
Shore	27	31	23	23	23	26	23
MPA	17	18	16	16	16	18	16
<u>Age likelihood</u>							
Total	3253	3255	3267	3266	3267	3250	3265
Trawl	669	675	664	664	664	670	664
Non-trawl	1330	1330	1335	1335	1335	1333	1334
Ocean	1255	1250	1269	1268	1269	1247	1267
<u>Parameters</u>							
Female M	0.19	0.19	0.19	0.19	0.19	0.19	0.19
Female L_∞	50.85	51.36	50.29	50.31	50.29	51.38	50.35
Female k	0.18	0.18	0.18	0.18	0.18	0.18	0.18
Male M	0.17	0.17	0.17	0.17	0.17	0.17	0.17
Male L_∞	44.88	45.32	44.48	44.49	44.48	45.38	44.52
Male k	0.22	0.22	0.23	0.23	0.23	0.22	0.23

Table 23: Likelihood, parameter and derived quantities from data (catchability) treatment sensitivities. (*continued*)

	Reference Model	AV q=1, small CV	Est AV q, small CV	AV q=1, CV =0.45	Est AV q, CV=0.45	Tag q 2015 value	Tag q 2015 value, xvar
$\ln(R_0)$	8.24	8.50	8.08	8.09	8.08	8.25	8.09
Non-trawl logQ	-5.82	-6.35	-5.33	-5.36	-5.33	-5.90	-5.38
Non-trawl extra SD	0.03	0.06	0.04	0.03	0.04	0.07	0.03
Ocean logQ	-9.23	-9.74	-8.79	-8.81	-8.79	-9.33	-8.84
Ocean extra SD	0.28	0.35	0.21	0.22	0.21	0.17	0.22
MPA logQ	-7.39	-7.99	-6.86	-6.88	-6.86	-7.44	-6.91
MPA extra SD	0.17	0.20	0.12	0.13	0.12	0.12	0.12
AV logQ	0.60	0.00	1.17	0.00	1.17	0.73	1.12
Tag logQ	-1.18	-1.66	-0.79	-0.80	-0.79	-1.39	-1.39
Tag extra SD	0.16	0.18	0.14	0.14	0.14	0.00	0.55
MRFSS logQ	-7.69	-8.00	-7.48	-7.48	-7.48	-7.81	-7.49
MRFSS extra SD	0.40	0.39	0.41	0.41	0.41	0.28	0.40
Trawl: Peak	52.53	51.90	53.23	53.21	53.23	51.80	53.16
Non-trawl: Peak	42.63	42.27	43.12	43.10	43.12	42.16	43.07
Non-trawl: Decend	2.16	2.27	1.76	1.78	1.76	2.31	1.82
Non-trawl: End	-1.58	-1.85	-1.10	-1.12	-1.10	-1.92	-1.16
Ocean: Peak	41.84	41.51	42.23	42.22	42.23	41.39	42.19
Ocean: Decend	2.69	2.73	2.62	2.62	2.62	2.75	2.63
Ocean: End	-0.98	-1.29	-0.56	-0.57	-0.56	-1.38	-0.60
Shore: Peak	22.07	21.66	22.37	22.35	22.37	22.26	22.35
MPA: Peak	43.33	42.82	43.86	43.85	43.86	42.79	43.80
MPA: Decend	1.23	1.92	0.42	0.44	0.42	1.96	0.53
MPA: End	-1.28	-1.75	-0.91	-0.92	-0.91	-1.84	-0.95
<u>Derived quantities</u>							
$S0_0$	1632.81	2194.04	1312.83	1326.79	1312.83	1730.45	1336.98
$S0_{2023}$	899.77	1867.42	414.19	436.60	414.19	844.18	448.42
$Bratio_{2023}$	0.55	0.85	0.32	0.33	0.32	0.49	0.34
MSY_{SPR}	454.96	591.99	379.58	383.20	379.58	465.78	385.05
F_{SPR}	0.06	0.07	0.06	0.06	0.06	0.07	0.06

Table 24: Likelihood, parameter and derived quantities from other data treatment sensitivities.

Table 24: Likelihood, parameter and derived quantities from other data treatment sensitivities. (*continued*)

	Reference Model	No AV	No tag	No non-trawl	No ORBS	No MRFSS	No MPA	Add SMURF	No data wts	Dirichlet wts	McAll wts	2015 catches
Male L_∞	44.88	44.48	44.87	44.57	44.78	44.80	44.90	44.88	45.96	45.97	45.75	44.90
Male k	0.22	0.23	0.22	0.23	0.22	0.22	0.22	0.22	0.21	0.21	0.22	0.22
$\ln(R_0)$	8.24	8.08	8.24	8.23	8.24	8.25	8.24	8.25	8.24	8.24	8.25	8.22
Non-trawl logQ	-5.82	-5.33	-5.81		-5.79	-5.80	-5.81	-5.82	-5.91	-5.91	-5.87	-5.83
Non-trawl extra SD	0.03	0.04	0.03		0.04	0.03	0.03	0.03	0.03	0.03	0.02	0.03
Ocean logQ	-9.23	-8.79	-9.23	-9.12		-9.22	-9.23	-9.23	-9.32	-9.32	-9.29	-9.25
Ocean extra SD	0.28	0.21	0.29	0.33		0.29	0.28	0.28	0.23	0.23	0.26	0.28
MPA logQ	-7.39	-6.86	-7.39	-7.28	-7.36	-7.37		-7.39	-7.48	-7.49	-7.44	-7.40
MPA extra SD	0.17	0.12	0.17	0.19	0.17	0.17		0.17	0.14	0.14	0.16	0.17
AV logQ	0.60		0.60	0.60	0.60	0.60	0.60	0.60	0.60	0.60	0.60	0.60
Tag logQ	-1.18	-0.79		-1.04	-1.14	-1.16	-1.17	-1.18	-1.36	-1.36	-1.24	-1.19
Tag extra SD	0.16	0.14		0.17	0.16	0.16	0.16	0.16	0.15	0.15	0.15	0.15
MRFSS logQ	-7.69	-7.48	-7.69	-7.63	-7.67		-7.69	-7.69	-7.75	-7.76	-7.78	-7.73
MRFSS extra SD	0.40	0.41	0.40	0.44	0.41		0.40	0.40	0.30	0.30	0.34	0.38
Trawl: Peak	52.53	53.23	52.55	52.95	52.66	52.62	52.52	52.53	51.31	51.28	52.05	52.46
Non-trawl: Peak	42.63	43.12	42.64	42.96	42.72	42.68	42.63	42.63	42.20	42.20	42.43	42.61
Non-trawl: Decend	2.16	1.76	2.16	1.93	2.12	2.14	2.16	2.16	2.36	2.36	2.12	2.17
Non-trawl: End	-1.58	-1.10	-1.57	-1.31	-1.51	-1.54	-1.58	-1.58	-2.26	-2.26	-1.56	-1.60
Ocean: Peak	41.84	42.23	41.85	42.10	41.91	41.89	41.84	41.84	41.46	41.45	41.65	41.82
Ocean: Decend	2.69	2.62	2.69	2.65	2.68	2.68	2.69	2.69	2.82	2.82	2.60	2.69
Ocean: End	-0.98	-0.56	-0.97	-0.74	-0.91	-0.93	-0.98	-0.98	-1.71	-1.71	-1.07	-0.99
Shore: Peak	22.07	22.37	22.07	22.04	22.06	22.02	22.08	22.07	22.45	22.30	22.00	21.95
MPA: Peak	43.33	43.86	43.33	43.68	43.43	43.39		43.33	42.78	42.77	43.13	43.31
MPA: Decend	1.23	0.42	1.23	0.71	1.09	1.14		1.23	2.05	2.03	1.44	1.26
MPA: End	-1.28	-0.91	-1.28	-1.05	-1.21	-1.24		-1.28	-2.16	-2.11	-1.42	-1.30
Derived quantities												
$S0_0$	1632.81	1312.83	1630.02	1553.14	1607.44	1623.37	1632.88	1633.86	1904.21	1910.06	1628.47	1600.47
$S0_{2023}$	899.77	414.19	898.95	855.44	887.58	890.12	899.31	898.66	1082.33	1083.56	904.36	904.77
$Bratio_{2023}$	0.55	0.32	0.55	0.55	0.55	0.55	0.55	0.55	0.57	0.57	0.56	0.57

Table 24: Likelihood, parameter and derived quantities from other data treatment sensitivities. (*continued*)

	Reference Model	No AV	No tag	No non-trawl	No ORBS	No MRFSS	No MPA	Add SMURF	No data wts	Dirichlet wts	McAll wts	2015 catches
MSY_{SPR}	454.96	379.58	454.43	442.29	450.80	454.35	454.72	455.27	460.43	461.52	461.95	445.10
F_{SPR}	0.06	0.06	0.06	0.06	0.06	0.06	0.06	0.06	0.07	0.07	0.07	0.06

Table 25: Likelihood, parameter and derived quantities from alternative model specification sensitivities.

Table 25: Likelihood, parameter and derived quantities from alternataive model specification sensitivities. (*continued*)

	Refer- ence Model	2015 M	Loren- zen M	Est M	Fix growth, est M	Fix growth	Est Lmin	Est CVlts	Bio mat	Fx- nal ogive	Fec = mat	No rec devs	Full rec devs	Logis- tic sel
Female Lmin	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Female L_∞	50.85	50.90	50.56	50.41	48.00	48.00	50.85	50.85	50.81	50.84	50.83	52.22	50.83	48.18
Female k	0.18	0.18	0.18	0.18	0.18	0.18	0.18	0.18	0.18	0.18	0.18	0.18	0.18	0.20
Female CV young	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10
Female CV old	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10
Female mat50	40.36	40.36	40.36	40.36	40.36	40.36	40.36	40.36	34.38	40.36	40.36	40.36	40.36	40.36
Female neg. slope	0.38	0.38	0.38	0.38	0.38	0.38	0.38	0.38	0.31	0.38	0.38	0.38	0.38	0.38
Male M	0.17			0.18	0.11	0.17	0.17	0.17	0.17	0.17	0.17	0.17	0.17	0.17
Male M break 1		0.17												
Male M break 2		0.17												
Male M Lorenz			0.17											
Male Lmin	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Male L_∞	44.88	45.08	44.34	44.30	44.87	44.87	44.88	44.88	44.84	44.87	44.86	46.55	44.89	42.93
Male k	0.22	0.22	0.23	0.23	0.21	0.21	0.22	0.22	0.22	0.22	0.22	0.21	0.22	0.24
Male CV young	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10
Male CV old	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10
$\ln(R_0)$	8.24	8.17	9.85	8.58	7.53	8.16	8.24	8.24	8.22	8.24	8.23	8.29	8.08	8.21
Non-trawl logQ	-5.8	-5.9	-5.7	-5.8	-6.2	-6.4	13.2	-5.8	-5.8	-5.8	-5.8	-6.0	-5.8	-5.5
Non-trawl extra SD	0.03	0.02	0.04	0.02	0.12	0.25	0.03	0.03	0.03	0.03	0.03	0.09	0.03	0.16
Ocean logQ	-9.2	-9.3	-9.1	-9.3	-9.7	-9.9	9.4	-9.2	-9.2	-9.2	-9.2	-9.4	-9.2	-8.8
Ocean extra SD	0.28	0.28	0.30	0.26	0.14	0.13	0.28	0.28	0.29	0.28	0.29	0.15	0.29	0.42
MPA logQ	-7.4	-7.4	-7.3	-7.4	-7.8	-7.9	12.4	-7.4	-7.4	-7.4	-7.4	-7.5	-7.4	-7.0
MPA extra SD	0.17	0.16	0.18	0.15	0.09	0.05	0.17	0.17	0.17	0.17	0.17	0.10	0.17	0.26
Tag logQ	-1.2	-1.2	-1.1	-1.2	-1.7	-1.9	19.4	-1.2	-1.2	-1.2	-1.2	-1.5	-1.2	-0.7
Tag extra SD	0.16	0.16	0.16	0.14	0.11	0.08	0.16	0.16	0.16	0.16	0.16	0.13	0.16	0.17
MRFSS logQ	-7.7	-7.7	-7.6	-7.7	-8.0	-8.4	-0.5	-7.7	-7.7	-7.7	-7.7	-8.1	-7.2	-7.5
MRFSS extra SD	0.40	0.38	0.44	0.36	0.11	0.06	0.40	0.40	0.39	0.40	0.39	0.20	0.08	0.59
Trawl: Peak	52.5	52.4	53.1	53.3	52.3	52.6	52.5	52.5	52.6	52.5	52.6	50.9	52.7	55.0

Table 25: Likelihood, parameter and derived quantities from alternataive model specification sensitivities. (*continued*)

	Refer- ence Model	2015 M	Loren- zen M	Est M	Fix growth, est M	Fix growth	Est Lmin	Est CVlts	Bio mat	Fx- nal ogive	Fec = mat	No rec devs	Full rec devs	Logis- tic sel
Non-trawl: Peak	42.6	42.5	43.2	43.2	41.5	41.8	42.6	42.6	42.7	42.6	42.6	41.9	42.6	42.9
Non-trawl: Ascend	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0	3.8
Non-trawl: Decend	2.2	2.2	1.5	1.6	2.3	2.2	2.2	2.2	2.2	2.2	2.2	2.3	2.2	-9.0
Non-trawl: End	-1.6	-1.6	-1.1	-1.0	-1.4	-1.4	-1.6	-1.6	-1.6	-1.6	-1.6	-2.2	-1.6	10.0
Ocean: Peak	41.8	41.7	42.4	42.3	40.6	41.0	41.8	41.8	41.9	41.8	41.9	41.3	41.9	43.0
Ocean: Ascend	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0
Ocean: Decend	2.7	2.7	2.5	2.6	2.7	2.5	2.7	2.7	2.7	2.7	2.7	2.7	2.7	-3.1
Ocean: End	-1.0	-1.1	-0.3	-0.3	-0.8	-0.8	-1.0	-1.0	-1.0	-1.0	-1.0	-1.7	-1.0	10.0
Shore: Peak	22.1	22.0	23.1	22.7	14.7	15.0	22.1	22.1	22.1	22.1	22.1	22.6	23.5	21.7
MPA: Peak	43.3	43.2	44.0	43.8	41.9	42.3	43.3	43.3	43.4	43.3	43.3	42.3	43.3	44.6
MPA: Decend	1.2	1.5	0.3	0.5	2.1	2.1	1.2	1.2	1.2	1.2	1.2	2.2	1.2	1.6
MPA: End	-1.3	-1.4	-0.8	-0.7	-1.4	-1.4	-1.3	-1.3	-1.3	-1.3	-1.3	-2.1	-1.3	-0.6
Derived quantities														
$S0_0$	1633	1773	1112	926	1244	1484	8597	1633	2094	1767	2865	1897	1385	1253
$S0_{2023}$	900	997	579	554	723	734	507	900	1273	984	1676	986	901	660
$Bratio_{2023}$	0.55	0.56	0.52	0.60	0.58	0.49	0.06	0.55	0.61	0.56	0.59	0.52	0.65	0.53
MSY_{SPR}	455	466	432	517	397	529	338	455	492	459	473	489	387	419
F_{SPR}	0.06	0.07	0.06	0.07	0.06	0.06	0.02	0.06	0.08	0.07	0.07	0.07	0.06	0.06

Table 26: The magnitude of retrospective pattern (Mohn's rho; Mohn, 1999) given the removal of ten years of data for fishing intensity (F), fraction unfished (Figure 64), recruitment, and spawning stock biomass (SSB ; Figure 63). Columns are a derivation of Mohn's rho (Hurtado-Ferro et al. 2015) used by the Alaska Fisheries Science Center (AFSC), as originally described in Mohn (1999), and a derivation of Mohn's rho (Woods Hole Mohn's rho; Legault 2009) used by the Northeast Fisheries Science Center (NEFSC).

Quantity	AFSC	Mohn	NEFSC
F	1.733824	17.33824	4.591969
Fraction unfished	-0.4494829	-4.494829	-0.4856366
Recruitment	-0.5536441	-5.536441	-2.727355
SSB	-0.6081382	-6.081382	-3.459641

Table 27: Projections of potential OFLs (mt), ABCs (mt), the buffer (ABC = buffer x OFL), estimated total biomass, spawning output, and fraction unfished.

Year	Predicted OFL	ABC Catch	Buffer	Summary Biomass	Spawning Output	Fraction Unfished
2023	541.9	496.5	1.00	7981.8	899.8	0.55
2024	-	-	-	-	-	-
2025	-	-	-	-	-	-
2026	-	-	-	-	-	-
2027	-	-	-	-	-	-
2028	-	-	-	-	-	-
2029	-	-	-	-	-	-
2030	-	-	-	-	-	-
2031	-	-	-	-	-	-
2032	-	-	-	-	-	-
2033	-	-	-	-	-	-
2034	-	-	-	-	-	-

Table 28: Decision table summary of 10 year projections beginning in 2023 for alternative states of nature based on an axis of uncertainty related to model structure relative to the reference model (i.e., estimate catchability, Q, associated with the acoustic-visual survey and no estimation of recruitment deviations) . Columns range over low (12.5 quantile), mid (reference model), and high states (87.5 quantile) of nature and rows range over different catch level assumptions. Values in italics indicate years where the stock size prevented the full catch removals.

Year	Catch	Estimate Q		Reference Model		No Rec Devs	
		Spawning Output	Fraction Unfished	Spawning Output	Fraction Unfished	Spawning Output	Fraction Unfished
2023	-	-	-	-	-	-	-
2024	-	-	-	-	-	-	-
2025	-	-	-	-	-	-	-
2026	-	-	-	-	-	-	-
2027	-	-	-	-	-	-	-
P*=0.45	2028	-	-	-	-	-	-
sigma=0.5	2029	-	-	-	-	-	-
	2030	-	-	-	-	-	-
	2031	-	-	-	-	-	-
	2032	-	-	-	-	-	-
	2033	-	-	-	-	-	-
	2034	-	-	-	-	-	-
	2023	-	-	-	-	-	-
	2024	-	-	-	-	-	-
	2025	-	-	-	-	-	-
	2026	-	-	-	-	-	-
	2027	-	-	-	-	-	-
P*=0.4	2028	-	-	-	-	-	-
sigma=0.5	2029	-	-	-	-	-	-
	2030	-	-	-	-	-	-
	2031	-	-	-	-	-	-
	2032	-	-	-	-	-	-
	2033	-	-	-	-	-	-
	2034	-	-	-	-	-	-
	2023	-	-	-	-	-	-
	2024	-	-	-	-	-	-
	2025	-	-	-	-	-	-
	2026	-	-	-	-	-	-
Equilibrium yield from FMSY proxy of SPR=0.5	2027	-	-	-	-	-	-
	2028	-	-	-	-	-	-
	2029	-	-	-	-	-	-
	2030	-	-	-	-	-	-
	2031	-	-	-	-	-	-
	2032	-	-	-	-	-	-
	2033	-	-	-	-	-	-
	2034	-	-	-	-	-	-

8 Figures




Figure 1: Map of the management and assessment areas for black rockfish.




Figure 2: Summary of data sources used in the reference model.




Figure 3: Landings by fleet used in the reference model where catches in metric tons by fleet are stacked.




Figure 4: Observed length-at-age by data source and sex.




Figure 5: Observed length-at-age by sex and year. Total samples are indicated in parentheses.




Figure 6: External fits to the observed length-at-age by sex.




Figure 7: Coefficient of variation of length by age by sex. Numbers indicate samples by age and colors indicate sex.




Figure 8: Model estimated length-at-age. Shaded area indicates 95 percent distribution of length-at-age around the estimated growth curve.




Figure 9: Ageing bias plots by reader comparisons.




Figure 10: Estimated bias relationships for each considered matrix. Reader 1 is always considered unbiased. Reader 1a and 1b is an intra-reader comparison. B refers to the bias type and S refers to the imprecision type in the model selection for the ageing error matrix.




Figure 11: Ageing error matrix standard deviation (SD) values by comparison. B refers to the bias type and S refers to the imprecision type in the model selection for the ageing error matrix.




Figure 12: Biological and functional maturity ogives. Biological maturity ogives are represented by the dashed lines (logistic regression and 95% CI is grey, flexible spline and 95% CI is light green). Functional maturity ogives are represented by the solid lines (logistic regression and 95% CI is grey, flexible spline and 95% CI is light blue).




Figure 13: Functional maturity ogives and sample size. Functional maturity ogives are represented by the solid lines (logistic regression and 95% CI is grey, flexible spline and 95% CI is light blue) and sample size is denoted by the size of the bubbles).




Figure 14: Maturity as a function of length (cm).




Figure 15: Fecundity (kg) as a function of length (cm).




Figure 16: Sex-specific length (cm)-weight (kg) data for Oregon black rockfish samples by source. MRFSS and ORBS are the ocean boat recreational fishery from early and late periods.





Figure 17: Sex-specific length (cm)-weight (kg) estimated power function relationships. Washington state estimate relationships are also provided for comparison.






Figure 18: Comparison of spawning output for black rockfish in waters off of Oregon between Stock Synthesis versions 3.24 and 3.30. Uncertainty envelops are 95% confidence intervals.




Figure 19: Comparison of spawning output for black rockfish in waters off of Oregon between Stock Synthesis versions 3.24 and 3.30. Uncertainty envelops are 95% confidence intervals.




Figure 20: Relative spawning output time series by data treatment compared to the reference model. Missing scenarios mean the spawning output was too large to show.




Figure 21: Spawning output (in millions of eggs) time series by data treatment compared to the reference model.




Figure 22: Jitter runs for the black rockfish reference model, with jitter run number on the x-axis and -log likelihood value on the y-axis. Blue dot are models that match the likelihood value of the reference model, while red dots deviate from the reference model. All red dots are above the blue dots, indicating no better fit to the reference model was found.




Figure 23: Pairs plots of the fastest mixing parameters from running 2000 posterior draws (and keep every draw) using the random walk Metropolis algorithm. Parameters that show little to no movement are recommended to be fixed to improve model speed and efficiency.




Figure 24: Pearson residuals for each fishing fleet and the MPA survey. Closed bubble are positive residuals (observed > expected) and open bubbles are negative residuals (observed < expected).




Figure 25: Mean length (cm) index from the commercial trawl fishery with 95 percent confidence intervals based on sample sizes and data weighting.




Figure 26: Mean length (cm) index from the commercial non-trawl fishery with 95 percent confidence intervals based on sample sizes and data weighting.




Figure 27: Mean length (cm) index from the recreational ocean boat fishery with 95 percent confidence intervals based on sample sizes and data weighting.




Figure 28: Mean length (cm) index from the recreational shore-based fishery with 95 percent confidence intervals based on sample sizes and data weighting.




Figure 29: Mean length (cm) index from the MPA survey with 95 percent confidence intervals based on sample sizes and data weighting.




Figure 30: Aggregated length (cm) compositions over all years.




Figure 31: Mean age from conditional age-at-length data for the commercial trawl fishery.




Figure 32: Mean age observations from the conditional age-at-length data from the non-trawl commercial fishery.




Figure 33: Mean age observations from the conditional age-at-length data from the ocean boat fishery.




Figure 34: Fit to the non-trawl commercial survey index of abundance.




Figure 35: Fit to the MRFSS recreational survey index of abundance.




Figure 36: Fit to the MRFSS recreational survey index of abundance.




Figure 37: Fit to the tagging survey index of abundance.




Figure 38: Fit to the MPA survey index of abundance.




Figure 39: Model estimated length-at-age in the beginning of the year. Shaded area indicates 95 percent distribution of length-at-age around the estimated growth curve.




Figure 40: Estimated time series of age-0 recruits (1000s).




Figure 41: Estimated time series of recruitment deviations.




Figure 42: Recruitment deviations variance by year. This plot tracks the information content contained in each recruitment deviation. Values below the red line (assumed recruitment variability) indicates years with more informed recruitment deviations.




Figure 43: Stock-recruit curve. Point colors indicate year, with warmer colors indicating earlier years and cooler colors indicating later years.




Figure 44: Recruitment bias adjustment applied in the reference model.




Figure 45: Recruitment deviations variance check. Low standard deviations indicate years with informative deviations .




Figure 46: Length-based selectivity curves for each fleet and survey.




Figure 47: Estimated time series of spawning output (in millions of eggs).




Figure 48: Estimated time series of total biomass (mt).




Figure 49: Estimated time series of fraction of unfished spawning output.




Figure 50: Comparison of the time series of spawning output between the 2015 and 2023 assessment results.




Figure 51: Comparison of the time series of relative spawning output between the 2015 and 2023 assessment results.




Figure 52: Log relative change ($\log((\text{Model_sensi}-\text{Model_ref})/\text{Model_ref}))$) in data treatment for 5 derived quantities. Colored boxes indicate 95 percent confidence interval of the reference model.




Figure 53: Spawning output (in millions of eggs) time series by data treatment compared to the reference model.




Figure 54: Relative spawning output time series by data treatment compared to the reference model. Missing scenarios mean the spawning output was too large to show.




Figure 55: Log relative change ($\log((\text{Model_sensi}-\text{Model_ref})/\text{Model_ref}))$) in model specification scenario for 5 derived quantities. Colored boxes indicate 95 percent confidence interval of the reference model.




Figure 56: Spawning output (in millions of eggs) time series by model specification scenario compared to the reference model. Missing scenarios mean the spawning output was too large to show.




Figure 57: Relative spawning output time series by model specification scenario compared to the reference model.




Figure 58: Acoustic-visual survey catchability likelihood profile (change in the negative log-likelihood across a range of catchability values) and derived quantities.




Figure 59: Acoustic-visual survey catchability likelihood profile for each of the likelihood components.




Figure 60: Beverton-Holt steepness parameter likelihood profile (change in the negative log-likelihood across a range of steepness values) and derived quantities.




Figure 61: Beverton-Holt steepness parameter likelihood profile for each of the likelihood components.




Figure 62: Female and male M multi-parameter likelihood profile and derived quantities. Red lines in the top left figure indicate significantly similar values compared to the reference model. Broken and solid lines in the bottom right figure indicate target and limit reference points, respectively.




Figure 63: Change in the estimate of spawning output when the most recent 10 years of data area removed sequentially.




Figure 64: Change in the estimate of fraction unfished when the most recent 10 years of data area removed sequentially.




Figure 65: Estimated 1 - relative spawning ratio (SPR) by year.




Figure 66: Phase plot of the relative biomass (also referred to as fraction unfished) versus the SPR ratio where each point represents the biomass ratio at the start of the year and the relative fishing intensity in that same year. Lines through the final point show the 95 percent intervals based on the asymptotic uncertainty for each dimension. The shaded ellipse is a 95 percent region which accounts for the estimated correlations between the biomass ratio and SPR ratio.




Figure 67: Equilibrium yield curve for the reference model. Values are based on the 2020 fishery selectivities and with steepness fixed at 0.72.

9 Appendix A: Detailed Fit to Length Composition Data




Figure 68: Length comps, whole catch, Trawl_wdis.'N adj.' is the input sample size after data-weighting adjustment. N eff. is the calculated effective sample size used in the McAllister-Ianelli tuning method.




Figure 69: Length comps, whole catch, Non-Trawl_wdis (plot 1 of 2). 'N adj.' is the input sample size after data-weighting adjustment. N eff. is the calculated effective sample size used in the McAllister-Ianelli tuning method.




Figure 70: Length comps, whole catch, Non-Trawl_wdis (plot 1 of 2). 'N adj.' is the input sample size after data-weighting adjustment. N eff. is the calculated effective sample size used in the McAllister-Ianelli tuning method. (plot 2 of 2).




Figure 71: Length comps, whole catch, Ocean (plot 1 of 2). 'N adj.' is the input sample size after data-weighting adjustment. N eff. is the calculated effective sample size used in the McAllister-Ianelli tuning method.




Figure 72: Length comps, whole catch, Ocean (plot 1 of 2). 'N adj.' is the input sample size after data-weighting adjustment. N eff. is the calculated effective sample size used in the McAllister-Ianelli tuning method. (plot 2 of 2).




Figure 73: Length comps, whole catch, Shore.'N adj.' is the input sample size after data-weighting adjustment. N eff. is the calculated effective sample size used in the McAllister-Ianelli tuning method.




Figure 74: Length comps, whole catch, MPA.'N adj.' is the input sample size after data-weighting adjustment. N eff. is the calculated effective sample size used in the McAllister-Ianelli tuning method.

10 Appendix B: Fit to Conditional-Age-at-Length Composition Data




Figure 75: Pearson residuals, whole catch, Trawl_wdis (max=13.44) (plot 1 of 3).




Figure 76: Pearson residuals, whole catch, Trawl_wdis (max=13.44) (plot 2 of 3).




Figure 77: Pearson residuals, whole catch, Trawl_wdis (max=13.44) (plot 3 of 3).




Figure 78: Pearson residuals, whole catch, Non-Trawl_wdis (max=12.48) (plot 1 of 5).




Figure 79: Pearson residuals, whole catch, Non-Trawl_wdis (max=12.48) (plot 2 of 5).




Figure 80: Pearson residuals, whole catch, Non-Trawl_wdis (max=12.48) (plot 3 of 5).




Figure 81: Pearson residuals, whole catch, Non-Trawl_wdis (max=12.48) (plot 4 of 5).




Figure 82: Pearson residuals, whole catch, Ocean (max=13.12) (plot 1 of 5).




Figure 83: Pearson residuals, whole catch, Ocean (max=13.12) (plot 2 of 5).




Figure 84: Pearson residuals, whole catch, Ocean (max=13.12) (plot 3 of 5).




Figure 85: Pearson residuals, whole catch, Ocean (max=13.12) (plot 4 of 5).




Figure 86: Pearson residuals, whole catch, Ocean (max=13.12) (plot 5 of 5).

11 Appendix C: Fit to Conditional-Age-at-Length Composition Data




Figure 87: Trawl conditional AAL plot (plot 1 of 5) showing mean age (left panel) and standard deviation (right panel. Shaded areas are 90 percent CIs).




Figure 88: Trawl conditional AAL plot (plot 2 of 5).




Figure 89: Trawl conditional AAL plot (plot 3 of 5).




Figure 90: Trawl' conditional AAL plot (plot 4 of 5).




Figure 91: Trawl conditional AAL plot (plot 5 of 5).




Figure 92: Non-trawl conditional AAL plot (plot 1 of 7) showing mean age (left panel) and standard deviation (right panel). Shaded areas are 90 percent CIs.




Figure 93: Non-trawl conditional AAL plot (plot 2 of 7).




Figure 94: Non-trawl conditional AAL plot (plot 3 of 7).




Figure 95: Non-trawl conditional AAL plot (plot 3 of 7).




Figure 96: Non-trawl conditional AAL plot (plot 4 of 7).




Figure 97: Non-trawl conditional AAL plot (plot 5 of 7).




Figure 98: Non-trawl conditional AAL plot (plot 6 of 7).




Figure 99: Non-trawl conditional AAL plot (plot 7 of 7).




Figure 100: Ocean boat conditional AAL plot (plot 1 of 6) showing mean age (left panel) and standard deviation (right panel. Shaded areas are 90 percent CIs).




Figure 101: Ocean boat conditional AAL plot (plot 2 of 6).




Figure 102: Ocean boat conditional AAL plot (plot 3 of 6).




Figure 103: Ocean boat conditional AAL plot (plot 3 of 6).




Figure 104: Ocean boat conditional AAL plot (plot 4 of 6).

12 Appendix D: Passive Fit to Marginal Age Composition Data




Figure 105: Excluded age comps, whole catch, Trawl_wdis.'N adj.' is the input sample size after data-weighting adjustment. N eff. is the calculated effective sample size used in the McAllister-Ianelli tuning method.




Figure 106: Excluded age comps, whole catch, Non-Trawl_wdis (plot 1 of 2). 'N adj.' is the input sample size after data-weighting adjustment. N eff. is the calculated effective sample size used in the McAllister-Ianelli tuning method.




Figure 107: Excluded age comps, whole catch, Non-Trawl_wdis (plot 1 of 2).‘N adj.’ is the input sample size after data-weighting adjustment. N eff. is the calculated effective sample size used in the McAllister-Ianelli tuning method. (plot 2 of 2).




Figure 108: Excluded age comps, whole catch, Ocean.'N adj.' is the input sample size after data-weighting adjustment. N eff. is the calculated effective sample size used in the McAllister-Ianelli tuning method.

13 Appendix E: Numbers at Age Plot

13.1 Females




Figure 109: Female black rockfish mean age over time.

13.2 Males




Figure 110: Male black rockfish mean age over time.

14 Appendix F: Numbers at Length Plot

14.1 Females




Figure 111: Female black rockfish mean length (cm) over time.

14.2 Males




Figure 112: Male black rockfish mean length over time.