

In the last class we discussed some basic notions of topology.

Recall that a subset  $U \subseteq \mathbb{C}$  is said to be open if given any point  $z_0 \in \mathbb{C}$ , there exists a real number  $\delta > 0$  such that  $B_\delta(z_0) \subseteq U$ . Here  $B_\delta(z_0)$  is the open ball of radius  $\delta$  around  $z_0$ .

In the last class we discussed some basic notions of topology.

Recall that a subset  $U \subseteq \mathbb{C}$  is said to be open if given any point  $z_0 \in \mathbb{C}$ , there exists a real number  $\delta > 0$  such that  $B_\delta(z_0) \subseteq U$ .

Here  $B_\delta(z_0)$  is the open ball of radius  $\delta$  around  $z_0$ .

Intuitively this means that if some point belongs to  $U$ , all sufficiently nearby points belong to  $U$ . The prototypical examples are  $\mathbb{C}$  and the open disc minus finitely many points.

A subset  $Z \subseteq \mathbb{C}$  is closed if its complement is open. Example: For any continuous function  $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ ,  $f(x, y) = 0$  defines a closed subset of  $\mathbb{C}$ .

In the last class we discussed some basic notions of topology.

Recall that a subset  $U \subseteq \mathbb{C}$  is said to be open if given any point  $z_0 \in \mathbb{C}$ , there exists a real number  $\delta > 0$  such that  $B_\delta(z_0) \subseteq U$ .

Here  $B_\delta(z_0)$  is the open ball of radius  $\delta$  around  $z_0$ .

Intuitively this means that if some point belongs to  $U$ , all sufficiently nearby points belong to  $U$ . The prototypical examples are  $\mathbb{C}$  and the open disc minus finitely many points.

A subset  $Z \subseteq \mathbb{C}$  is closed if its complement is open. Example: For any continuous function  $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ ,  $f(x, y) = 0$  defines a closed subset of  $\mathbb{C}$ .

Its a more non-trivial fact that  $\mathbb{C}$  has no subsets which are both open and closed other than  $\mathbb{C}$  and  $\emptyset$

# MA 205 Complex Analysis: CR Equations

August 20, 2020

# Introduction

In the last class, we introduced complex numbers and studied complex valued functions defined on a domain in  $\mathbb{C}$ .

# Introduction

In the last class, we introduced complex numbers and studied complex valued functions defined on a domain in  $\mathbb{C}$ . We stated the fact that every polynomial of degree  $n$  with complex coefficients has exactly  $n$  roots in  $\mathbb{C}$ .

# Introduction

In the last class, we introduced complex numbers and studied complex valued functions defined on a domain in  $\mathbb{C}$ . We stated the fact that every polynomial of degree  $n$  with complex coefficients has exactly  $n$  roots in  $\mathbb{C}$ . This is called the fundamental theorem of algebra.

# Introduction

In the last class, we introduced complex numbers and studied complex valued functions defined on a domain in  $\mathbb{C}$ . We stated the fact that every polynomial of degree  $n$  with complex coefficients has exactly  $n$  roots in  $\mathbb{C}$ . This is called the fundamental theorem of algebra. We introduced complex-differentiability of a function  $f : \Omega \subset \mathbb{C} \rightarrow \mathbb{C}$ , where  $\Omega$  is an open subset of  $\mathbb{C}$ .



In the last class, we introduced complex numbers and studied complex valued functions defined on a domain in  $\mathbb{C}$ . We stated the fact that every polynomial of degree  $n$  with complex coefficients has exactly  $n$  roots in  $\mathbb{C}$ . This is called the fundamental theorem of algebra. We introduced complex-differentiability of a function  $f : \Omega \subset \mathbb{C} \rightarrow \mathbb{C}$ , where  $\Omega$  is an open subset of  $\mathbb{C}$ . We also stated the fact that if  $f$  is once differentiable in  $\Omega$ , then it is infinitely many times differentiable in  $\Omega$ .

# Cauchy-Riemann Equations

Today, first we'll derive the so called Cauchy-Riemann equations. There are two Cauchy-Riemann equations, and these are partial differential equations; i.e., equations containing partial derivatives.

# Cauchy-Riemann Equations

Today, first we'll derive the so called Cauchy-Riemann equations. There are two Cauchy-Riemann equations, and these are partial differential equations; i.e., equations containing partial derivatives. If  $f$  is complex differentiable at a point  $z_0 = a + \imath b$ , then these two equations will be satisfied at the point  $(a, b)$ .

# Cauchy-Riemann Equations

Today, first we'll derive the so called Cauchy-Riemann equations. There are two Cauchy-Riemann equations, and these are partial differential equations; i.e., equations containing partial derivatives. If  $f$  is complex differentiable at a point  $z_0 = a + \imath b$ , then these two equations will be satisfied at the point  $(a, b)$ .

Let  $f : \Omega \subset \mathbb{C} \rightarrow \mathbb{C}$  be differentiable at  $z_0 \in \Omega$ . Thus,

$$\lim_{z \rightarrow z_0} \frac{f(z) - f(z_0)}{z - z_0} = f'(z_0)$$

exists.

# Cauchy-Riemann Equations

Today, first we'll derive the so called Cauchy-Riemann equations. There are two Cauchy-Riemann equations, and these are partial differential equations; i.e., equations containing partial derivatives. If  $f$  is complex differentiable at a point  $z_0 = a + \imath b$ , then these two equations will be satisfied at the point  $(a, b)$ .

Let  $f : \Omega \subset \mathbb{C} \rightarrow \mathbb{C}$  be differentiable at  $z_0 \in \Omega$ . Thus,

$$\lim_{z \rightarrow z_0} \frac{f(z) - f(z_0)}{z - z_0} = f'(z_0)$$

exists. In the last class, we have stressed the point that the existence of this complex limit means a lot; the limit exists as  $z$  approaches  $z_0$  along any path. To derive the CR equations, we'll in particular look at the existence of this limit as  $z \rightarrow z_0$  along the  $x$ -direction and the  $y$ -direction.

# Cauchy-Riemann Equations

Let  $z = x + iy$  and  $f(z) = u(x, y) + iv(x, y)$ . Now, as  $z \rightarrow z_0$  in the  $x$ -direction:

# Cauchy-Riemann Equations

Let  $z = x + iy$  and  $f(z) = u(x, y) + iv(x, y)$ . Now, as  $z \rightarrow z_0$  in the  $x$ -direction:

$$\begin{aligned}f'(z_0) &= \lim_{h \rightarrow 0} \frac{f(z_0 + h) - f(z_0)}{h} \\&= \lim_{h \rightarrow 0} \left[ \frac{u(a + h, b) - u(a, b)}{h} + i \frac{v(a + h, b) - v(a, b)}{h} \right] \\&= \lim_{h \rightarrow 0} \frac{u(a + h, b) - u(a, b)}{h} + i \lim_{h \rightarrow 0} \frac{v(a + h, b) - v(a, b)}{h} \\&= u_x(a, b) + iv_x(a, b).\end{aligned}$$

# Cauchy-Riemann Equations

Let  $z = x + iy$  and  $f(z) = u(x, y) + iv(x, y)$ . Now, as  $z \rightarrow z_0$  in the  $x$ -direction:

$$\begin{aligned}f'(z_0) &= \lim_{h \rightarrow 0} \frac{f(z_0 + h) - f(z_0)}{h} \\&= \lim_{h \rightarrow 0} \left[ \frac{u(a + h, b) - u(a, b)}{h} + i \frac{v(a + h, b) - v(a, b)}{h} \right] \\&= \lim_{h \rightarrow 0} \frac{u(a + h, b) - u(a, b)}{h} + i \lim_{h \rightarrow 0} \frac{v(a + h, b) - v(a, b)}{h} \\&= u_x(a, b) + iv_x(a, b).\end{aligned}$$

Similarly, in the  $y$ -direction, we get

$$f'(z_0) = \lim_{k \rightarrow 0} \frac{f(z_0 + ik) - f(z_0)}{ik} = v_y(a, b) - iu_y(a, b).$$



# Cauchy-Riemann Equations

Thus, differentiability of  $f = u + iv$  at  $z_0 = a + ib$  implies that  $u_x, u_y, v_x, v_y$  exist at  $(a, b)$  and they satisfy

$$u_x = v_y \text{ \& } u_y = -v_x$$

# Cauchy-Riemann Equations

Thus, differentiability of  $f = u + iv$  at  $z_0 = a + ib$  implies that  $u_x, u_y, v_x, v_y$  exist at  $(a, b)$  and they satisfy

$$u_x = v_y \text{ \& } u_y = -v_x$$

at  $(a, b)$ . These are the CR equations. If CR equations are not satisfied at a point, then  $f$  is not differentiable at that point.

Example: Consider  $f(z) = |z|^2$ . Here,  $u(x, y) = x^2 + y^2$ ,  $v(x, y) = 0$ . Thus CR equations are satisfied only at the point  $(0, 0)$ .

# Cauchy-Riemann Equations

Thus, differentiability of  $f = u + iv$  at  $z_0 = a + ib$  implies that  $u_x, u_y, v_x, v_y$  exist at  $(a, b)$  and they satisfy

$$u_x = v_y \text{ \& } u_y = -v_x$$

at  $(a, b)$ . These are the CR equations. If CR equations are not satisfied at a point, then  $f$  is not differentiable at that point.

Example: Consider  $f(z) = |z|^2$ . Here,  $u(x, y) = x^2 + y^2$ ,  $v(x, y) = 0$ . Thus CR equations are satisfied only at the point  $(0, 0)$ . We conclude that  $f$  is not differentiable at any point other than  $(0, 0)$ . Can we conclude that  $f$  is differentiable at  $(0, 0)$ ?

# Cauchy-Riemann Equations

Thus, differentiability of  $f = u + iv$  at  $z_0 = a + ib$  implies that  $u_x, u_y, v_x, v_y$  exist at  $(a, b)$  and they satisfy

$$u_x = v_y \text{ \& } u_y = -v_x$$

at  $(a, b)$ . These are the CR equations. If CR equations are not satisfied at a point, then  $f$  is not differentiable at that point.

Example: Consider  $f(z) = |z|^2$ . Here,  $u(x, y) = x^2 + y^2$ ,  $v(x, y) = 0$ . Thus CR equations are satisfied only at the point  $(0, 0)$ . We conclude that  $f$  is not differentiable at any point other than  $(0, 0)$ . Can we conclude that  $f$  is differentiable at  $(0, 0)$ ? Well, we need to check; CR equations give only one direction. In other words, real and imaginary parts of  $f$  satisfying CR equations at a point is necessary but not sufficient for  $f$  to be differentiable at that point. In this example:

$$\lim_{z \rightarrow 0} \frac{f(z) - f(0)}{z - 0} = \lim_{z \rightarrow 0} \frac{|z|^2}{z} = 0.$$

# Cauchy-Riemann Equations

Example:

$$f(z) = \begin{cases} \frac{\bar{z}^2}{z} & \text{if } z \neq 0 \\ 0 & \text{if } z = 0. \end{cases}$$

# Cauchy-Riemann Equations

Example:

$$f(z) = \begin{cases} \frac{\bar{z}^2}{z} & \text{if } z \neq 0 \\ 0 & \text{if } z = 0. \end{cases}$$

Here,

$$u(x, y) = \begin{cases} \frac{x^3 - 3xy^2}{x^2 + y^2} & \text{if } (x, y) \neq (0, 0) \\ 0 & \text{if } (x, y) = (0, 0), \end{cases}$$

$$v(x, y) = \begin{cases} \frac{-3x^2y + y^3}{x^2 + y^2} & \text{if } (x, y) \neq (0, 0) \\ 0 & \text{if } (x, y) = (0, 0). \end{cases}$$

# Cauchy-Riemann Equations

Example:

$$f(z) = \begin{cases} \frac{\bar{z}^2}{z} & \text{if } z \neq 0 \\ 0 & \text{if } z = 0. \end{cases}$$

Here,

$$u(x, y) = \begin{cases} \frac{x^3 - 3xy^2}{x^2 + y^2} & \text{if } (x, y) \neq (0, 0) \\ 0 & \text{if } (x, y) = (0, 0), \end{cases}$$

$$v(x, y) = \begin{cases} \frac{-3x^2y + y^3}{x^2 + y^2} & \text{if } (x, y) \neq (0, 0) \\ 0 & \text{if } (x, y) = (0, 0). \end{cases}$$

Check that CR equations are satisfied at  $(0, 0)$ . You'll get  $u_x = v_y = 1$  and  $u_y = -v_x = 0$  at  $(0, 0)$ .

# Cauchy-Riemann Equations

But,  $f$  is not differentiable at 0.



# Cauchy-Riemann Equations

But,  $f$  is not differentiable at 0.

$$\lim_{z \rightarrow 0} \frac{f(z) - f(0)}{z - 0} = \lim_{z \rightarrow 0} \frac{\bar{z}^2}{z^2} = \lim_{(x,y) \rightarrow (0,0)} \frac{(x - iy)^2}{(x + iy)^2}.$$

# Cauchy-Riemann Equations

But,  $f$  is not differentiable at 0.

$$\lim_{z \rightarrow 0} \frac{f(z) - f(0)}{z - 0} = \lim_{z \rightarrow 0} \frac{\bar{z}^2}{z^2} = \lim_{(x,y) \rightarrow (0,0)} \frac{(x - iy)^2}{(x + iy)^2}.$$

If  $(x, y) \rightarrow (0, 0)$  via either of the axes, this limit is 1. If  $(x, y) \rightarrow (0, 0)$  via  $y = x$ , this limit is  $-1$ . So limit does not exist.

# Cauchy-Riemann Equations

If  $z = x + iy$ , then,

$$x = \frac{z + \bar{z}}{2}, \quad y = \frac{z - \bar{z}}{2i}.$$

# Cauchy-Riemann Equations

If  $z = x + iy$ , then,

$$x = \frac{z + \bar{z}}{2}, \quad y = \frac{z - \bar{z}}{2i}.$$

Suppose for a moment that  $z$  and  $\bar{z}$  are independent variables!  
Formally applying chain rule:

$$\frac{\partial f}{\partial z} = \frac{\partial f}{\partial x} \cdot \frac{1}{2} + \frac{\partial f}{\partial y} \cdot \frac{1}{2i} = \frac{1}{2} \left( \frac{\partial f}{\partial x} - i \frac{\partial f}{\partial y} \right).$$

# Cauchy-Riemann Equations

If  $z = x + iy$ , then,

$$x = \frac{z + \bar{z}}{2}, \quad y = \frac{z - \bar{z}}{2i}.$$

Suppose for a moment that  $z$  and  $\bar{z}$  are independent variables!  
Formally applying chain rule:

$$\frac{\partial f}{\partial z} = \frac{\partial f}{\partial x} \cdot \frac{1}{2} + \frac{\partial f}{\partial y} \cdot \frac{1}{2i} = \frac{1}{2} \left( \frac{\partial f}{\partial x} - i \frac{\partial f}{\partial y} \right).$$

Similarly,

$$\frac{\partial f}{\partial \bar{z}} = \frac{1}{2} \left( \frac{\partial f}{\partial x} + i \frac{\partial f}{\partial y} \right).$$

# Cauchy-Riemann Equations

Motivated by this, we introduce the symbols:

# Cauchy-Riemann Equations

Motivated by this, we introduce the symbols:

$$\frac{\partial}{\partial z} = \frac{1}{2} \left( \frac{\partial}{\partial x} - i \frac{\partial}{\partial y} \right); \quad \frac{\partial}{\partial \bar{z}} = \frac{1}{2} \left( \frac{\partial}{\partial x} + i \frac{\partial}{\partial y} \right).$$

# Cauchy-Riemann Equations

Motivated by this, we introduce the symbols:

$$\frac{\partial}{\partial z} = \frac{1}{2} \left( \frac{\partial}{\partial x} - i \frac{\partial}{\partial y} \right); \quad \frac{\partial}{\partial \bar{z}} = \frac{1}{2} \left( \frac{\partial}{\partial x} + i \frac{\partial}{\partial y} \right).$$

Note that CR equations now can be written as

$$\frac{\partial f}{\partial \bar{z}} = 0.$$



# Cauchy-Riemann Equations

Motivated by this, we introduce the symbols:

$$\frac{\partial}{\partial z} = \frac{1}{2} \left( \frac{\partial}{\partial x} - i \frac{\partial}{\partial y} \right); \quad \frac{\partial}{\partial \bar{z}} = \frac{1}{2} \left( \frac{\partial}{\partial x} + i \frac{\partial}{\partial y} \right).$$

Note that CR equations now can be written as

$$\frac{\partial f}{\partial \bar{z}} = 0.$$

Since  $\frac{\partial f}{\partial \bar{z}} = \frac{1}{2}(u_x + iv_x + iu_y - v_y)$

# Cauchy-Riemann Equations

We can of course view  $f : \Omega \subset \mathbb{C} \rightarrow \mathbb{C}$  as a function of two real variables;

$$f(x, y) = (u(x, y), v(x, y)).$$

For such functions, in MA 105, you have seen the notion of the total derivative.

# Cauchy-Riemann Equations

We can of course view  $f : \Omega \subset \mathbb{C} \rightarrow \mathbb{C}$  as a function of two real variables;

$$f(x, y) = (u(x, y), v(x, y)).$$

For such functions, in MA 105, you have seen the notion of the total derivative. Recall:  $f$  is differentiable at  $(a, b)$  if there exists a  $2 \times 2$  matrix  $Df(a, b)$  such that

$$\lim_{(h,k) \rightarrow (0,0)} \frac{\|f(a+h, b+k) - f(a, b) - Df(a, b) \begin{bmatrix} h \\ k \end{bmatrix}\|}{\|(h, k)\|} = 0.$$

# Cauchy-Riemann Equations

We can of course view  $f : \Omega \subset \mathbb{C} \rightarrow \mathbb{C}$  as a function of two real variables;

$$f(x, y) = (u(x, y), v(x, y)).$$

For such functions, in MA 105, you have seen the notion of the total derivative. Recall:  $f$  is differentiable at  $(a, b)$  if there exists a  $2 \times 2$  matrix  $Df(a, b)$  such that

$$\lim_{(h,k) \rightarrow (0,0)} \frac{\|f(a+h, b+k) - f(a, b) - Df(a, b) \begin{bmatrix} h \\ k \end{bmatrix}\|}{\|(h, k)\|} = 0.$$

Of course, if total derivative exists, then all the partial derivatives exist, and

$$Df = \begin{bmatrix} u_x & u_y \\ v_x & v_y \end{bmatrix}.$$

# Cauchy-Riemann Equations

We can of course view  $f : \Omega \subset \mathbb{C} \rightarrow \mathbb{C}$  as a function of two real variables;

$$f(x, y) = (u(x, y), v(x, y)).$$

For such functions, in MA 105, you have seen the notion of the total derivative. Recall:  $f$  is differentiable at  $(a, b)$  if there exists a  $2 \times 2$  matrix  $Df(a, b)$  such that

$$\lim_{(h,k) \rightarrow (0,0)} \frac{\|f(a+h, b+k) - f(a, b) - Df(a, b) \begin{bmatrix} h \\ k \end{bmatrix}\|}{\|(h, k)\|} = 0.$$

Of course, if total derivative exists, then all the partial derivatives exist, and

$$Df = \begin{bmatrix} u_x & u_y \\ v_x & v_y \end{bmatrix}.$$

Existence of partial derivatives does not imply the existence of total derivative, but existence of partial derivatives which are continuous throughout the domain does imply the existence of total derivative.

# Cauchy-Riemann Equations

Exercise: Show that if  $f$  is complex differentiable, then  $f$  is real differentiable; i.e.,  $f$  has a total derivative as a function of two real variables. Show that the converse is not true.

(At the moment solve this exercise assuming the continuity of the first partial derivatives of  $u$  and  $v$ . We shall see later that this assumption can be removed (it is automatic)).

# Cauchy-Riemann Equations

Exercise: Show that if  $f$  is complex differentiable, then  $f$  is real differentiable; i.e.,  $f$  has a total derivative as a function of two real variables. Show that the converse is not true.

(At the moment solve this exercise assuming the continuity of the first partial derivatives of  $u$  and  $v$ . We shall see later that this assumption can be removed (it is automatic)).

Thus, complex differentiability implies:

- real differentiability
- real and imaginary parts satisfy CR.

# Cauchy-Riemann Equations

Exercise: Show that if  $f$  is complex differentiable, then  $f$  is real differentiable; i.e.,  $f$  has a total derivative as a function of two real variables. Show that the converse is not true.

(At the moment solve this exercise assuming the continuity of the first partial derivatives of  $u$  and  $v$ . We shall see later that this assumption can be removed (it is automatic)).

Thus, complex differentiability implies:

- real differentiability
- real and imaginary parts satisfy CR.

What if we assume both these? Can we then say  $f$  is complex differentiable? And the answer is Yes.

Proof: Since  $f = u + iv$  is real differentiable,

$$\lim_{(x,y) \rightarrow (a,b)} \frac{\left\| \begin{bmatrix} u(x,y) \\ v(x,y) \end{bmatrix} - \begin{bmatrix} u(a,b) \\ v(a,b) \end{bmatrix} - \begin{bmatrix} u_x & u_y \\ v_x & v_y \end{bmatrix} \begin{bmatrix} x-a \\ y-b \end{bmatrix} \right\|}{\|(x-a, y-b)\|} = 0.$$



# Cauchy-Riemann Equations

Note that the numerator is nothing but

$$|f(z) - f(z_0) - \alpha(x - a) - \beta(y - b)|,$$

where  $\alpha = u_x + iv_x$ ,  $\beta = u_y + iv_y$ .

# Cauchy-Riemann Equations

Note that the numerator is nothing but

$$|f(z) - f(z_0) - \alpha(x - a) - \beta(y - b)|,$$

where  $\alpha = u_x + iv_x$ ,  $\beta = u_y + iv_y$ . Define

$$\eta(z) = \frac{f(z) - f(z_0) - \alpha(x - a) - \beta(y - b)}{z - z_0}.$$

Observe that

$$\lim_{z \rightarrow z_0} \eta(z) = 0.$$

# Cauchy-Riemann Equations

Note that the numerator is nothing but

$$|f(z) - f(z_0) - \alpha(x - a) - \beta(y - b)|,$$

where  $\alpha = u_x + iv_x$ ,  $\beta = u_y + iv_y$ . Define

$$\eta(z) = \frac{f(z) - f(z_0) - \alpha(x - a) - \beta(y - b)}{z - z_0}.$$

Observe that

$$\lim_{z \rightarrow z_0} \eta(z) = 0.$$

Thus,

$$f(z) - f(z_0) = \alpha(x - a) + \beta(y - b) + \eta(z)(z - z_0),$$

with  $\eta(z) \rightarrow 0$  as  $z \rightarrow z_0$ .

# Cauchy-Riemann Equations

Note that the numerator is nothing but

$$|f(z) - f(z_0) - \alpha(x - a) - \beta(y - b)|,$$

where  $\alpha = u_x + iv_x$ ,  $\beta = u_y + iv_y$ . Define

$$\eta(z) = \frac{f(z) - f(z_0) - \alpha(x - a) - \beta(y - b)}{z - z_0}.$$

Observe that

$$\lim_{z \rightarrow z_0} \eta(z) = 0.$$

Thus,

$$f(z) - f(z_0) = \alpha(x - a) + \beta(y - b) + \eta(z)(z - z_0),$$

with  $\eta(z) \rightarrow 0$  as  $z \rightarrow z_0$ . Write this as

$$f(z) - f(z_0) = \frac{\alpha - i\beta}{2}(z - z_0) + \frac{\alpha + i\beta}{2}\overline{z - z_0} + \eta(z)(z - z_0).$$

# Cauchy-Riemann Equations

Thus,

$$\frac{f(z) - f(z_0)}{z - z_0} = \frac{\partial f}{\partial z}(z_0) + \frac{\partial f}{\partial \bar{z}}(z_0) \frac{\overline{z - z_0}}{z - z_0} + \eta(z).$$

# Cauchy-Riemann Equations

Thus,

$$\frac{f(z) - f(z_0)}{z - z_0} = \frac{\partial f}{\partial z}(z_0) + \frac{\partial f}{\partial \bar{z}}(z_0) \frac{\overline{z - z_0}}{z - z_0} + \eta(z).$$

Question is whether the lhs limit exists as  $z \rightarrow z_0$ . This exists if and only if the rhs limit exists.

# Cauchy-Riemann Equations

Thus,

$$\frac{f(z) - f(z_0)}{z - z_0} = \frac{\partial f}{\partial z}(z_0) + \frac{\partial f}{\partial \bar{z}}(z_0) \frac{\overline{z - z_0}}{z - z_0} + \eta(z).$$

Question is whether the lhs limit exists as  $z \rightarrow z_0$ . This exists if and only if the rhs limit exists. Since,

$$\lim_{z \rightarrow z_0} \frac{\overline{z - z_0}}{z - z_0}$$

does not exist (why?) and  $\lim_{z \rightarrow z_0} \eta(z)$  exists, this happens if and only if

$$\frac{\partial f}{\partial \bar{z}}(z_0) = 0.$$

i.e., CR equations are satisfied at  $z_0$ .

# Cauchy-Riemann Equations

Thus,

$$\frac{f(z) - f(z_0)}{z - z_0} = \frac{\partial f}{\partial z}(z_0) + \frac{\partial f}{\partial \bar{z}}(z_0) \frac{\overline{z - z_0}}{z - z_0} + \eta(z).$$

Question is whether the lhs limit exists as  $z \rightarrow z_0$ . This exists if and only if the rhs limit exists. Since,

$$\lim_{z \rightarrow z_0} \frac{\overline{z - z_0}}{z - z_0}$$

does not exist (why?) and  $\lim_{z \rightarrow z_0} \eta(z)$  exists, this happens if and only if

$$\frac{\partial f}{\partial \bar{z}}(z_0) = 0.$$

i.e., CR equations are satisfied at  $z_0$ . Also, if this is the case,

$$f'(z_0) = \frac{\partial f}{\partial z}(z_0),$$

since  $\lim_{z \rightarrow z_0} \eta(z) = 0$ .



# Cauchy-Riemann Equations

Corollary: Let  $f : \Omega \subset \mathbb{C} \rightarrow \mathbb{C}$  be such that it has continuous partial derivatives throughout  $\Omega$ . Then if they satisfy the CR equations at a point,  $f$  is differentiable at that point. (Proof?)

# Cauchy-Riemann Equations

Corollary: Let  $f : \Omega \subset \mathbb{C} \rightarrow \mathbb{C}$  be such that it has continuous partial derivatives throughout  $\Omega$ . Then if they satisfy the CR equations at a point,  $f$  is differentiable at that point. (Proof?)

Exercise: Show that  $f(z) = e^x(\cos y + i \sin y)$  is holomorphic throughout  $\mathbb{C}$ . Note that  $f'(z) = f(z)$ . This is the complex exponential function.

# Cauchy-Riemann Equations

Corollary: Let  $f : \Omega \subset \mathbb{C} \rightarrow \mathbb{C}$  be such that it has continuous partial derivatives throughout  $\Omega$ . Then if they satisfy the CR equations at a point,  $f$  is differentiable at that point. (Proof?)

Exercise: Show that  $f(z) = e^x(\cos y + i \sin y)$  is holomorphic throughout  $\mathbb{C}$ . Note that  $f'(z) = f(z)$ . This is the complex exponential function.

The assumptions in the statement of the corollary can be weakened. In fact, the following is true:

# Cauchy-Riemann Equations

Corollary: Let  $f : \Omega \subset \mathbb{C} \rightarrow \mathbb{C}$  be such that it has continuous partial derivatives throughout  $\Omega$ . Then if they satisfy the CR equations at a point,  $f$  is differentiable at that point. (Proof?)

Exercise: Show that  $f(z) = e^x(\cos y + i \sin y)$  is holomorphic throughout  $\mathbb{C}$ . Note that  $f'(z) = f(z)$ . This is the complex exponential function.

The assumptions in the statement of the corollary can be weakened. In fact, the following is true:

## Theorem

*Let  $f$  be continuous on  $\Omega$ . Suppose the partial derivatives exist and satisfy the Cauchy-Riemann equations at every point in  $\Omega$ . Then  $f$  is holomorphic in  $\Omega$ .*

We shall not prove this theorem.

# Cauchy-Riemann Equations

Exercise: Show that the CR equations take the form

$$u_r = \frac{1}{r}v_\theta \text{ \& \; } v_r = -\frac{1}{r}u_\theta$$

in polar coordinates.

# Harmonic Functions

A real valued function  $u : U \subset \mathbb{R}^2 \rightarrow \mathbb{R}$  is called harmonic if it is twice continuously differentiable and satisfies  $u_{xx} + u_{yy} = 0$  on  $U$ .

# Harmonic Functions

A real valued function  $u : U \subset \mathbb{R}^2 \rightarrow \mathbb{R}$  is called harmonic if it is twice continuously differentiable and satisfies  $u_{xx} + u_{yy} = 0$  on  $U$ . If  $f = u + iv$  is holomorphic on  $\Omega$ , then both  $u$  and  $v$  are harmonic on  $\Omega$ .

# Harmonic Functions

A real valued function  $u : U \subset \mathbb{R}^2 \rightarrow \mathbb{R}$  is called harmonic if it is twice continuously differentiable and satisfies  $u_{xx} + u_{yy} = 0$  on  $U$ . If  $f = u + iv$  is holomorphic on  $\Omega$ , then both  $u$  and  $v$  are harmonic on  $\Omega$ . Indeed, CR equations tell us that

$$u_x = v_y \text{ \& } u_y = -v_x.$$

Thus,

$$u_{xx} + u_{yy} = v_{xy} - v_{xy} = 0.$$

Similarly for  $v$ .



# Harmonic Functions

Suppose  $u$  and  $v$  are harmonic functions on  $\Omega$ . We say that  $v$  is a harmonic conjugate of  $u$  if  $f = u + iv$  is holomorphic in  $\Omega$ .

# Harmonic Functions

Suppose  $u$  and  $v$  are harmonic functions on  $\Omega$ . We say that  $v$  is a harmonic conjugate of  $u$  if  $f = u + iv$  is holomorphic in  $\Omega$ .

Example:  $v(x, y) = 2xy$  is a harmonic conjugate of  $x^2 - y^2$  in any domain. Indeed,  $f(z) = z^2$  is holomorphic everywhere.

Note that  $v$  is a harmonic conjugate of  $u$  does not mean that  $u$  is a harmonic conjugate of  $v$ ! In fact:

# Harmonic Functions

Suppose  $u$  and  $v$  are harmonic functions on  $\Omega$ . We say that  $v$  is a harmonic conjugate of  $u$  if  $f = u + iv$  is holomorphic in  $\Omega$ .

Example:  $v(x, y) = 2xy$  is a harmonic conjugate of  $x^2 - y^2$  in any domain. Indeed,  $f(z) = z^2$  is holomorphic everywhere.

Note that  $v$  is a harmonic conjugate of  $u$  does not mean that  $u$  is a harmonic conjugate of  $v$ ! In fact:

Exercise: If  $u$  and  $v$  are harmonic conjugates of each other, show that they are constant functions.

# Harmonic Functions

Suppose  $u$  and  $v$  are harmonic functions on  $\Omega$ . We say that  $v$  is a harmonic conjugate of  $u$  if  $f = u + iv$  is holomorphic in  $\Omega$ .

Example:  $v(x, y) = 2xy$  is a harmonic conjugate of  $x^2 - y^2$  in any domain. Indeed,  $f(z) = z^2$  is holomorphic everywhere.

Note that  $v$  is a harmonic conjugate of  $u$  does not mean that  $u$  is a harmonic conjugate of  $v$ ! In fact:

Exercise: If  $u$  and  $v$  are harmonic conjugates of each other, show that they are constant functions.

Here's a general method to find a harmonic conjugate: given a harmonic  $u$ , find  $u_x$ . Equate  $v_y = u_x$  and integrate wrt  $y$ . You'll get  $v(x, y) = \dots + \phi(x)$ . Now  $v_x = \dots + \phi'(x)$ . Equate this to  $-u_y$  to find  $\phi(x)$ . That gives you  $v$ .

# Harmonic Functions

Suppose  $u$  and  $v$  are harmonic functions on  $\Omega$ . We say that  $v$  is a harmonic conjugate of  $u$  if  $f = u + iv$  is holomorphic in  $\Omega$ .

Example:  $v(x, y) = 2xy$  is a harmonic conjugate of  $x^2 - y^2$  in any domain. Indeed,  $f(z) = z^2$  is holomorphic everywhere.

Note that  $v$  is a harmonic conjugate of  $u$  does not mean that  $u$  is a harmonic conjugate of  $v$ ! In fact:

Exercise: If  $u$  and  $v$  are harmonic conjugates of each other, show that they are constant functions.

Here's a general method to find a harmonic conjugate: given a harmonic  $u$ , find  $u_x$ . Equate  $v_y = u_x$  and integrate wrt  $y$ . You'll get  $v(x, y) = \dots + \phi(x)$ . Now  $v_x = \dots + \phi'(x)$ . Equate this to  $-u_y$  to find  $\phi(x)$ . That gives you  $v$ .

This might give you an impression that you can always find a harmonic conjugate, but this is not so.

# Harmonic Functions

Suppose  $u$  and  $v$  are harmonic functions on  $\Omega$ . We say that  $v$  is a harmonic conjugate of  $u$  if  $f = u + iv$  is holomorphic in  $\Omega$ .

Example:  $v(x, y) = 2xy$  is a harmonic conjugate of  $x^2 - y^2$  in any domain. Indeed,  $f(z) = z^2$  is holomorphic everywhere.

Note that  $v$  is a harmonic conjugate of  $u$  does not mean that  $u$  is a harmonic conjugate of  $v$ ! In fact:

Exercise: If  $u$  and  $v$  are harmonic conjugates of each other, show that they are constant functions.

Here's a general method to find a harmonic conjugate: given a harmonic  $u$ , find  $u_x$ . Equate  $v_y = u_x$  and integrate wrt  $y$ . You'll get  $v(x, y) = \dots + \phi(x)$ . Now  $v_x = \dots + \phi'(x)$ . Equate this to  $-u_y$  to find  $\phi(x)$ . That gives you  $v$ .

This might give you an impression that you can always find a harmonic conjugate, but this is not so.

*Unfortunately this method fails in general. Try and think of the reason !*

# Harmonic Functions

But if  $\Omega$  is “nice”, then every harmonic  $u$  on  $\Omega$  has a harmonic conjugate. Conversely, if every harmonic  $u$  on  $\Omega$  has a harmonic conjugate, then  $\Omega$  has to be “nice”.

# Harmonic Functions

But if  $\Omega$  is “nice”, then every harmonic  $u$  on  $\Omega$  has a harmonic conjugate. Conversely, if every harmonic  $u$  on  $\Omega$  has a harmonic conjugate, then  $\Omega$  has to be “nice”. Thus, the question in analysis: “does every harmonic function has a harmonic conjugate?” is answered by geometry: “answer depends on the shape of the domain”. It’s relevant at this point to recall from MA 105 that curl of grad is always zero but curl free is certainly a grad of something only when the domain is “nice” (for example  $\mathbb{C}$  or a disc in  $\mathbb{C}$ ). Remember this !