

# Signal Processing - 1 by One

Sibi Raj B. Pillai  
Dept of Electrical Engineering  
IIT Bombay



- So Far: Impulse, Sampling, Replacement
- Previous Week: Convolution ( $*$ ) and Interpolation
- Today: Drawing with Sinusoids




- So Far: Impulse, Sampling, Replacement
- Previous Week: Convolution ( $*$ ) and Interpolation
- Today: Drawing with Sinusoids




- So Far: Impulse, Sampling, Replacement
- Previous Week: Convolution ( $*$ ) and Interpolation
- Today: Drawing with Sinusoids



# Sinusoidal Leopards




# Sinusoidal Leopards



Can we plot this picture using sine waves, i.e.  $\sin(2\pi f x)$ .




# Sinusoidal Leopards



Can we plot this picture using sine waves, i.e.  $\sin(2\pi f x)$ .



# Sinusoidal Leopards




Can we plot this picture using sine waves, i.e.  $\sin(2\pi f x)$ .







# Min-Max Error




# Min-Max Error



# Min-Max Error



# Min-Max Error



# Mean Squared Error (MSE)

**DC Approximation:** Find the constant value  $a$  which minimizes

$$\text{MSE} := \int_{\mathbb{T}} |y(t) - a|^2 dt,$$

where we take  $\mathbb{T} = [0, T_d]$  as the interval of interest.



# Mean Squared Error (MSE)

**DC Approximation:** Find the constant value  $a$  which minimizes

$$\text{MSE} := \int_{\mathbb{T}} |y(t) - a|^2 dt,$$

where we take  $\mathbb{T} = [0, T_d]$  as the interval of interest.

*Notation:*

$$\mu = \frac{1}{T_d} \int_{\mathbb{T}} y(t) dt \quad (\text{Mean Value})$$

$$\|y(t)\|_{\ell_2} = \left( \int_{\mathbb{T}} |y(t)|^2 dt \right)^{\frac{1}{2}} \quad (\ell_2\text{-norm})$$

$$\langle y(t), x(t) \rangle = \int_{\mathbb{T}} y(t) x^*(t) dt \quad (\text{Dot product, } \mathbb{C} \text{ valued}).$$



# Mean Squared Error (MSE)

**DC Approximation:** Find the constant value  $a$  which minimizes

$$\text{MSE} := \int_{\mathbb{T}} |y(t) - a|^2 dt,$$

where we take  $\mathbb{T} = [0, T_d]$  as the interval of interest.

*Notation:*

$$\mu = \frac{1}{T_d} \int_{\mathbb{T}} y(t) dt \quad (\text{Mean Value})$$

$$\|y(t)\|_{\ell_2} = \left( \int_{\mathbb{T}} |y(t)|^2 dt \right)^{\frac{1}{2}} \quad (\ell_2\text{-norm})$$

$$\langle y(t), x(t) \rangle = \int_{\mathbb{T}} y(t) x^*(t) dt \quad (\text{Dot product, } \mathbb{C} \text{ valued}).$$

DC Approximation :  $y_{\text{approx}}(t) = \alpha_0, \forall t \in [0, T_d]$  where

$$\alpha_0 = \underset{a \in \mathbb{R}}{\text{argmin}} \int_{\mathbb{T}} |y(t) - a|^2 dt$$



# Minimizing Functions

$$\int_{\mathbb{T}} |y(t) - a|^2 dt = \int_{\mathbb{T}} |y(t) - \mu + \mu - a|^2 dt$$





# Minimizing Functions

$$\begin{aligned}\int_{\mathbb{T}} |y(t) - a|^2 dt &= \int_{\mathbb{T}} |y(t) - \mu + \mu - a|^2 dt \\ &= \int_{\mathbb{T}} |y(t) - \mu|^2 dt + \int_{\mathbb{T}} |a - \mu|^2 dt - 2 \int_{\mathbb{T}} (y(t) - \mu)(a - \mu) dt\end{aligned}$$



# Minimizing Functions

$$\begin{aligned}\int_{\mathbb{T}} |y(t) - a|^2 dt &= \int_{\mathbb{T}} |y(t) - \mu + \mu - a|^2 dt \\&= \int_{\mathbb{T}} |y(t) - \mu|^2 dt + \int_{\mathbb{T}} |a - \mu|^2 dt - 2 \int_{\mathbb{T}} (y(t) - \mu)(a - \mu) dt \\&= \int_{\mathbb{T}} |y(t) - \mu|^2 dt + |a - \mu|^2 T_d - 2(a - \mu) \int_{\mathbb{T}} (y(t) - \mu) dt.\end{aligned}$$



# Minimizing Functions

$$\begin{aligned}\int_{\mathbb{T}} |y(t) - a|^2 dt &= \int_{\mathbb{T}} |y(t) - \mu + \mu - a|^2 dt \\&= \int_{\mathbb{T}} |y(t) - \mu|^2 dt + \int_{\mathbb{T}} |a - \mu|^2 dt - 2 \int_{\mathbb{T}} (y(t) - \mu)(a - \mu) dt \\&= \int_{\mathbb{T}} |y(t) - \mu|^2 dt + |a - \mu|^2 T_d - 2(a - \mu) \int_{\mathbb{T}} (y(t) - \mu) dt.\end{aligned}$$

Notice that  $\int_{\mathbb{T}} (y(t) - \mu) dt = 0$ , by the definition of  $\mu$ .

$$\int_{\mathbb{T}} |y(t) - a|^2 dt \geq \int_{\mathbb{T}} |y(t) - \mu|^2 dt, \quad \forall a \in \mathbb{R}.$$



# Minimizing Functions

$$\begin{aligned}\int_{\mathbb{T}} |y(t) - a|^2 dt &= \int_{\mathbb{T}} |y(t) - \mu + \mu - a|^2 dt \\&= \int_{\mathbb{T}} |y(t) - \mu|^2 dt + \int_{\mathbb{T}} |a - \mu|^2 dt - 2 \int_{\mathbb{T}} (y(t) - \mu)(a - \mu) dt \\&= \int_{\mathbb{T}} |y(t) - \mu|^2 dt + |a - \mu|^2 T_d - 2(a - \mu) \int_{\mathbb{T}} (y(t) - \mu) dt.\end{aligned}$$


Notice that  $\int_{\mathbb{T}} (y(t) - \mu) dt = 0$ , by the definition of  $\mu$ .

$$\int_{\mathbb{T}} |y(t) - a|^2 dt \geq \int_{\mathbb{T}} |y(t) - \mu|^2 dt, \quad \forall a \in \mathbb{R}.$$


Mean Value is the best DC approximation for MSE.



# Mean Animal



# Mean Animal



Why not fit the sinusoids we started with?




# Fitting a Sine Wave

$$b_1 = \operatorname{argmin}_{a \in \mathbb{R}} \int_{\mathbb{T}} \left| y(t) - a \sin\left(\frac{2\pi}{T_d} t\right) \right|^2 dt.$$



# Fitting a Sine Wave




$$b_1 = \operatorname{argmin}_{a \in \mathbb{R}} \int_{\mathbb{T}} \left| y(t) - a \sin\left(\frac{2\pi}{T_d} t\right) \right|^2 dt.$$






# Fitting a Sine Wave



$$b_1 = \operatorname{argmin}_{a \in \mathbb{R}} \int_{\mathbb{T}} \left| y(t) - a \sin\left(\frac{2\pi}{T_d} t\right) \right|^2 dt.$$



# Fitting a Sine Wave



$$b_1 = \operatorname{argmin}_{a \in \mathbb{R}} \int_{\mathbb{T}} \left| y(t) - a \sin\left(\frac{2\pi}{T_d} t\right) \right|^2 dt.$$



# 'Tailored' Approximation

$$\gamma = \operatorname{argmin}_{a \in \mathbb{R}} \int_{\mathbb{T}} |y(t) - a f(t)|^2 dt.$$

The derivative of the MSE term should be zero at  $a = \gamma$ , i.e.

$$\frac{d}{da} \int_{\mathbb{T}} |y(t) - \gamma f(t)|^2 dt = 0$$



## 'Tailored' Approximation

$$\gamma = \operatorname{argmin}_{a \in \mathbb{R}} \int_{\mathbb{T}} |y(t) - a f(t)|^2 dt.$$

The derivative of the MSE term should be zero at  $a = \gamma$ , i.e.

$$\frac{d}{da} \int_{\mathbb{T}} |y(t) - \gamma f(t)|^2 dt = 0$$



## 'Tailored' Approximation

$$\gamma = \operatorname{argmin}_{a \in \mathbb{R}} \int_{\mathbb{T}} |y(t) - af(t)|^2 dt.$$

The derivative of the MSE term should be zero at  $a = \gamma$ , i.e.

$$\frac{d}{da} \int_{\mathbb{T}} |y(t) - \gamma f(t)|^2 dt = 0$$

$$\Rightarrow \langle y(t) - af(t), f(t) \rangle = 0 \text{ at } a = \gamma$$



## 'Tailored' Approximation

$$\gamma = \operatorname{argmin}_{a \in \mathbb{R}} \int_{\mathbb{T}} |y(t) - af(t)|^2 dt.$$

The derivative of the MSE term should be zero at  $a = \gamma$ , i.e.

$$\frac{d}{da} \int_{\mathbb{T}} |y(t) - \gamma f(t)|^2 dt = 0$$

$$\Rightarrow \langle y(t) - af(t), f(t) \rangle = 0 \text{ at } a = \gamma$$

$$\Rightarrow \gamma = \frac{\langle y(t), f(t) \rangle}{\langle f(t), f(t) \rangle}$$



## 'Tailored' Approximation

$$\gamma = \operatorname{argmin}_{a \in \mathbb{R}} \int_{\mathbb{T}} |y(t) - af(t)|^2 dt.$$

The derivative of the MSE term should be zero at  $a = \gamma$ , i.e.

$$\frac{d}{da} \int_{\mathbb{T}} |y(t) - \gamma f(t)|^2 dt = 0$$

$$\Rightarrow \langle y(t) - af(t), f(t) \rangle = 0 \text{ at } a = \gamma$$

$$\Rightarrow \gamma = \frac{\langle y(t), f(t) \rangle}{\langle f(t), f(t) \rangle} = \frac{\langle y(t), f(t) \rangle}{\|f\|_{\ell_2}^2}$$



# Sines and Cosines

$$f(t) = \sin\left(\frac{2\pi}{T_d} t\right)$$





# Sines and Cosines

$$f(t) = \sin\left(\frac{2\pi}{T_d} t\right)$$

Since  $\|f(t)\|_{\ell_2}^2 = \frac{T_d}{2}$ , we get

$$b_1 = \frac{2}{T_d} \int_{\mathbb{T}} y(t) \sin\left(\frac{2\pi}{T_d} t\right) dt.$$



# Sines and Cosines

$$f(t) = \sin\left(\frac{2\pi}{T_d} t\right)$$

Since  $\|f(t)\|_{\ell_2}^2 = \frac{T_d}{2}$ , we get

$$b_1 = \frac{2}{T_d} \int_{\mathbb{T}} y(t) \sin\left(\frac{2\pi}{T_d} t\right) dt.$$

To be fair, we should fit  $\cos\left(\frac{2\pi}{T_d} t\right)$  as well.

$$a_1 = \frac{2}{T_d} \int_{\mathbb{T}} y(t) \cos\left(\frac{2\pi}{T_d} t\right) dt.$$



