

MA 205 Complex Analysis: Some More Theorems

October 2, 2020

Choice of contour for integration

It might be a bit of a mystery as to which contour one should consider for a given contour integral. Here is a general recipe.

Choice of contour for integration

It might be a bit of a mystery as to which contour one should consider for a given contour integral. Here is a general recipe.

Suppose the improper integral is of the form $\int_{-\infty}^{\infty} f(x)dx$. The general idea of course is to find a contour which contains the real line as part of the contour in the “limit”. The choice should be made so that by residue theory one knows the integral over the full contour and such that the integral over the extra added part goes to zero in the limit.

Choice of contour for integration

It might be a bit of a mystery as to which contour one should consider for a given contour integral. Here is a general recipe.

Suppose the improper integral is of the form $\int_{-\infty}^{\infty} f(x)dx$. The general idea of course is to find a contour which contains the real line as part of the contour in the “limit”. The choice should be made so that by residue theory one knows the integral over the full contour and such that the integral over the extra added part goes to zero in the limit.

I. For instance suppose there exists a constant C such that $|f(z)| \leq \frac{C}{|z|^r}$ for sufficiently large $|z|$ and for some $r > 1$ (here $f(z)$ is an extension of $f(x)$ to a function of the complex variable).

Note that this happens for instance in the case when $f(x) = P(x)/Q(x)$ where $\deg Q(x) \geq \deg P(x) + 2$. Then close up the interval with a semicircle into the upper half plane and integrate along the contour and take limit as the radius of semicircle goes to infinity. Use ML inequality to show that the integral along the semicircle goes to zero as radius goes to ∞ .

Choice of contour for integration

In case the integral is from 0 to ∞ , try and relate it to some integral from $-\infty$ to ∞ . For instance the function may have a natural continuation to the negative reals. In case this is not possible, often because $f(x)$ has a singularity at origin; usually a pole, then try using a half annular region $A(0; r, R)$ like we have done in earlier examples.

Choice of contour for integration

In case the integral is from 0 to ∞ , try and relate it to some integral from $-\infty$ to ∞ . For instance the function may have a natural continuation to the negative reals. In case this is not possible, often because $f(x)$ has a singularity at origin; usually a pole, then try using a half annular region $A(0; r, R)$ like we have done in earlier examples. This will avoid the pole and then show that in the limit as R tends to ∞ and r tends to zero, the integrals along the larger and smaller circle tend to zero. Hence in the limit we will get integral over the real line. Then try and relate the integral over the positive real to that over the negative reals.

Choice of contour for integration

In case the integral is from 0 to ∞ , try and relate it to some integral from $-\infty$ to ∞ . For instance the function may have a natural continuation to the negative reals. In case this is not possible, often because $f(x)$ has a singularity at origin; usually a pole, then try using a half annular region $A(0; r, R)$ like we have done in earlier examples. This will avoid the pole and then show that in the limit as R tends to ∞ and r tends to zero, the integrals along the larger and smaller circle tend to zero. Hence in the limit we will get integral over the real line. Then try and relate the integral over the positive real to that over the negative reals. See the example of $\int_0^\infty \frac{\log(x)}{1+x^2}$ as an illustration of this. Otherwise, try a rectangular contour and show that the integral over the extra sides goes to zero in the limit.

Choice of contour for integration

II. If the integrand is of the form $\int_{-\infty}^{\infty} \frac{P(x)}{Q(x)} \sin(x) dx$, where $P(x)$ and $Q(x)$ are polynomials with $\deg(Q(x))$ at least one more than that of $P(x)$, close up the interval by the semicircular region in the upper half plane and use Jordan's lemma to show that the integral over the semicircle goes to zero. (We have seen this when we integrated $\sin(x)/x$).

Choice of contour for integration

II. If the integrand is of the form $\int_{-\infty}^{\infty} \frac{P(x)}{Q(x)} \sin(x) dx$, where $P(x)$ and $Q(x)$ are polynomials with $\deg(Q(x))$ at least one more than that of $P(x)$, close up the interval by the semicircular region in the upper half plane and use Jordan's lemma to show that the integral over the semicircle goes to zero. (We have seen this when we integrated $\sin(x)/x$).

III. If the integrand is of the type $\int_0^{2\pi} P(\cos(t), \sin(t)) dt$, set $z = e^{it}$ and use $\cos(t) = \frac{z+z^{-1}}{2}$ and $\sin(t) = \frac{z-z^{-1}}{2i}$. dt becomes $\frac{dz}{iz}$ and then the integral assumes the form $\int_{|z|=1} P\left(\frac{z+z^{-1}}{2}, \frac{z-z^{-1}}{2i}\right) \frac{dz}{iz}$ which can then be computed by using residue theorem.

Choice of contour for integration

II. If the integrand is of the form $\int_{-\infty}^{\infty} \frac{P(x)}{Q(x)} \sin(x) dx$, where $P(x)$ and $Q(x)$ are polynomials with $\deg(Q(x))$ at least one more than that of $P(x)$, close up the interval by the semicircular region in the upper half plane and use Jordan's lemma to show that the integral over the semicircle goes to zero. (We have seen this when we integrated $\sin(x)/x$).

III. If the integrand is of the type $\int_0^{2\pi} P(\cos(t), \sin(t)) dt$, set $z = e^{it}$ and use $\cos(t) = \frac{z+z^{-1}}{2}$ and $\sin(t) = \frac{z-z^{-1}}{2i}$. dt becomes $\frac{dz}{iz}$ and then the integral assumes the form $\int_{|z|=1} P\left(\frac{z+z^{-1}}{2}, \frac{z-z^{-1}}{2i}\right) \frac{dz}{iz}$ which can then be computed by using residue theorem.

IV. If the integrand has infinitely many poles going to infinity, you are usually better off using a rectangular contour which encompasses only finitely many poles.

Choice of contour for integration

As before one tries to show that in the limit, the integral over the extra added vertical sides goes to zero in the limit and the integrals over the two horizontal sides are related ; usually proportional to each other. Thus taking limit as the length of the rectangular sides goes to infinity, one gets the desired answer.

Choice of contour for integration

As before one tries to show that in the limit, the integral over the extra added vertical sides goes to zero in the limit and the integrals over the two horizontal sides are related ; usually proportional to each other. Thus taking limit as the length of the rectangular sides goes to infinity, one gets the desired answer.

V. In case the function involves a branch cut, choose a contour which avoids (goes around) the branch cut like in the earlier example.

Choice of contour for integration

As before one tries to show that in the limit, the integral over the extra added vertical sides goes to zero in the limit and the integrals over the two horizontal sides are related ; usually proportional to each other. Thus taking limit as the length of the rectangular sides goes to infinity, one gets the desired answer.

V. In case the function involves a branch cut, choose a contour which avoids (goes around) the branch cut like in the earlier example.

The following rather obvious lemma is often useful for contour integration:

Choice of contour for integration

As before one tries to show that in the limit, the integral over the extra added vertical sides goes to zero in the limit and the integrals over the two horizontal sides are related ; usually proportional to each other. Thus taking limit as the length of the rectangular sides goes to infinity, one gets the desired answer.

V. In case the function involves a branch cut, choose a contour which avoids (goes around) the branch cut like in the earlier example.

The following rather obvious lemma is often useful for contour integration:

Fractional Residue Theorem : Suppose z_0 is a simple pole of $f(z)$ and C_δ is an arc of the circle $|z - z_0| = \delta$ of angle α , then

$$\lim_{\delta \rightarrow 0} \int_{C_\delta} f(z) dz = \alpha i \operatorname{Res}(f(z), z_0)$$

Argument principle

The following is a useful variant of the residue theorem and is known as the argument principle. It is used to count zero's and poles of a meromorphic function on a domain.

Argument principle

The following is a useful variant of the residue theorem and is known as the argument principle. It is used to count zero's and poles of a meromorphic function on a domain.

Argument Principle : Let γ be a simple closed contour contained in \mathbb{C} and let $f(z)$ be a meromorphic function on an open set containing γ and its interior such that γ does not pass through any of the zeros and poles of $f(z)$. Then

$$\frac{1}{2\pi i} \int_{\gamma} \frac{f'(z)}{f(z)} dz = N - P$$

where N and P denote the number of zero's and poles enclosed by γ with each zero and pole counted as many times as its order.

Argument principle

Proof: Let z_1, \dots, z_r be the set of zeros of $f(z)$ inside γ with multiplicities m_i at z_i and let p_1, \dots, p_s be the set of poles of $f(z)$ inside γ with multiplicities n_j at p_j . As in the proof of the residue theorem, we break up the integral into sum of integrals along each of the zeros and poles. Let γ_i 's be the loops around the zero's and Γ_j 's be the loops around the poles. Thus

$$\frac{1}{2\pi i} \int_{\gamma} \frac{f'(z)}{f(z)} dz = \frac{1}{2\pi i} \sum_{i=1}^r \int_{\gamma_i} \frac{f'(z)}{f(z)} dz + \frac{1}{2\pi i} \sum_{j=1}^s \int_{\Gamma_j} \frac{f'(z)}{f(z)} dz$$

Consider the zero z_i with multiplicity m_i . Then locally around z_i , $f(z) = (z - z_i)^{m_i} g(z)$ for some holomorphic function $g(z)$ with $g(z) \neq 0$ at any point inside γ_i . Then $\frac{f'(z)}{f(z)} = \frac{m_i}{z - z_i} + \frac{g'(z)}{g(z)}$. Since $g(z)$ has no zero's in γ_i , it follows that $\int_{\gamma_i} \frac{g'(z)}{g(z)} dz = 0$ by Cauchy's theorem. Hence $\int_{\gamma_i} \frac{f'(z)}{f(z)} dz = m_i$. Similar argument shows that every pole of order n_j contributes $-n_j$ to the integral. Hence we get $\frac{1}{2\pi i} \int_{\gamma} \frac{f'(z)}{f(z)} dz = N - P$ as desired.

Rouche's Theorem

A nice and useful corollary of the argument principle is the following theorem:

Rouche's Theorem

A nice and useful corollary of the argument principle is the following theorem:

Rouche's Theorem: Let γ be a simple closed contour and let $f(z)$ and $g(z)$ be two functions holomorphic on an open set containing γ and its interior. Suppose $|f(z) - g(z)| < |f(z)|$ at all points on γ . Then γ encloses the same number of zero's of $f(z)$ and $g(z)$.

Rouche's Theorem

A nice and useful corollary of the argument principle is the following theorem:

Rouche's Theorem: Let γ be a simple closed contour and let $f(z)$ and $g(z)$ be two functions holomorphic on an open set containing γ and its interior. Suppose $|f(z) - g(z)| < |f(z)|$ at all points on γ . Then γ encloses the same number of zero's of $f(z)$ and $g(z)$. The proof of Rouché's theorem follows fairly easily from the argument principle I'll leave the proof as an exercise. You can also find a proof in most of the standard books in Complex Analysis.

Remark: The hypothesis $|f(z) - g(z)| < |f(z)|$ on γ implies that f and g have no zeros on γ .

Example

Let us compute the number of zero's of

$f(z) = z^6 + 11z^4 + z^3 + 2z + 4$ inside the unit disc.

Example

Let us compute the number of zero's of

$f(z) = z^6 + 11z^4 + z^3 + 2z + 4$ inside the unit disc.

Take $g(z) = 11z^4$. Then $|g(z) - f(z)| = |z^6 + z^3 + 2z + 4| < 8$ on the unit circle. Hence $|g(z) - f(z)| < |g(z)|$ on the unit circle.

Hence $g(z)$ has the same number of roots as $f(z)$ inside the unit circle. But the number of roots of $g(z)$ inside unit circle is 4 (counting mutiplicity) which therefore equals number of roots of $f(z)$.

Example

Lets count number of roots of $f(z) = e^z - 2z - 1$ inside the unit circle.

Example

Lets count number of roots of $f(z) = e^z - 2z - 1$ inside the unit circle.

Let us consider $g(z) = -2z$. Then

$$|g(z) - f(z)| = |e^z - 1| = \left| \sum_{n=1}^{\infty} \frac{z^n}{n!} \right| \leq \sum_{n=1}^{\infty} \frac{|z^n|}{n!} = e - 1 < |g(z)|$$

on the unit circle. Hence by Rouché's theorem $f(z)$ and $g(z)$ have equal number of roots in the unit circle, namely 1.

Example

Let us determine the number of zeros of $2z^5 - 6z^2 + z + 1$ in the annulus $1 < |z| < 2$. Consider the open disc $|z| < 1$ bounded by the unit circle $C_1 : |z| = 1$. Taking $f(z) = 2z^5 - 6z^2 + z + 1$ and $g(z) = -6z^2$, we find that

$|f(z) - g(z)| = |2z^5 + z + 1| < |-6z^2| = 6$ on C_1 . Hence $f(z)$ has the same number of roots in the open unit disc as $g(z)$ which is 2.

Now consider the region $|z| < 2$ bounded by $C_2 : |z| = 2$. Then on C_2 , $|f(z) - 2z^5| = |-6z^2 + z + 1| < |27| < |2z^5| = 64$ on C_2 .

Hence $f(z)$ has the same number of roots as $2z^5$ inside C_2 . Since $2z^5$ has 5 roots counting multiplicity in the region $|z| < 2$, it follows that $f(z)$ has 5 roots inside the disc of radius 2.

Therefore $f(z)$ has 3 roots in the region $1 \leq |z| < 2$.

By the remark following Rouché's theorem, f has no zeros on $|z| = 1$ and hence $f(z)$ has 3 roots in the open annulus $1 < |z| < 2$ as well.

Here's another quick and pretty proof of FTA using Rouché's theorem.

Let $f(z) = a_0 + a_1z + \cdots + z^n$ be a non-constant polynomial. Take $g(z) = z^n$. Then on a sufficiently large circle around 0 of radius R , $|f(z) - g(z)| < |f(z)|$. Hence $f(z)$ and $g(z)$ have same number of zero's in the disc of radius R . Since $g(z)$ has n zero's, so does $f(z)$!

Harmonic Functions

We now revisit a topic we studied at the beginning of the course, namely harmonic functions. Recall that a function $u(x, y)$ of real variables is said to be harmonic if it is twice differentiable and $u_{xx} + u_{yy} = 0$. It turns out that harmonic functions share many properties similar to holomorphic functions. We'll see some of them.

Harmonic Functions

We now revisit a topic we studied at the beginning of the course, namely harmonic functions. Recall that a function $u(x, y)$ of real variables is said to be harmonic if it is twice differentiable and $u_{xx} + u_{yy} = 0$. It turns out that harmonic functions share many properties similar to holomorphic functions. We'll see some of them.

Recall that if u is a harmonic function, then a harmonic conjugate of u is another harmonic function v such that $u + iv$ is holomorphic. We saw some examples of computing harmonic conjugates and that time I commented that if the domain is “nice”, then a harmonic conjugate always exists. The mathematical notion that replaces “nice” is simply connectedness.

Harmonic Functions

We now revisit a topic we studied at the beginning of the course, namely harmonic functions. Recall that a function $u(x, y)$ of real variables is said to be harmonic if it is twice differentiable and $u_{xx} + u_{yy} = 0$. It turns out that harmonic functions share many properties similar to holomorphic functions. We'll see some of them.

Recall that if u is a harmonic function, then a harmonic conjugate of u is another harmonic function v such that $u + iv$ is holomorphic. We saw some examples of computing harmonic conjugates and that time I commented that if the domain is “nice”, then a harmonic conjugate always exists. The mathematical notion that replaces “nice” is simply connectedness.

Theorem: Let U be a simply-connected domain in \mathbb{C} and let u be a harmonic function on U . Then u admits exactly one harmonic conjugate up to a constant.

Harmonic Functions

Proof: Let's dismiss the uniqueness first. Suppose u has a harmonic conjugate v . Let $f(z) = u + iv$. By CR equations, v_x and v_y are determined and hence v is determined upto a constant. To prove existence, let $g(z) = u_x - iu_y$. Then by CR equations, $g(z)$ is holomorphic. Now fix $z_0 \in U$, and define f to be the anti-derivative of g :

Harmonic Functions

Proof: Let's dismiss the uniqueness first. Suppose u has a harmonic conjugate v . Let $f(z) = u + iv$. By CR equations, v_x and v_y are determined and hence v is determined upto a constant. To prove existence, let $g(z) = u_x - iu_y$. Then by CR equations, $g(z)$ is holomorphic. Now fix $z_0 \in U$, and define f to be the anti-derivative of g :

$f(z) = u(z_0) + \int_{z_0}^z g(z)dz$ with the integral being along a path in U connecting z_0 and z . As U is simply-connected, this function is well-defined. By construction, f is holomorphic and $f' = g = u_x - iu_y$.

Harmonic Functions

Proof: Let's dismiss the uniqueness first. Suppose u has a harmonic conjugate v . Let $f(z) = u + iv$. By CR equations, v_x and v_y are determined and hence v is determined upto a constant. To prove existence, let $g(z) = u_x - iu_y$. Then by CR equations, $g(z)$ is holomorphic. Now fix $z_0 \in U$, and define f to be the anti-derivative of g :

$f(z) = u(z_0) + \int_{z_0}^z g(z)dz$ with the integral being along a path in U connecting z_0 and z . As U is simply-connected, this function is well-defined. By construction, f is holomorphic and $f' = g = u_x - iu_y$. Letting $\tilde{u} = \operatorname{Re}(f)$, by the Cauchy-Riemann for f , $f' = \tilde{u}_x - i\tilde{u}_y$. On comparing the two equations, one concludes that $\tilde{u}_x = u_x$ and $\tilde{u}_y = u_y$.

Harmonic Functions

Proof: Let's dismiss the uniqueness first. Suppose u has a harmonic conjugate v . Let $f(z) = u + iv$. By CR equations, v_x and v_y are determined and hence v is determined upto a constant. To prove existence, let $g(z) = u_x - iu_y$. Then by CR equations, $g(z)$ is holomorphic. Now fix $z_0 \in U$, and define f to be the anti-derivative of g :

$f(z) = u(z_0) + \int_{z_0}^z g(z)dz$ with the integral being along a path in U connecting z_0 and z . As U is simply-connected, this function is well-defined. By construction, f is holomorphic and $f' = g = u_x - iu_y$. Letting $\tilde{u} = \operatorname{Re}(f)$, by the Cauchy-Riemann for f , $f' = \tilde{u}_x - i\tilde{u}_y$. On comparing the two equations, one concludes that $\tilde{u}_x = u_x$ and $\tilde{u}_y = u_y$. Therefore $u - \tilde{u}$ is a constant. Since u and \tilde{u} are equal at z_0 , they are equal throughout.

Corollary: Harmonic functions are infinitely differentiable.

This follows since on a small open disc around each point, every harmonic function admits a harmonic conjugate by simply-connectedness of open discs. Since holomorphic functions are infinitely differentiable, the corollary follows.

Harmonic functions

Let us now derive an expression analogous to the CIF.

Harmonic functions

Let us now derive an expression analogous to the CIF.

Theorem: (Mean-Value Property): Let u be a harmonic function on a disc of radius R . Then for any $r < R$, we have,

$$u(w) = \frac{1}{2\pi} \int_0^{2\pi} u(w + re^{i\theta}) d\theta.$$

In particular, u does not attain its maximum at any interior point unless it is constant.

Harmonic functions

Let us now derive an expression analogous to the CIF.

Theorem: (Mean-Value Property): Let u be a harmonic function on a disc of radius R . Then for any $r < R$, we have,

$$u(w) = \frac{1}{2\pi} \int_0^{2\pi} u(w + re^{i\theta}) d\theta.$$

In particular, u does not attain its maximum at any interior point unless it is constant.

Proof: Can assume $w = 0$ without loss of generality. Since u is harmonic and the domain is simply-connected, there exists a holomorphic function $f(z)$ such that $\operatorname{Re}(f) = u$. By Cauchy's integral formula,

$$f(0) = \frac{1}{2\pi i} \int_{|z|=r} \frac{f(z)}{z} dz$$

Hence

$$f(0) = \frac{1}{2\pi} \int_0^{2\pi} f(re^{i\theta}) d\theta$$

Identity Principle

Taking real parts on both sides, we get the desired result.

Identity Principle

Taking real parts on both sides, we get the desired result.

Note that this in particular implies that if u is a harmonic function on a domain Ω and $z \in \Omega$. Suppose $u(z) \geq u(w)$ for all w in a neighborhood of z , then u is a constant. By considering $-u$ instead of u , the same holds for \geq replaced by \leq .

Identity Principle

Taking real parts on both sides, we get the desired result.

Note that this in particular implies that if u is a harmonic function on a domain Ω and $z \in \Omega$. Suppose $u(z) \geq u(w)$ for all w in a neighborhood of z , then u is a constant. By considering $-u$ instead of u , the same holds for \geq replaced by \leq .

Identity Principle: Let u be a harmonic function on a domain $\Omega \subset \mathbb{C}$. If $u = 0$ on a non-empty open subset $U \subseteq \Omega$, then $u = 0$ throughout Ω .

Identity Principle

Taking real parts on both sides, we get the desired result.

Note that this in particular implies that if u is a harmonic function on a domain Ω and $z \in \Omega$. Suppose $u(z) \geq u(w)$ for all w in a neighborhood of z , then u is a constant. By considering $-u$ instead of u , the same holds for \geq replaced by \leq .

Identity Principle: Let u be a harmonic function on a domain $\Omega \subset \mathbb{C}$. If $u = 0$ on a non-empty open subset $U \subseteq \Omega$, then $u = 0$ throughout Ω .

Proof: Set $f = u_x - iu_y$. Then as before, f is holomorphic on Ω . Since $u = 0$ on U then so is f . Hence, by the Identity Principle for the holomorphic functions $f = 0$ on Ω , and consequently, $u_x = u_y = 0$ on Ω . Therefore u is constant on Ω , and as it is zero on U , it must be zero on Ω .

Remark: Recall that the identity theorem for holomorphic functions is stronger; namely if a holomorphic function vanishes on a set of points having a limit point, then it is identically zero. This is not true for harmonic functions. The function $Re(z)$ vanishes identically on imaginary axis but is non-zero elsewhere.