

Signal Processing - 1 by One

Sibi Raj B. Pillai
Dept of Electrical Engineering
IIT Bombay




- So Far: Impulse, Sampling, Replacement
- Previous Week: Convolution ($*$) and Interpolation
- Previous Class: Fourier Series
- Today: Fourier Series Contd..




- So Far: Impulse, Sampling, Replacement
- Previous Week: Convolution ($*$) and Interpolation
- Previous Class: Fourier Series
- Today: Fourier Series Contd..




Frequency of Square Waves



Frequency of Square Waves




Frequency of Square Waves



Interval of interest $\mathbb{T} = [-\frac{T}{2}, +\frac{T}{2}]$.



Frequency of Square Waves




Interval of interest $\mathbb{T} = [-\frac{T}{2}, +\frac{T}{2}]$.

$$\alpha_0 = \frac{1}{T} \int_{-\frac{T}{2}}^{\frac{T}{2}} f(t) dt = \frac{\tau}{T}$$



Frequency of Square Waves




Interval of interest $\mathbb{T} = [-\frac{T}{2}, +\frac{T}{2}]$.

$$\alpha_0 = \frac{1}{T} \int_{-\frac{T}{2}}^{\frac{T}{2}} f(t) dt = \frac{\tau}{T}$$

$$b_n = \frac{2}{T} \int_{-\frac{T}{2}}^{\frac{T}{2}} f(t) \sin\left(\frac{2\pi}{T}nt\right) dt = 0 \quad (\text{odd function})$$



Frequency of Square Waves



Interval of interest $\mathbb{T} = [-\frac{T}{2}, +\frac{T}{2}]$.


$$\alpha_0 = \frac{1}{T} \int_{-\frac{T}{2}}^{\frac{T}{2}} f(t) dt = \frac{\tau}{T}$$

$$b_n = \frac{2}{T} \int_{-\frac{T}{2}}^{\frac{T}{2}} f(t) \sin\left(\frac{2\pi}{T} nt\right) dt = 0 \quad (\text{odd function})$$

$$a_n = \frac{2}{T} \int_{-\frac{T}{2}}^{\frac{T}{2}} f(t) \cos\left(\frac{2\pi}{T} nt\right) dt = \frac{2}{n\pi} \sin\left(n\pi \frac{\tau}{T}\right)$$



Frequency of Square Waves



Interval of interest $\mathbb{T} = [-\frac{T}{2}, +\frac{T}{2}]$.

$$\alpha_0 = \frac{1}{T} \int_{-\frac{T}{2}}^{\frac{T}{2}} f(t) dt = \frac{\tau}{T}$$

$$b_n = \frac{2}{T} \int_{-\frac{T}{2}}^{\frac{T}{2}} f(t) \sin\left(\frac{2\pi}{T} nt\right) dt = 0 \quad (\text{odd function})$$

$$a_n = \frac{2}{T} \int_{-\frac{T}{2}}^{\frac{T}{2}} f(t) \cos\left(\frac{2\pi}{T} nt\right) dt = \frac{2}{n\pi} \sin\left(n\pi \frac{\tau}{T}\right) = 2\frac{\tau}{T} \text{sinc}\left(n\frac{\tau}{T}\right) \text{ for } (n \neq 0).$$



Fourier Series

$$\begin{aligned}f(t) &= \alpha_0 + \sum_{n \in \mathbb{Z} \setminus \{0\}} a_n \cos\left(\frac{2\pi}{T_d} nt\right) + \sum_{n \in \mathbb{Z}} b_n \sin\left(\frac{2\pi}{T_d} nt\right) \\&= \alpha_0 + \sum_{n \in \mathbb{Z} \setminus \{0\}} a_n \frac{\left(e^{j\frac{2\pi}{T_d} nt} + e^{-j\frac{2\pi}{T_d} nt}\right)}{2} + \sum_{n \in \mathbb{Z}} b_n \frac{\left(e^{j\frac{2\pi}{T_d} nt} - e^{-j\frac{2\pi}{T_d} nt}\right)}{2j} \\&= \sum_{m \in \mathbb{Z}} \alpha_m e^{j\frac{2\pi}{T_d} mt}.\end{aligned}$$



Fourier Series

$$\begin{aligned}f(t) &= \alpha_0 + \sum_{n \in \mathbb{Z} \setminus \{0\}} a_n \cos\left(\frac{2\pi}{T_d} nt\right) + \sum_{n \in \mathbb{Z}} b_n \sin\left(\frac{2\pi}{T_d} nt\right) \\&= \alpha_0 + \sum_{n \in \mathbb{Z} \setminus \{0\}} a_n \frac{\left(e^{j\frac{2\pi}{T_d} nt} + e^{-j\frac{2\pi}{T_d} nt}\right)}{2} + \sum_{n \in \mathbb{Z}} b_n \frac{\left(e^{j\frac{2\pi}{T_d} nt} - e^{-j\frac{2\pi}{T_d} nt}\right)}{2j} \\&= \sum_{m \in \mathbb{Z}} \alpha_m e^{j\frac{2\pi}{T_d} mt}.\end{aligned}$$

$$\alpha_m = \frac{a_m - jb_m}{2}, \quad m \in \mathbb{Z}^+ \quad \text{and} \quad \alpha_m = \frac{a_m + jb_m}{2}, \quad m \in \mathbb{Z}^-.$$



Fourier Series

$$\begin{aligned}f(t) &= \alpha_0 + \sum_{n \in \mathbb{Z} \setminus \{0\}} a_n \cos\left(\frac{2\pi}{T_d} nt\right) + \sum_{n \in \mathbb{Z}} b_n \sin\left(\frac{2\pi}{T_d} nt\right) \\&= \alpha_0 + \sum_{n \in \mathbb{Z} \setminus \{0\}} a_n \frac{\left(e^{j\frac{2\pi}{T_d} nt} + e^{-j\frac{2\pi}{T_d} nt}\right)}{2} + \sum_{n \in \mathbb{Z}} b_n \frac{\left(e^{j\frac{2\pi}{T_d} nt} - e^{-j\frac{2\pi}{T_d} nt}\right)}{2j} \\&= \sum_{m \in \mathbb{Z}} \alpha_m e^{j\frac{2\pi}{T_d} mt}.\end{aligned}$$

$$\alpha_m = \frac{a_m - jb_m}{2}, \quad m \in \mathbb{Z}^+ \quad \text{and} \quad \alpha_m = \frac{a_m + jb_m}{2}, \quad m \in \mathbb{Z}^-.$$

$$\alpha_m = \frac{\langle f(t), \exp(j\frac{2\pi}{T_d} mt) \rangle}{T_d} = \frac{1}{T_d} \int_{-\frac{T_d}{2}}^{\frac{T_d}{2}} f(t) \exp(-j\frac{2\pi}{T_d} mt) dt.$$



Fourier Series

$$\begin{aligned}f(t) &= \alpha_0 + \sum_{n \in \mathbb{Z} \setminus \{0\}} a_n \cos\left(\frac{2\pi}{T_d} nt\right) + \sum_{n \in \mathbb{Z}} b_n \sin\left(\frac{2\pi}{T_d} nt\right) \\&= \alpha_0 + \sum_{n \in \mathbb{Z} \setminus \{0\}} a_n \frac{\left(e^{j\frac{2\pi}{T_d} nt} + e^{-j\frac{2\pi}{T_d} nt}\right)}{2} + \sum_{n \in \mathbb{Z}} b_n \frac{\left(e^{j\frac{2\pi}{T_d} nt} - e^{-j\frac{2\pi}{T_d} nt}\right)}{2j} \\&= \sum_{m \in \mathbb{Z}} \alpha_m e^{j\frac{2\pi}{T_d} mt}.\end{aligned}$$

“Why is this True?”

$$\alpha_m = \frac{a_m - jb_m}{2}, \quad m \in \mathbb{Z}^+ \quad \text{and} \quad \alpha_m = \frac{a_m + jb_m}{2}, \quad m \in \mathbb{Z}^-.$$

$$\alpha_m = \frac{\langle f(t), \exp(j\frac{2\pi}{T_d} mt) \rangle}{T_d} = \frac{1}{T_d} \int_{-\frac{T_d}{2}}^{\frac{T_d}{2}} f(t) \exp(-j\frac{2\pi}{T_d} mt) dt.$$



Convergence

Proposition

No two continuous functions in $[-\frac{T_d}{2}, \frac{T_d}{2}]$ will have all the Fourier Series coefficients the same.



Proposition

No two continuous functions in $[-\frac{T_d}{2}, \frac{T_d}{2}]$ will have all the Fourier Series coefficients the same.

Lemma

Let $f(t)$ be a signal locally integrable in $[-\frac{T}{2}, \frac{T}{2}]$.

If the FS coefficients $\alpha_m = 0$, identically $\forall m \in \mathbb{Z}$, then $f(t) = 0$, whenever it is continuous at t .



Proposition

No two continuous functions in $[-\frac{T_d}{2}, \frac{T_d}{2}]$ will have all the Fourier Series coefficients the same.

Lemma

Let $f(t)$ be a signal locally integrable in $[-\frac{T}{2}, \frac{T}{2}]$.

If the FS coefficients $\alpha_m = 0$, identically $\forall m \in \mathbb{Z}$, then $f(t) = 0$, whenever it is continuous at t .

Proof Idea: Reduction or Contradiction



Proposition

No two continuous functions in $[-\frac{T_d}{2}, \frac{T_d}{2}]$ will have all the Fourier Series coefficients the same.

Lemma

Let $f(t)$ be a signal locally integrable in $[-\frac{T}{2}, \frac{T}{2}]$.

If the FS coefficients $\alpha_m = 0$, identically $\forall m \in \mathbb{Z}$, then $f(t) = 0$, whenever it is continuous at t .

Proof Idea: Reduction or Contradiction

Suppose FS coefficients $\alpha_m = 0$ for all $m \in \mathbb{Z}$ and $f(t)$ is continuous at $t = 0$, with $f(0) > 0$.



Proposition

No two continuous functions in $[-\frac{T_d}{2}, \frac{T_d}{2}]$ will have all the Fourier Series coefficients the same.

Lemma

Let $f(t)$ be a signal locally integrable in $[-\frac{T}{2}, \frac{T}{2}]$.

If the FS coefficients $\alpha_m = 0$, identically $\forall m \in \mathbb{Z}$, then $f(t) = 0$, whenever it is continuous at t .

Proof Idea: Reduction or Contradiction

Suppose FS coefficients $\alpha_m = 0$ for all $m \in \mathbb{Z}$ and $f(t)$ is continuous at $t = 0$, with $f(0) > 0$. **We will show this is ABSURD!**



Proof [Stein-Schakarchi]

From our assumption of FS coefficients being zero, i.e.

$$\frac{1}{T} \int_{\mathbb{T}} f(t) \exp(-j \frac{2\pi}{T} m t) dt = 0, \forall m \in \mathbb{Z}$$

Furthermore

$$\int_{\mathbb{T}} f(t) (P(t))^k dt = 0, \forall k = 0, 1, 2, \dots$$

Take for some parameter $\epsilon > 0$,

$$P(t) = \epsilon + \cos\left(\frac{2\pi}{T} t\right).$$



Proof [Stein-Schakarchi]

From our assumption of FS coefficients being zero, i.e.

$$\frac{1}{T} \int_{\mathbb{T}} f(t) \exp(-j \frac{2\pi}{T} m t) dt = 0, \forall m \in \mathbb{Z} \Rightarrow \int_{\mathbb{T}} f(t) P(t) dt = 0,$$

for any trigonometric polynomial $P(t)$ of the form

$$P(t) = c_0 + c_1 \exp(-j \frac{2\pi}{T} t) + c_2 \exp(j \frac{2\pi}{T} t).$$

Furthermore

$$\int_{\mathbb{T}} f(t) (P(t))^k dt = 0, \forall k = 0, 1, 2, \dots$$

Take for some parameter $\epsilon > 0$,

$$P(t) = \epsilon + \cos\left(\frac{2\pi}{T} t\right).$$



Proof [Stein-Schakarchi]

From our assumption of FS coefficients being zero, i.e.

$$\frac{1}{T} \int_{\mathbb{T}} f(t) \exp(-j \frac{2\pi}{T} m t) dt = 0, \forall m \in \mathbb{Z} \Rightarrow \int_{\mathbb{T}} f(t) P(t) dt = 0,$$

for any trigonometric polynomial $P(t)$ of the form

$$P(t) = c_0 + c_1 \exp(-j \frac{2\pi}{T} t) + c_2 \exp(j \frac{2\pi}{T} t).$$

Furthermore

$$\int_{\mathbb{T}} f(t) (P(t))^k dt = 0, \forall k = 0, 1, 2, \dots$$

Take for some parameter $\epsilon > 0$,

$$P(t) = \epsilon + \cos\left(\frac{2\pi}{T} t\right).$$



Proof [Stein-Schakarchi]

From our assumption of FS coefficients being zero, i.e.

$$\frac{1}{T} \int_{\mathbb{T}} f(t) \exp(-j \frac{2\pi}{T} m t) dt = 0, \forall m \in \mathbb{Z} \Rightarrow \int_{\mathbb{T}} f(t) P(t) dt = 0,$$

for any trigonometric polynomial $P(t)$ of the form

$$P(t) = c_0 + c_1 \exp(-j \frac{2\pi}{T} t) + c_2 \exp(j \frac{2\pi}{T} t).$$

Furthermore


$$\int_{\mathbb{T}} f(t) (P(t))^k dt = 0, \forall k = 0, 1, 2, \dots$$

Take for some parameter $\epsilon > 0$,


$$P(t) = \epsilon + \cos\left(\frac{2\pi}{T} t\right).$$




Proof in Picture




Proof in Picture



Proof in Picture




Proof in Picture




$$\epsilon + \cos \frac{2\pi}{T} \delta = 1$$




Proof in Picture




Proof in Picture




Proof in Picture



Proof in Picture



Proof in Picture



$$\begin{aligned}\int f(t)P^k(t)dt &= \int_{\delta \leq |t| \leq \frac{T}{2}} f(t)P^k(t)dt \\ &\quad + \int_{\nu \leq |t| \leq \delta} f(t)P^k(t)dt \\ &\quad + \int_{|t| \leq \nu} f(t)P^k(t)dt\end{aligned}$$

1. $I_1 > -\infty$ as $f(t)$ is locally integrable.
2. $I_2 > 0$.
3. $I_3 \geq \frac{f(0)}{2}(1 + \frac{\epsilon}{2})^k \rightarrow \infty$ when $k \uparrow \infty$, as $\epsilon > 0$.



$$\begin{aligned}\int f(t)P^k(t)dt &= \int_{\delta \leq |t| \leq \frac{T}{2}} f(t)P^k(t)dt \\ &\quad + \int_{\nu \leq |t| \leq \delta} f(t)P^k(t)dt \\ &\quad + \int_{|t| \leq \nu} f(t)P^k(t)dt \\ &= l_1 + l_2 + l_3.\end{aligned}$$

1. $l_1 > -\infty$ as $f(t)$ is locally integrable.
2. $l_2 > 0$.
3. $l_3 \geq \frac{f(0)}{2}(1 + \frac{\epsilon}{2})^k \rightarrow \infty$ when $k \uparrow \infty$, as $\epsilon > 0$.



$$\begin{aligned}\int f(t)P^k(t)dt &= \int_{\delta \leq |t| \leq \frac{T}{2}} f(t)P^k(t)dt \\ &\quad + \int_{\nu \leq |t| \leq \delta} f(t)P^k(t)dt \\ &\quad + \int_{|t| \leq \nu} f(t)P^k(t)dt \\ &= I_1 + I_2 + I_3.\end{aligned}$$

1. $I_1 > -\infty$ as $f(t)$ is locally integrable.
2. $I_2 > 0$.
3. $I_3 \geq \frac{f(0)}{2}(1 + \frac{\epsilon}{2})^k \rightarrow \infty$ when $k \uparrow \infty$, as $\epsilon > 0$.



$$\begin{aligned}\int f(t)P^k(t)dt &= \int_{\delta \leq |t| \leq \frac{T}{2}} f(t)P^k(t)dt \\ &\quad + \int_{\nu \leq |t| \leq \delta} f(t)P^k(t)dt \\ &\quad + \int_{|t| \leq \nu} f(t)P^k(t)dt \\ &= I_1 + I_2 + I_3.\end{aligned}$$

1. $I_1 > -\infty$ as $f(t)$ is locally integrable.
2. $I_2 > 0$.
3. $I_3 \geq \frac{f(0)}{2}(1 + \frac{\epsilon}{2})^k \rightarrow \infty$ when $k \uparrow \infty$, as $\epsilon > 0$.



$$\begin{aligned}\int f(t)P^k(t)dt &= \int_{\delta \leq |t| \leq \frac{T}{2}} f(t)P^k(t)dt \\ &\quad + \int_{\nu \leq |t| \leq \delta} f(t)P^k(t)dt \\ &\quad + \int_{|t| \leq \nu} f(t)P^k(t)dt \\ &= l_1 + l_2 + l_3.\end{aligned}$$

1. $l_1 > -\infty$ as $f(t)$ is locally integrable.
2. $l_2 > 0$.
3. $l_3 \geq \frac{f(0)}{2}(1 + \frac{\epsilon}{2})^k \rightarrow \infty$ when $k \uparrow \infty$, as $\epsilon > 0$.



$$\begin{aligned}\int f(t)P^k(t)dt &= \int_{\delta \leq |t| \leq \frac{T}{2}} f(t)P^k(t)dt \\ &\quad + \int_{\nu \leq |t| \leq \delta} f(t)P^k(t)dt \\ &\quad + \int_{|t| \leq \nu} f(t)P^k(t)dt \\ &= l_1 + l_2 + l_3.\end{aligned}$$

1. $l_1 > -\infty$ as $f(t)$ is locally integrable.
2. $l_2 > 0$.
3. $l_3 \geq \frac{f(0)}{2}(1 + \frac{\epsilon}{2})^k \rightarrow \infty$ when $k \uparrow \infty$, as $\epsilon > 0$.

We have our contradiction, thus $f(0)$ has to be zero.



What this was about!

Assume a function $f(t)$ integrable in $[-\frac{T}{2}, \frac{T}{2}]$, and

$$\alpha_m = \frac{1}{T} \int_{-\frac{T}{2}}^{\frac{T}{2}} f(t) \exp(-j\frac{2\pi}{T}mt) dt, m \in \mathbb{Z}.$$



What this was about!

Assume a function $f(t)$ integrable in $[-\frac{T}{2}, \frac{T}{2}]$, and

$$\alpha_m = \frac{1}{T} \int_{-\frac{T}{2}}^{\frac{T}{2}} f(t) \exp(-j\frac{2\pi}{T}mt) dt, m \in \mathbb{Z}.$$

Now consider the function

$$g(t) = \sum_{n \in \mathbb{Z}} \alpha_n \exp(j\frac{2\pi}{T}nt) dt.$$

Clearly, $g(t)$ also has Fourier Series coefficients $\alpha_n, n \in \mathbb{Z}$.



What this was about!

Assume a function $f(t)$ integrable in $[-\frac{T}{2}, \frac{T}{2}]$, and

$$\alpha_m = \frac{1}{T} \int_{-\frac{T}{2}}^{\frac{T}{2}} f(t) \exp(-j\frac{2\pi}{T}mt) dt, m \in \mathbb{Z}.$$

Now consider the function

$$g(t) = \sum_{n \in \mathbb{Z}} \alpha_n \exp(j\frac{2\pi}{T}nt) dt.$$

Clearly, $g(t)$ also has Fourier Series coefficients $\alpha_n, n \in \mathbb{Z}$.

Our lemma says $g(t) = f(t)$, if $f(t)$ is continuous at t . Therefore,

$$\sum_{n \in \mathbb{Z}} \alpha_n \exp\left(j\frac{2\pi}{T}nt\right) dt = f(t)$$

at instants where $f(t)$ is continuous.

