

MA 205 Complex Analysis: Examples

September 11, 2020

Correction from Lecture 7

Let me begin with a correction from lecture 7.

Correction from Lecture 7

Let me begin with a correction from lecture 7. I stated that if Ω is an open subset of \mathbb{C} and f a holomorphic function on Ω , then the radius of convergence of the power series expanded around a point $z_0 \in \Omega$ is the radius of the largest circle around z_0 contained in Ω .

Correction from Lecture 7

Let me begin with a correction from lecture 7. I stated that if Ω is an open subset of \mathbb{C} and f a holomorphic function on Ω , then the radius of convergence of the power series expanded around a point $z_0 \in \Omega$ is the radius of the largest circle around z_0 contained in Ω . Really what I meant is that the radius of convergence of the power series expanded around z_0 is the radius of the largest circle centered at z_0 on which f extends as a holomorphic function.

Correction from Lecture 7

Let me begin with a correction from lecture 7. I stated that if Ω is an open subset of \mathbb{C} and f a holomorphic function on Ω , then the radius of convergence of the power series expanded around a point $z_0 \in \Omega$ is the radius of the largest circle around z_0 contained in Ω . Really what I meant is that the radius of convergence of the power series expanded around z_0 is the radius of the largest circle centered at z_0 on which f extends as a holomorphic function. The largest circle centered at z_0 contained in Ω may be smaller.

Example

Consider $f(z) = \frac{e^z}{\sin z + \cos z}$. Let's expand it as a power series centered at 0. First note that f is holomorphic at 0. The nearest pole to 0 for this function is when $\sin z + \cos z = 0$. On squaring, when this happens $\sin(2z) = -1$. Lets understand when this happens.

Example

Consider $f(z) = \frac{e^z}{\sin z + \cos z}$. Let's expand it as a power series centered at 0. First note that f is holomorphic at 0. The nearest pole to 0 for this function is when $\sin z + \cos z = 0$. On squaring, when this happens $\sin(2z) = -1$. Lets understand when this happens.

We use the fact that $\sin(x + iy) = \sin x \cosh y + i \cos x \sinh y$.
(Exercise !)

Example

Consider $f(z) = \frac{e^z}{\sin z + \cos z}$. Let's expand it as a power series centered at 0. First note that f is holomorphic at 0. The nearest pole to 0 for this function is when $\sin z + \cos z = 0$. On squaring, when this happens $\sin(2z) = -1$. Let's understand when this happens.

We use the fact that $\sin(x + iy) = \sin x \cosh y + i \cos x \sinh y$. (Exercise !)

Thus $\sin(x + iy) = -1$ when $\sin x \cosh y = -1$ and $\cos x \sinh y = 0$. Thus either $\cos x = 0$ or $\sinh y = 0$. If $\cos x = 0$, then $x = \pm \frac{\pi}{2} \pm n\pi$.

Example

Consider $f(z) = \frac{e^z}{\sin z + \cos z}$. Let's expand it as a power series centered at 0. First note that f is holomorphic at 0. The nearest pole to 0 for this function is when $\sin z + \cos z = 0$. On squaring, when this happens $\sin(2z) = -1$. Lets understand when this happens.

We use the fact that $\sin(x + iy) = \sin x \cosh y + i \cos x \sinh y$. (Exercise !)

Thus $\sin(x + iy) = -1$ when $\sin x \cosh y = -1$ and $\cos x \sinh y = 0$. Thus either $\cos x = 0$ or $\sinh y = 0$. If $\cos x = 0$, then $x = \pm \frac{\pi}{2} \pm n\pi$. We are interested in the nearest value to 0. When $x = \pi/2$, $\sin x = 1$ and so $\sin x \cosh y > 0$ and so cannot be -1 . When $x = -\pi/2$ and $y = 0$, then $\cos x \sinh y = 0$ and $\sin x \cosh y = -1$ are both satisfied and so this is the nearest value to 0 for which $\sin(x + iy) = -1$.

Example

Consider $f(z) = \frac{e^z}{\sin z + \cos z}$. Let's expand it as a power series centered at 0. First note that f is holomorphic at 0. The nearest pole to 0 for this function is when $\sin z + \cos z = 0$. On squaring, when this happens $\sin(2z) = -1$. Let's understand when this happens.

We use the fact that $\sin(x + iy) = \sin x \cosh y + i \cos x \sinh y$. (Exercise!) Thus $\sin(x + iy) = -1$ when $\sin x \cosh y = -1$ and $\cos x \sinh y = 0$. Thus either $\cos x = 0$ or $\sinh y = 0$. If $\cos x = 0$, then $x = \pm \frac{\pi}{2} \pm n\pi$. We are interested in the nearest value to 0. When $x = \pi/2$, $\sin x = 1$ and so $\sin x \cosh y > 0$ and so cannot be -1 . When $x = -\pi/2$ and $y = 0$, then $\cos x \sinh y = 0$ and $\sin x \cosh y = -1$ are both satisfied and so this is the nearest value to 0 for which $\sin(x + iy) = -1$. The analysis when $\sinh y = 0$ gives the same result. Thus the nearest pole for the function is when $\sin(2z) = -1$ and the smallest value of z for which this happens is when $z = -\pi/4$.

Example

Consider $f(z) = \frac{e^z}{\sin z + \cos z}$. Let's expand it as a power series centered at 0. First note that f is holomorphic at 0. The nearest pole to 0 for this function is when $\sin z + \cos z = 0$. On squaring, when this happens $\sin(2z) = -1$. Let's understand when this happens.

We use the fact that $\sin(x + iy) = \sin x \cosh y + i \cos x \sinh y$. (Exercise !). Thus $\sin(x + iy) = -1$ when $\sin x \cosh y = -1$ and $\cos x \sinh y = 0$. Thus either $\cos x = 0$ or $\sinh y = 0$. If $\cos x = 0$, then $x = \pm \frac{\pi}{2} \pm n\pi$. We are interested in the nearest value to 0. When $x = \pi/2$, $\sin x = 1$ and so $\sin x \cosh y > 0$ and so cannot be -1 . When $x = -\pi/2$ and $y = 0$, then $\cos x \sinh y = 0$ and $\sin x \cosh y = -1$ are both satisfied and so this is the nearest value to 0 for which $\sin(x + iy) = -1$. The analysis when $\sinh y = 0$ gives the same result. Thus the nearest pole for the function is when $\sin(2z) = -1$ and the smallest value of z for which this happens is when $z = -\pi/4$.

Hence radius of convergence is $\pi/4$.

Definition

Definition: If $z_0 \in \mathbb{C}$ is any point and γ is any closed contour not passing through z_0 , then $\int_{\gamma} \frac{1}{z-z_0} dz$ is an integer multiple of $2\pi i$. This integer is (suggestively) called the **winding number** of γ around z_0 and counts the number of times the curves winds around z_0 . Note that this integer could be negative which happens when the curve winds around in clockwise orientation. If γ is a finite union of simple closed curves and z_0 lies outside γ (which makes sense by Jordan curve theorem), then this integer is zero. If γ is a simple closed curve, and z_0 lies in the interior of γ then this integer is 0.

Let us now see some computational applications of Cauchy Integral Formula.

Definition

Definition: If $z_0 \in \mathbb{C}$ is any point and γ is any closed contour not passing through z_0 , then $\int_{\gamma} \frac{1}{z-z_0} dz$ is an integer multiple of $2\pi i$. This integer is (suggestively) called the **winding number** of γ around z_0 and counts the number of times the curves winds around z_0 . Note that this integer could be negative which happens when the curve winds around in clockwise orientation. If γ is a finite union of simple closed curves and z_0 lies outside γ (which makes sense by Jordan curve theorem), then this integer is zero. If γ is a simple closed curve, and z_0 lies in the interior of γ then this integer is 0.

Let us now see some computational applications of Cauchy Integral Formula.

Recall that if Ω is a domain in \mathbb{C} and f is a holomorphic function on and inside a simply closed contour γ and z_0 is an interior point of γ , then

$$f^{(n)}(z_0) = \frac{n!}{2\pi i} \int \frac{f(z)}{(z-z_0)^{n+1}} dz$$

Definition

Definition: If $z_0 \in \mathbb{C}$ is any point and γ is any closed contour not passing through z_0 , then $\int_{\gamma} \frac{1}{z-z_0} dz$ is an integer multiple of $2\pi i$. This integer is (suggestively) called the **winding number** of γ around z_0 and counts the number of times the curves winds around z_0 . Note that this integer could be negative which happens when the curve winds around in clockwise orientation. If γ is a finite union of simple closed curves and z_0 lies outside γ (which makes sense by Jordan curve theorem), then this integer is zero. If γ is a simple closed curve, and z_0 lies in the interior of γ then this integer is 0.

Let us now see some computational applications of Cauchy Integral Formula.

Recall that if Ω is a domain in \mathbb{C} and f is a holomorphic function on and inside a simply closed contour γ and z_0 is an interior point of γ , then

$$f^{(n)}(z_0) = \frac{n!}{2\pi i} \int \frac{f(z)}{(z-z_0)^{n+1}} dz$$

Example 1:

$$\begin{aligned} \int_{|z|=2} \frac{z^2}{z-1} dz \\ = 2\pi i [z^2]_{z=1} = 2\pi i \end{aligned}$$

Example 1:

$$\begin{aligned} & \int_{|z|=2} \frac{z^2}{z-1} dz \\ &= 2\pi i [z^2]_{z=1} = 2\pi i \end{aligned}$$

Example 2:

$$\begin{aligned} & \int_{|z|=2} \frac{e^z}{z^2(z-1)} dz \\ &= \int_{|z|=\epsilon} \frac{e^z/z-1}{z^2} + \int_{|z-1|=\epsilon} \frac{e^z/z^2}{z-1} dz \\ &= 2\pi i \left[\frac{d}{dz} \left(\frac{e^z}{z-1} \right) \right]_{z=0} + 2\pi i \left[\frac{e^z}{z^2} \right]_{z=1} \\ &= -4\pi i + (2\pi i)e = 2\pi i(e - 2) \end{aligned}$$

Example 3

$$\int_{|z-1|=1} \frac{z^2-4z+3}{z^2-z-1} dz$$

Example 3

$$\int_{|z-1|=1} \frac{z^2-4z+3}{z^2-z-1} dz$$

$$= \int_{|z-1|=1} \frac{z^2-z-1-3z+4}{z^2-z-1} dz$$

Example 3

$$\int_{|z-1|=1} \frac{z^2-4z+3}{z^2-z-1} dz$$

$$= \int_{|z-1|=1} \frac{z^2-z-1-3z+4}{z^2-z-1} dz$$

$$= \int_{|z-1|=1} \left[1 - \frac{3z-4}{z^2-z-1} \right] dz$$

Example 3

$$\int_{|z-1|=1} \frac{z^2-4z+3}{z^2-z-1} dz$$

$$= \int_{|z-1|=1} \frac{z^2-z-1-3z+4}{z^2-z-1} dz$$

$$= \int_{|z-1|=1} \left[1 - \frac{3z-4}{z^2-z-1} \right] dz$$

$$= \int_{|z-1|=1} 1 dz - \int_{|z-1|=1} \frac{3z-4}{(z-\frac{1+\sqrt{5}}{2})(z-\frac{1-\sqrt{5}}{2})} dz$$

Example 3

$$\begin{aligned} & \int_{|z-1|=1} \frac{z^2-4z+3}{z^2-z-1} dz \\ &= \int_{|z-1|=1} \frac{z^2-z-1-3z+4}{z^2-z-1} dz \\ &= \int_{|z-1|=1} \left[1 - \frac{3z-4}{z^2-z-1} \right] dz \\ &= \int_{|z-1|=1} 1 dz - \int_{|z-1|=1} \frac{3z-4}{(z-\frac{1+\sqrt{5}}{2})(z-\frac{1-\sqrt{5}}{2})} \\ &= 0 - \int_{|z-1|=1} \frac{\left(\frac{3z-4}{z-\frac{1-\sqrt{5}}{2}} \right)}{(z-\frac{1+\sqrt{5}}{2})} \end{aligned}$$

Example 3

$$\begin{aligned} & \int_{|z-1|=1} \frac{z^2-4z+3}{z^2-z-1} dz \\ &= \int_{|z-1|=1} \frac{z^2-z-1-3z+4}{z^2-z-1} dz \\ &= \int_{|z-1|=1} \left[1 - \frac{3z-4}{z^2-z-1} \right] dz \\ &= \int_{|z-1|=1} 1 dz - \int_{|z-1|=1} \frac{3z-4}{(z-\frac{1+\sqrt{5}}{2})(z-\frac{1-\sqrt{5}}{2})} \\ &= 0 - \int_{|z-1|=1} \frac{\left(\frac{3z-4}{z-\frac{1-\sqrt{5}}{2}} \right)}{(z-\frac{1+\sqrt{5}}{2})} \\ &= 0 - 2\pi i \left[(3z-4)/(z-(1-\sqrt{5})/2) \right]_{z=(1+\sqrt{5})/2} \\ &= (\sqrt{5}-3)\pi i \end{aligned}$$

Example 4

$$\begin{aligned} I &= \int_{|z|=3} \frac{z^9+1}{z^6-1} \\ &= \int_{|z|=3} \frac{z^3(z^6-1)+z^3+1}{z^6-1} \\ &= \int_{|z|=3} z^3 + \int_{|z|=3} \frac{z^3+1}{z^6-1} \\ &= 0 + \int_{|z|=3} \frac{1}{z^3-1} \\ &= 0 \text{ (by an earlier exercise)} \end{aligned}$$

Example 5

Lets compute $\int_{|z|=1} \frac{dz}{z(z^3+3z-7)}.$

Example 5

Lets compute $\int_{|z|=1} \frac{dz}{z(z^3+3z-7)}$. This is an example where we can estimate the location of poles without actually locating them precisely.

Example 5

Lets compute $\int_{|z|=1} \frac{dz}{z(z^3+3z-7)}$. This is an example where we can estimate the location of poles without actually locating them precisely. The function $z^3 + 3z - 7$ restricted to real axis is an odd degree polynomial and so it has either 3 real roots or 1 real root and 2 complex conjugate roots.

Example 5

Lets compute $\int_{|z|=1} \frac{dz}{z(z^3+3z-7)}$. This is an example where we can estimate the location of poles without actually locating them precisely. The function $z^3 + 3z - 7$ restricted to real axis is an odd degree polynomial and so it has either 3 real roots or 1 real root and 2 complex conjugate roots. Now just by observation , at $z = 1$, this function is -3 and at $z=2$, its value is 7 and so by intermediate value theorem it has a real root between 1 and 2.

Example 5

Lets compute $\int_{|z|=1} \frac{dz}{z(z^3+3z-7)}$. This is an example where we can estimate the location of poles without actually locating them precisely. The function $z^3 + 3z - 7$ restricted to real axis is an odd degree polynomial and so it has either 3 real roots or 1 real root and 2 complex conjugate roots. Now just by observation, at $z = 1$, this function is -3 and at $z=2$, its value is 7 and so by intermediate value theorem it has a real root between 1 and 2. Since the derivative of $z^3 + 3z - 7$ is $3z^2 + 3$ which is always positive for real values of z , it is a strictly increasing function and so can have at most real root β .

Example 5

Lets compute $\int_{|z|=1} \frac{dz}{z(z^3+3z-7)}$. This is an example where we can estimate the location of poles without actually locating them precisely. The function $z^3 + 3z - 7$ restricted to real axis is an odd degree polynomial and so it has either 3 real roots or 1 real root and 2 complex conjugate roots. Now just by observation, at $z = 1$, this function is -3 and at $z=2$, its value is 7 and so by intermediate value theorem it has a real root between 1 and 2. Since the derivative of $z^3 + 3z - 7$ is $3z^2 + 3$ which is always positive for real values of z , it is a strictly increasing function and so can have at most real root β . We have located one real root between 1 and 2, this is the only real root. Hence the other two roots are complex conjugates, say α and $\bar{\alpha}$ and hence have the same magnitude.

Example 5

Lets compute $\int_{|z|=1} \frac{dz}{z(z^3+3z-7)}$. This is an example where we can estimate the location of poles without actually locating them precisely. The function $z^3 + 3z - 7$ restricted to real axis is an odd degree polynomial and so it has either 3 real roots or 1 real root and 2 complex conjugate roots. Now just by observation, at $z = 1$, this function is -3 and at $z=2$, its value is 7 and so by intermediate value theorem it has a real root between 1 and 2. Since the derivative of $z^3 + 3z - 7$ is $3z^2 + 3$ which is always positive for real values of z , it is a strictly increasing function and so can have at most real root β . We have located one real root between 1 and 2, this is the only real root. Hence the other two roots are complex conjugates, say α and $\bar{\alpha}$ and hence have the same magnitude. Since the product of roots is 7, we find that $\beta|\alpha|^2$ is 7. Hence both the complex conjugate roots have magnitude more than one and hence $z^3 + 3z - 7$ is holomorphic inside the unit circle. Hence by CIF

Example 5

Lets compute $\int_{|z|=1} \frac{dz}{z(z^3+3z-7)}$. This is an example where we can estimate the location of poles without actually locating them precisely. The function $z^3 + 3z - 7$ restricted to real axis is an odd degree polynomial and so it has either 3 real roots or 1 real root and 2 complex conjugate roots. Now just by observation, at $z = 1$, this function is -3 and at $z=2$, its value is 7 and so by intermediate value theorem it has a real root between 1 and 2. Since the derivative of $z^3 + 3z - 7$ is $3z^2 + 3$ which is always positive for real values of z , it is a strictly increasing function and so can have at most real root β . We have located one real root between 1 and 2, this is the only real root. Hence the other two roots are complex conjugates, say α and $\bar{\alpha}$ and hence have the same magnitude. Since the product of roots is 7, we find that $\beta|\alpha|^2$ is 7. Hence both the complex conjugate roots have magnitude more than one and hence $z^3 + 3z - 7$ is holomorphic inside the unit circle. Hence by CIF

$$\int_{|z|=1} \frac{1}{z(z^3+3z-7)} = \frac{2\pi i}{z^3+3z-7} \text{ evaluated at } 0 = \frac{-2\pi i}{7}.$$

Many times, one has a situation where Ω is an open set and f is a holomorphic function on the complement of a certain subset.

Singularities

Many times, one has a situation where Ω is an open set and f is a holomorphic function on the complement of a certain subset. The points of this subset are called **singularities** of the function. Given the rigid nature of holomorphic functions, we can get a lot of information on the nature of the singularities; essentially by looking at the function in small punctured neighborhoods of those points. Let us see this in more detail.

Definitions

Singularity of a function: The set of points in Ω where f is not defined or not holomorphic are called the singularities of Ω .
For example $1/z$ has a singularity at 0.

Definitions

Singularity of a function: The set of points in Ω where f is not defined or not holomorphic are called the singularities of Ω .

For example $1/z$ has a singularity at 0.

Singularities are of 2 types, isolated and non-isolated singularities.

A singular point is said to be isolated if the function is holomorphic in a punctured disc around that point.

For example $1/z$ is holomorphic in any punctured disc around 0.

$\frac{1}{z(z-1)}$ has 2 singular points 0 and 1, both of which are isolated singularities; the function is holomorphic in a punctured disc of radius 1 around both of them.

Definitions

Singularity of a function: The set of points in Ω where f is not defined or not holomorphic are called the singularities of Ω .

For example $1/z$ has a singularity at 0.

Singularities are of 2 types, isolated and non-isolated singularities.

A singular point is said to be isolated if the function is holomorphic in a punctured disc around that point.

For example $1/z$ is holomorphic in any punctured disc around 0.

$\frac{1}{z(z-1)}$ has 2 singular points 0 and 1, both of which are isolated singularities; the function is holomorphic in a punctured disc of radius 1 around both of them.

A singularity is non-isolated if it is not isolated ! That is, in no punctured neighborhood of the singularity is the function holomorphic.

For example $f(z) = |z|$ has all points as singularities and hence no point is an isolated singularity.

Removable and Non-Removable Singularities

Isolated singularities are of two types; removable and non-removable singularities.

Removable and Non-Removable Singularities

Isolated singularities are of two types; removable and non-removable singularities.

If an isolated singularity can be removed by defining the function by assigning a certain value at that point, we say that the singularity is removable. If not we say it is non-removable. For instance, the function $f(z) = \frac{\sin(z)}{z}$ is a removable singularity.

Removable and Non-Removable Singularities

Isolated singularities are of two types; removable and non-removable singularities.

If an isolated singularity can be removed by defining the function by assigning a certain value at that point, we say that the singularity is removable. If not we say it is non-removable. For instance, the function $f(z) = \frac{\sin(z)}{z}$ is a removable singularity. By redefining the function to be $f(z) = \frac{\sin(z)}{z}$ for $z \neq 0$ and 1 for $z = 0$, we get a function which is holomorphic even at 0. More generally take any holomorphic function f on Ω and for some point $z_0 \in \Omega$, define $g(z)$ to be $\frac{f(z)-f(z_0)}{z-z_0}$ for $z \neq z_0$. $g(z)$ has a removable singularity at z_0 , which is removed by defining the value at $g(z_0) = f'(z_0)$. The function $1/z$ on \mathbb{C}^* has a non-removable singularity at 0.

Removable and Non-Removable Singularities

Isolated singularities are of two types; removable and non-removable singularities.

If an isolated singularity can be removed by defining the function by assigning a certain value at that point, we say that the singularity is removable. If not we say it is non-removable. For instance, the function $f(z) = \frac{\sin(z)}{z}$ is a removable singularity. By redefining the function to be $f(z) = \frac{\sin(z)}{z}$ for $z \neq 0$ and 1 for $z = 0$, we get a function which is holomorphic even at 0. More generally take any holomorphic function f on Ω and for some point $z_0 \in \Omega$, define $g(z)$ to be $\frac{f(z)-f(z_0)}{z-z_0}$ for $z \neq z_0$. $g(z)$ has a removable singularity at z_0 , which is removed by defining the value at $g(z_0) = f'(z_0)$. The function $1/z$ on \mathbb{C}^* has a non-removable singularity at 0.

Note that if an isolated singularity at z_0 is removable, then $\lim_{z \rightarrow z_0} f(z)$ exists. The converse is also true and that is the Riemann's Removable Singularity Theorem.

Riemann's Removable Singularity Theorem

Theorem: z_0 is removable iff $\lim_{z \rightarrow z_0} f(z)$ exists. Clearly removable singularity implies this limit exists.

Riemann's Removable Singularity Theorem

Theorem: z_0 is removable iff $\lim_{z \rightarrow z_0} f(z)$ exists. Clearly removable singularity implies this limit exists. For the converse, suppose this limit exists. Then $\lim_{z \rightarrow z_0} (z - z_0)f(z) = 0$. Then define

$$g(z) = \begin{cases} (z - z_0)^2 f(z) & \text{if } z \neq z_0 \\ 0 & \text{if } z = z_0. \end{cases}$$

Riemann's Removable Singularity Theorem

Theorem: z_0 is removable iff $\lim_{z \rightarrow z_0} f(z)$ exists. Clearly removable singularity implies this limit exists. For the converse, suppose this limit exists. Then $\lim_{z \rightarrow z_0} (z - z_0)f(z) = 0$. Then define

$$g(z) = \begin{cases} (z - z_0)^2 f(z) & \text{if } z \neq z_0 \\ 0 & \text{if } z = z_0. \end{cases}$$

If f is analytic in a punctured neighbourhood of z_0 , then clearly g is analytic throughout that neighbourhood.

Riemann's Removable Singularity Theorem

Theorem: z_0 is removable iff $\lim_{z \rightarrow z_0} f(z)$ exists. Clearly removable singularity implies this limit exists. For the converse, suppose this limit exists. Then $\lim_{z \rightarrow z_0} (z - z_0)f(z) = 0$. Then define

$$g(z) = \begin{cases} (z - z_0)^2 f(z) & \text{if } z \neq z_0 \\ 0 & \text{if } z = z_0. \end{cases}$$

If f is analytic in a punctured neighbourhood of z_0 , then clearly g is analytic throughout that neighbourhood. Write

$$g(z) = c_0 + c_1(z - z_0) + c_2(z - z_0)^2 + \dots$$

Note that $c_0 = g(z_0) = 0$ and $c_1 = g'(z_0) = 0$.

Riemann's Removable Singularity Theorem

Theorem: z_0 is removable iff $\lim_{z \rightarrow z_0} f(z)$ exists. Clearly removable singularity implies this limit exists. For the converse, suppose this limit exists. Then $\lim_{z \rightarrow z_0} (z - z_0)f(z) = 0$. Then define

$$g(z) = \begin{cases} (z - z_0)^2 f(z) & \text{if } z \neq z_0 \\ 0 & \text{if } z = z_0. \end{cases}$$

If f is analytic in a punctured neighbourhood of z_0 , then clearly g is analytic throughout that neighbourhood. Write

$$g(z) = c_0 + c_1(z - z_0) + c_2(z - z_0)^2 + \dots$$

Note that $c_0 = g(z_0) = 0$ and $c_1 = g'(z_0) = 0$. Thus,

$$g(z) = c_2(z - z_0)^2 + c_3(z - z_0)^3 + \dots$$

If we define $f(z_0) = c_2$, then f is holomorphic throughout. i.e., z_0 is a removable singularity.

Intuitively a pole is a point at which the function blows up from all directions. An isolated singularity z_0 is said to be a pole if $\lim_{z \rightarrow z_0} f(z)$ is ∞ (that is the function takes values outside any bounded set in any small punctured neighborhood of z_0). In this case the function $g(z) = \frac{1}{f(z)}$ is holomorphic at z_0 with $g(0) = 0$. (Why ?).

Intuitively a pole is a point at which the function blows up from all directions. An isolated singularity z_0 is said to be a pole if $\lim_{z \rightarrow z_0} f(z)$ is ∞ (that is the function takes values outside any bounded set in any small punctured neighborhood of z_0). In this case the function $g(z) = \frac{1}{f(z)}$ is holomorphic at z_0 with $g(z_0) = 0$. (Why ?). Since $g(z)$ is not identically equal to zero (Why ?), it follows that there exists a positive integer m such that $g(z) = (z - z_0)^m h(z)$ for some holomorphic function $h(z)$ defined in a neighborhood of z_0 with $h(z_0) \neq 0$. Note that such an m and therefore such a $h(z)$ is uniquely defined.

Intuitively a pole is a point at which the function blows up from all directions. An isolated singularity z_0 is said to be a pole if $\lim_{z \rightarrow z_0} f(z)$ is ∞ (that is the function takes values outside any bounded set in any small punctured neighborhood of z_0). In this case the function $g(z) = \frac{1}{f(z)}$ is holomorphic at z_0 with $g(z_0) = 0$. (Why ?). Since $g(z)$ is not identically equal to zero (Why ?), it follows that there exists a positive integer m such that $g(z) = (z - z_0)^m h(z)$ for some holomorphic function $h(z)$ defined in a neighborhood of z_0 with $h(z_0) \neq 0$. Note that such an m and therefore such a $h(z)$ is uniquely defined. Thus $f(z) = (z - z_0)^{-m} f_1(z)$ for some holomorphic function $f_1(z)$ in a punctured neighborhood of z_0 . m is called the order of the pole and is a measure of how fast the function blows up at z_0 . If m is one, we say that the pole is a **simple pole**.

Casorati-Weierstrass Theorem

A function $f(z)$ defined on an open set except at all the poles is called a **meromorphic function**. An isolated singularity that is neither a pole nor a removable singularity is called an **essentially singularity**. These are the most interesting to understand. Like before we have an important theorem on the values attained by a function near an essential singularity.

Casorati-Weierstrass Theorem

A function $f(z)$ defined on an open set except at all the poles is called a **meromorphic function**. An isolated singularity that is neither a pole nor a removable singularity is called an **essentially singularity**. These are the most interesting to understand. Like before we have an important theorem on the values attained by a function near an essential singularity.

Theorem: If z_0 is an isolated singularity, then it essential if and only if the values of f come arbitrarily close to every complex number in a neighborhood of z_0 .

Casorati-Weierstrass Theorem

A function $f(z)$ defined on an open set except at all the poles is called a **meromorphic function**. An isolated singularity that is neither a pole nor a removable singularity is called an **essentially singularity**. These are the most interesting to understand. Like before we have an important theorem on the values attained by a function near an essential singularity.

Theorem: If z_0 is an isolated singularity, then it is essential if and only if the values of f come arbitrarily close to every complex number in a neighborhood of z_0 .

The if part is obvious. For the only if part, suppose f has an essential singularity. Let a be any complex number. Suppose f does not attain values arbitrarily close to a , then

$$\lim_{z \rightarrow z_0} (z - z_0) \frac{1}{(f(z) - a)} = 0.$$
 Hence by Riemann's theorem above, it has a removable singularity at z_0 .

Depending on whether the singularity can be removed by assigning the value to be zero or a non-zero value, $f(z)$ will have a pole or a removable singularity at z_0 respectively. In either case we have a contradiction.

Depending on whether the singularity can be removed by assigning the value to be zero or a non-zero value, $f(z)$ will have a pole or a removable singularity at z_0 respectively. In either case we have a contradiction. For example, the function $e^{1/z}$ has an essential

singularity at 0. (Check !)

Remark: Its important to note that we only talk about the singularity being removable, pole or essential if it is an isolated singularity. There might be small discrepancies in the convention from one reference to another.

A good mathematical joke is better, and better mathematics, than a dozen mediocre papers. - J E Littlewood.

A good mathematical joke is better, and better mathematics, than a dozen mediocre papers. - J E Littlewood.

(Obviously to be taken in jocular vein !)

So time for a mathematical joke ...

There was a transatlantic flight and the pilot and copilot dropped dead. A desperate flight attendant asked if anyone knew how to fly a plane. An old polish man said: "Well, I used to fly planes in WW II, but nothing like this". When he brought him into the cockpit, his jaw dropped. There were so many buttons, levers, and fancy dials. "What's wrong?" the flight attendant asked.



"I'm just a simple pole in a complex plane", he responded.

Another (non-mathematical) Polish Joke

A Polish immigrant went to the DMV to apply for a driver's license. First, of course, he had to take an eyesight test. The optician showed him a card with the letters:

'C Z W I X N O S T A C Z.'

"Can you read this?" the optician asked.

"Read it?" the Polish guy replied, "I know the guy."