


Signal Processing - 1 by One

Sibi Raj B. Pillai
Dept of Electrical Engineering
IIT Bombay




Cubic Interpolation

Let us interpolate in the interval $[0, T]$ between two samples.




Cubic Interpolation

Let us interpolate in the interval $[0, T]$ between two samples.




Cubic Interpolation

Let us interpolate in the interval $[0, T]$ between two samples.



Cubic Interpolation

Let us interpolate in the interval $[0, T]$ between two samples.




Key: Find the *fitting* polynomial $p(t) = at^3 + bt^2 + ct + d$ in $[0, 1]$.



Cubic Interpolation

Let us interpolate in the interval $[0, T]$ between two samples.



Key: Find the *fitting* polynomial $p(t) = at^3 + bt^2 + ct + d$ in $[0, 1]$.


Eg: For $\delta(t)$ as input, the output in $t \in [0, 1]$

$$p(t) = \frac{3}{2}t^3 - \frac{5}{2}t^2 + 1.$$



Cubic Interpolation

Let us interpolate in the interval $[0, T]$ between two samples.



Key: Find the *fitting* polynomial $p(t) = at^3 + bt^2 + ct + d$ in $[0, 1]$.


Eg: For $\delta(t)$ as input, the output in $t \in [0, 1]$

$$p(t) = \frac{3}{2}t^3 - \frac{5}{2}t^2 + 1.$$

We need to do slightly more work!



Cubic Details




Take $T = 1$ and samples of $f(t)$ at instants $t = 0, \pm 1, \dots$ are given.

$$p(t) = at^3 + bt^2 + ct + d, 0 \leq t \leq 1.$$

$$f(0) = d; f(1) = a + b + c + d;$$



Cubic Details




Take $T = 1$ and samples of $f(t)$ at instants $t = 0, \pm 1, \dots$ are given.

$$p(t) = at^3 + bt^2 + ct + d, 0 \leq t \leq 1.$$

$$f(0) = d; f(1) = a + b + c + d;$$



Cubic Details




Take $T = 1$ and samples of $f(t)$ at instants $t = 0, \pm 1, \dots$ are given.

$$p(t) = at^3 + bt^2 + ct + d, 0 \leq t \leq 1.$$

$$f(0) = d; f(1) = a + b + c + d; f'(0) = c; f'(1) = 3a + 2b + c$$



Cubic Details




Take $T = 1$ and samples of $f(t)$ at instants $t = 0, \pm 1, \dots$ are given.

$$p(t) = at^3 + bt^2 + ct + d, 0 \leq t \leq 1.$$

$$f(0) = d; f(1) = a + b + c + d; f'(0) = c; f'(1) = 3a + 2b + c$$



Cubic Details




Take $T = 1$ and samples of $f(t)$ at instants $t = 0, \pm 1, \dots$ are given.

$$p(t) = at^3 + bt^2 + ct + d, 0 \leq t \leq 1.$$

$$f(0) = d; f(1) = a + b + c + d; f'(0) = c; f'(1) = 3a + 2b + c$$



Cubic Details




Take $T = 1$ and samples of $f(t)$ at instants $t = 0, \pm 1, \dots$ are given.

$$p(t) = at^3 + bt^2 + ct + d, 0 \leq t \leq 1.$$

$$f(0) = d; f(1) = a + b + c + d; f'(0) = c; f'(1) = 3a + 2b + c$$



Cubic Details



Take $T = 1$ and samples of $f(t)$ at instants $t = 0, \pm 1, \dots$ are given.

$$p(t) = at^3 + bt^2 + ct + d, 0 \leq t \leq 1.$$

$$f(0) = d; f(1) = a + b + c + d; f'(0) = c; f'(1) = 3a + 2b + c$$



Cubic Coefficients

$$\begin{bmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 1 & 1 & 1 & 1 \\ 3 & 2 & 1 & 0 \end{bmatrix} \begin{bmatrix} a \\ b \\ c \\ d \end{bmatrix} = \begin{bmatrix} f(0) \\ f'(0) \\ f(1) \\ f'(1) \end{bmatrix}$$



Cubic Coefficients

$$\begin{bmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 1 & 1 & 1 & 1 \\ 3 & 2 & 1 & 0 \end{bmatrix} \begin{bmatrix} a \\ b \\ c \\ d \end{bmatrix} = \begin{bmatrix} f(0) \\ f'(0) \\ f(1) \\ f'(1) \end{bmatrix}$$

$$a = 2f(0) + f'(0) - 2f(1) + f'(1)$$

$$b = -3f(0) - 2f'(0) + 3f(1) - f'(1)$$



Cubic Coefficients

$$\begin{bmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 1 & 1 & 1 & 1 \\ 3 & 2 & 1 & 0 \end{bmatrix} \begin{bmatrix} a \\ b \\ c \\ d \end{bmatrix} = \begin{bmatrix} f(0) \\ f'(0) \\ f(1) \\ f'(1) \end{bmatrix}$$

$$a = 2f(0) + f'(0) - 2f(1) + f'(1)$$

$$b = -3f(0) - 2f'(0) + 3f(1) - f'(1)$$

In interval $[0, T]$ use

$$p\left(\frac{t}{T}\right) = a\left(\frac{t}{T}\right)^3 + b\left(\frac{t}{T}\right)^2 + c\left(\frac{t}{T}\right) + d.$$



Cubic Coefficients

$$\begin{bmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 1 & 1 & 1 & 1 \\ 3 & 2 & 1 & 0 \end{bmatrix} \begin{bmatrix} a \\ b \\ c \\ d \end{bmatrix} = \begin{bmatrix} f(0) \\ f'(0) \\ f(1) \\ f'(1) \end{bmatrix}$$

$$a = 2f(0) + f'(0) - 2f(1) + f'(1)$$

$$b = -3f(0) - 2f'(0) + 3f(1) - f'(1)$$

In interval $[0, T]$ use


$$p\left(\frac{t}{T}\right) = a\left(\frac{t}{T}\right)^3 + b\left(\frac{t}{T}\right)^2 + c\left(\frac{t}{T}\right) + d.$$

Popular Choice:




$$f'(0) = \frac{f(1) - f(-1)}{2}; \quad f'(1) = \frac{f(2) - f(0)}{2};$$






Superposition and Shift



Superposition and Shift



Superposition and Shift




Time Shifting: Shifting the input results in a shifted output.



LTI Systems

Linear System: A system such that for any $x_1(t)$ and $x_2(t)$, if



LTI Systems

Linear System: A system such that for any $x_1(t)$ and $x_2(t)$, if

$$x_1(t) \longrightarrow \boxed{\text{System}} \longrightarrow y_1(t) \quad \text{and}$$

$$x_2(t) \longrightarrow \boxed{\text{System}} \longrightarrow y_2(t) \quad , \text{ then}$$

$$a_1 x_1(t) + a_2 x_2(t) \longrightarrow \boxed{\text{System}} \longrightarrow a_1 y_1(t) + a_2 y_2(t)$$



LTI Systems

Linear System: A system such that for any $x_1(t)$ and $x_2(t)$, if

$$x_1(t) \longrightarrow \boxed{\text{System}} \longrightarrow y_1(t) \quad \text{and}$$

$$x_2(t) \longrightarrow \boxed{\text{System}} \longrightarrow y_2(t) \quad , \text{ then}$$

$$a_1 x_1(t) + a_2 x_2(t) \longrightarrow \boxed{\text{System}} \longrightarrow a_1 y_1(t) + a_2 y_2(t)$$

Time Invariant System



LTI Systems

Linear System: A system such that for any $x_1(t)$ and $x_2(t)$, if

$$x_1(t) \longrightarrow \boxed{\text{System}} \longrightarrow y_1(t) \quad \text{and}$$

$$x_2(t) \longrightarrow \boxed{\text{System}} \longrightarrow y_2(t) \quad , \text{ then}$$

$$a_1 x_1(t) + a_2 x_2(t) \longrightarrow \boxed{\text{System}} \longrightarrow a_1 y_1(t) + a_2 y_2(t)$$

Time Invariant System

$$x_1(t - \tau) \longrightarrow \boxed{\text{System}} \longrightarrow y_1(t - \tau), \quad \forall \tau \in \mathbb{R}$$



LTI Systems

Linear System: A system such that for any $x_1(t)$ and $x_2(t)$, if

$$x_1(t) \longrightarrow \boxed{\text{System}} \longrightarrow y_1(t) \quad \text{and}$$

$$x_2(t) \longrightarrow \boxed{\text{System}} \longrightarrow y_2(t) \quad , \text{ then}$$

$$a_1 x_1(t) + a_2 x_2(t) \longrightarrow \boxed{\text{System}} \longrightarrow a_1 y_1(t) + a_2 y_2(t)$$


Time Invariant System

$$x_1(t - \tau) \longrightarrow \boxed{\text{System}} \longrightarrow y_1(t - \tau), \quad \forall \tau \in \mathbb{R}$$


LTI system: knowing the output to an impulse is good enough.




Cubic Interpolation as LTI System




Cubic Interpolation as LTI System




Cubic Interpolation as LTI System



Cubic Interpolation as LTI System




Cubic Interpolation as LTI System



The signal $h(t)$ is called the **impulse response** of cubic interpoltr.



Cubic Interpolation as LTI System




The signal $h(t)$ is called the **impulse response** of cubic interpoltr.

When $x(t) = \sum_{n \in \mathbb{Z}} x_n \delta(t - nT)$ is input:

$$y(t) = \sum_{n \in \mathbb{Z}} x_n h(t - nT)$$



Cubic Interpolation as LTI System




The signal $h(t)$ is called the **impulse response** of cubic interpoltr.

When $x(t) = \sum_{n \in \mathbb{Z}} x_n \delta(t - nT)$ is input:

$$\begin{aligned} y(t) &= \sum_{n \in \mathbb{Z}} x_n h(t - nT) \\ &\stackrel{?}{=} \left(\sum_{n \in \mathbb{Z}} x_n \delta(t - nT) \right) * h(t) \end{aligned}$$



Cubic Interpolation as LTI System




The signal $h(t)$ is called the **impulse response** of cubic interpoltr.

When $x(t) = \sum_{n \in \mathbb{Z}} x_n \delta(t - nT)$ is input:


$$\begin{aligned} y(t) &= \sum_{n \in \mathbb{Z}} x_n h(t - nT) \\ &\stackrel{?}{=} \left(\sum_{n \in \mathbb{Z}} x_n \delta(t - nT) \right) * h(t) = x(t) * h(t) \end{aligned}$$





For input $\delta(t)$, the output $y(t) > 0$ at $t = -2 + \epsilon \Rightarrow h(t)$ is **non-causal**.





For input $\delta(t)$, the output $y(t) > 0$ at $t = -2 + \epsilon \Rightarrow h(t)$ is **non-causal**.

Practical Implementation:

Use the causal version $h(t - 2T)$ instead \Rightarrow **delay at the output!**

