

MA 205 Complex Analysis: Laurent Series and Examples

September 14, 2020

Recall

Last time we looked at some examples of contour integration. We then began discussing singularities. There are two of singularities: isolated and non-isolated singularities.

Recall

Last time we looked at some examples of contour integration. We then began discussing singularities. There are two of singularities: isolated and non-isolated singularities. Consider $\tan(1/z)$. Is the singularity at 0 isolated or non-isolated ?

Recall

Last time we looked at some examples of contour integration. We then began discussing singularities. There are two of singularities: isolated and non-isolated singularities. Consider $\tan(1/z)$. Is the singularity at 0 isolated or non-isolated ?

Isolated singularities are of 3 types: Removable singularity, Pole and Essential Singularity.

Last time we looked at some examples of contour integration. We then began discussing singularities. There are two of singularities: isolated and non-isolated singularities. Consider $\tan(1/z)$. Is the singularity at 0 isolated or non-isolated ?

Isolated singularities are of 3 types: Removable singularity, Pole and Essential Singularity.

A singularity at z_0 is **removable** if $\lim_{z \rightarrow z_0} f(z)$ exists. In particular $f(z)$ is bounded in a neighborhood of z_0 . Also, if $f(z)$ is bounded in a neighborhood of z_0 , then f has a removable singularity at z_0 . A singularity at z_0 is a **pole** if $f(z) \rightarrow \infty$ as $z \rightarrow z_0$. In particular the function takes unbounded values in any punctured neighborhood of z_0 .

A singularity at z_0 is an **essential singularity** if it is neither a removable singularity nor a pole.

Casorati-Weierstrass Theorem

We also discussed the Casorati Weierstrass theorem which said that restricted to any punctured neighborhood of an essential singularity, the image of $f(z)$ is dense, i.e, comes arbitrarily close to every complex number.

$e^{1/z}$ has an essential singularity at 0. (Why ?)

The Casorati-Weierstrass theorem has a deep generalization, namely the Big Picard Theorem.

Big Picard Theorem: If z_0 is an essential singularity for a function $f(z)$, then in any punctured neighborhood of z_0 , f attains every value except possibly one, infinitely many times.

Big Picard Theorem: If z_0 is an essential singularity for a function $f(z)$, then in any punctured neighborhood of z_0 , f attains every value except possibly one, infinitely many times.

The Big Picard theorem implies another striking and beautiful theorem, namely the **Little Picard Theorem** which states that the image of a non-constant entire function can at most miss one point. Of course I sketched a proof of the fact that if it does miss one point then it is of the form $e^{g(z)}$ for some entire function $g(z)$.

Big Picard Theorem: If z_0 is an essential singularity for a function $f(z)$, then in any punctured neighborhood of z_0 , f attains every value except possibly one, infinitely many times.

The Big Picard theorem implies another striking and beautiful theorem, namely the **Little Picard Theorem** which states that the image of a non-constant entire function can at most miss one point. Of course I sketched a proof of the fact that if it does miss one point then it is of the form $e^{g(z)}$ for some entire function $g(z)$.

I will make some more comments about these theorems at the end of the course, although a complete proof is beyond the scope of this course.

Laurent Series

Recall how we derived the power series representation of a holomorphic function on a disc centered around z_0 . We used

$$f(z) = \frac{1}{2\pi i} \int_{|w-z_0|=r} \frac{f(w)}{w-z} dw,$$

and manipulated $\frac{1}{w-z}$ as

$$\frac{1}{w-z_0} \cdot \frac{1}{1 - \frac{z-z_0}{w-z_0}}.$$


Now suppose z_0 is an isolated singularity for f . Consider an annulus with radii $R > r$ centered at z_0 such that f is holomorphic there. CIF takes the form:

Now suppose z_0 is an isolated singularity for f . Consider an annulus with radii $R > r$ centered at z_0 such that f is holomorphic there. CIF takes the form:

$$f(z) = \frac{1}{2\pi i} \int_{|w-z_0|=R} \frac{f(w)}{w-z} dw - \frac{1}{2\pi i} \int_{|w-z_0|=r} \frac{f(w)}{w-z} dw.$$

Now suppose z_0 is an isolated singularity for f . Consider an annulus with radii $R > r$ centered at z_0 such that f is holomorphic there. CIF takes the form:

$$f(z) = \frac{1}{2\pi i} \int_{|w-z_0|=R} \frac{f(w)}{w-z} dw - \frac{1}{2\pi i} \int_{|w-z_0|=r} \frac{f(w)}{w-z} dw.$$



Laurent Series

The first integral gives rise to $\sum_{n=1}^{\infty} a_n (z - z_0)^n$ with

$$a_n = \frac{1}{2\pi i} \int_{|z-z_0|=R} \frac{f(z)}{(z-z_0)^{n+1}} dz,$$

exactly as before.

In the second integral, write

$$\frac{-1}{w-z} = \frac{1}{z-z_0} \cdot \frac{1}{1 - \frac{w-z_0}{z-z_0}},$$

Note that $\left| \frac{w-z_0}{z-z_0} \right| < 1$ for all w with $|w - z_0| = r$.

Expand to get $\sum_{n=1}^{\infty} \frac{b_n}{(z - z_0)^n}$ with :

Laurent Series

The first integral gives rise to $\sum_{n=1}^{\infty} a_n (z - z_0)^n$ with

$$a_n = \frac{1}{2\pi i} \int_{|z-z_0|=R} \frac{f(z)}{(z-z_0)^{n+1}} dz,$$

exactly as before.

In the second integral, write

$$\frac{-1}{w-z} = \frac{1}{z-z_0} \cdot \frac{1}{1 - \frac{w-z_0}{z-z_0}},$$

Note that $|\frac{w-z_0}{z-z_0}| < 1$ for all w with $|w-z_0| = r$.

Expand to get $\sum_{n=1}^{\infty} \frac{b_n}{(z-z_0)^n}$ with :

$$b_n = \frac{1}{2\pi i} \int_{|z-z_0|=r} \frac{f(z)}{(z-z_0)^{-n+1}} dz.$$

Laurent Series

The first integral gives rise to $\sum_{n=1}^{\infty} a_n (z - z_0)^n$ with

$$a_n = \frac{1}{2\pi i} \int_{|z-z_0|=R} \frac{f(z)}{(z-z_0)^{n+1}} dz,$$

exactly as before.

In the second integral, write

$$\frac{-1}{w-z} = \frac{1}{z-z_0} \cdot \frac{1}{1 - \frac{w-z_0}{z-z_0}},$$

Note that $|\frac{w-z_0}{z-z_0}| < 1$ for all w with $|w-z_0| = r$.

Expand to get $\sum_{n=1}^{\infty} \frac{b_n}{(z-z_0)^n}$ with :

$$b_n = \frac{1}{2\pi i} \int_{|z-z_0|=r} \frac{f(z)}{(z-z_0)^{-n+1}} dz.$$

We write both together as $\sum_{n=-\infty}^{\infty} a_n(z - z_0)^n$. This is the Laurent series around the isolated singularity z_0 . The negative part is called the principal part of the Laurent series. If z_0 is an isolated singularity of f , then f is holomorphic in an annulus $0 < |z - z_0| < R$ for some R . The corresponding Laurent expansion is called the Laurent expansion around z_0 .

Residue

We write both together as $\sum_{n=-\infty}^{\infty} a_n(z - z_0)^n$. This is the Laurent series around the isolated singularity z_0 . The negative part is called the principal part of the Laurent series. If z_0 is an isolated singularity of f , then f is holomorphic in an annulus $0 < |z - z_0| < R$ for some R . The corresponding Laurent expansion is called the Laurent expansion around z_0 . Consider the 1st coefficient of this Laurent series.

$$a_{-1} = \frac{1}{2\pi i} \int_{\gamma} f(z) dz.$$

If you integrate a Laurent series, only a_{-1} remains; other terms vanish. What remains is usually called a residue.

$$a_{-1} = \text{Res}(f; z_0).$$

Often a_{-1} is easy to compute from $f(z)$ and if that's the case integration has become easy.

Cauchy Residue Theorem

Suppose f is given and γ is given.

Cauchy Residue Theorem

Suppose f is given and γ is given. Suppose there are finitely many isolated singularities of f inside γ ; say z_1, z_2, \dots, z_n .

Cauchy Residue Theorem

Suppose f is given and γ is given. Suppose there are finitely many isolated singularities of f inside γ ; say z_1, z_2, \dots, z_n . What's $\int_{\gamma} f(z) dz$?


Cauchy Residue Theorem

Suppose f is given and γ is given. Suppose there are finitely many isolated singularities of f inside γ ; say z_1, z_2, \dots, z_n . What's $\int_{\gamma} f(z) dz$?

Theorem (Cauchy Residue Theorem)

$$\int_{\gamma} f(z) dz = 2\pi i \cdot \sum_{i=1}^n \operatorname{Res}(f, z_i).$$

Proof : We have already seen the proof in the previous lectures. The following figure should remind you of the proof. (Here the case of 2 singularities is considered; similarly one can handle more singular points)




Cauchy Residue Theorem

Suppose f is given and γ is given. Suppose there are finitely many isolated singularities of f inside γ ; say z_1, z_2, \dots, z_n . What's $\int_{\gamma} f(z) dz$?

Theorem (Cauchy Residue Theorem)

$$\int_{\gamma} f(z) dz = 2\pi i \cdot \sum_{i=1}^n \operatorname{Res}(f, z_i).$$

Proof : We have already seen the proof in the previous lectures. The following figure should remind you of the proof. (Here the case of 2 singularities is considered; similarly one can handle more singular points)



Why did the mathematician name his dog Cauchy?

Why did the mathematician name his dog Cauchy?




Why did the mathematician name his dog Cauchy?




B'coz it left a residue at every pole!

What's the contour integral over Western Europe?

What's the contour integral over Western Europe?



What's the contour integral over Western Europe?



Zero. All the poles are in Eastern Europe.

Modification: Actually there are poles in Western Europe but they are all removable !!

Principal Part of the Laurent Series

If $\sum_{n=-\infty}^{\infty} a_n(z - z_0)^n$ is the Laurent expansion around z_0 , then its principal part is

$$\sum_{n=-\infty}^{-1} a_n(z - z_0)^n.$$

Principal Part of the Laurent Series

If $\sum_{n=-\infty}^{\infty} a_n(z - z_0)^n$ is the Laurent expansion around z_0 , then its principal part is

$$\sum_{n=-\infty}^{-1} a_n(z - z_0)^n.$$

Note that:

- removable iff principal part is zero.
- pole iff principal part is finite.
- essential iff principal part is infinite.

Proof (Easy exercise using previous slides).

Residue at a Pole

If the isolated singularity is removable, then the residue is trivial. If the isolated singularity is a pole, then the residue is trivial to compute. If z_0 is a pole, can write

$$f(z) = \frac{a_{-m}}{(z - z_0)^m} + \dots + \frac{a_{-1}}{(z - z_0)} + a_0 + a_1(z - z_0) + a_2(z - z_0)^2 + \dots$$

Residue at a Pole

If the isolated singularity is removable, then the residue is trivial. If the isolated singularity is a pole, then the residue is trivial to compute. If z_0 is a pole, can write

$$f(z) = \frac{a_{-m}}{(z-z_0)^m} + \dots + \frac{a_{-1}}{(z-z_0)} + a_0 + a_1(z-z_0) + a_2(z-z_0)^2 + \dots$$

Thus,

$$g(z) = (z-z_0)^m f(z) = a_{-m} + \dots + a_{-1}(z-z_0)^{m-1} + a_0(z-z_0)^m + \dots$$

Residue at a Pole

If the isolated singularity is removable, then the residue is trivial. If the isolated singularity is a pole, then the residue is trivial to compute. If z_0 is a pole, can write

$$f(z) = \frac{a_{-m}}{(z - z_0)^m} + \dots + \frac{a_{-1}}{(z - z_0)} + a_0 + a_1(z - z_0) + a_2(z - z_0)^2 + \dots$$

Thus,

$$g(z) = (z - z_0)^m f(z) = a_{-m} + \dots + a_{-1}(z - z_0)^{m-1} + a_0(z - z_0)^m + \dots$$

Thus, g is holomorphic and

$$a_{-1} = \frac{g^{(m-1)}(z_0)}{(m-1)!}.$$

Example

Find the Laurent Series expansion of $\frac{1}{(z-1)(z-2)}$ in the annulus $1 < |z| < 2$.

Example

Find the Laurent Series expansion of $\frac{1}{(z-1)(z-2)}$ in the annulus $1 < |z| < 2$.

$$\frac{1}{(z-1)(z-2)} = \frac{1}{z-2} - \frac{1}{z-1}.$$

For the first term, $\frac{1}{z-2} = -\frac{1}{2}\left(\frac{1}{1-\frac{z}{2}}\right) = -\frac{1}{2} \sum_{n=0}^{\infty} \left(\frac{z}{2}\right)^n.$

For the second term, $\frac{1}{z-1} = \frac{1}{z}\left(\frac{1}{1-1/z}\right) = \sum_{n=1}^{\infty} \frac{1}{z^n}.$

Example

Find the Laurent Series expansion of $\frac{1}{(z-1)(z-2)}$ in the annulus $1 < |z| < 2$.

$$\frac{1}{(z-1)(z-2)} = \frac{1}{z-2} - \frac{1}{z-1}.$$

For the first term, $\frac{1}{z-2} = -\frac{1}{2}\left(\frac{1}{1-\frac{z}{2}}\right) = -\frac{1}{2} \sum_{n=0}^{\infty} \left(\frac{z}{2}\right)^n.$

For the second term, $\frac{1}{z-1} = \frac{1}{z}\left(\frac{1}{1-1/z}\right) = \sum_{n=1}^{\infty} \frac{1}{z^n}.$

Putting the two together we get the desired Laurent Series Expansion.

Example

Determine the Laurent series of $e^{1/z}$ around the point 0.

$$\begin{aligned} e^{1/z} &= \sum_0^{\infty} \frac{1}{n!} \frac{1}{z^n} \\ &= \sum_{-\infty}^0 \frac{z^n}{(-n)!} \end{aligned}$$

Example

Determine the Laurent series of $\sin\left(\frac{1}{z-1}\right)$ in the range $1 < |z| < 2$.

$\sin\left(\frac{1}{z-1}\right)$

Example

Determine the Laurent series of $\sin\left(\frac{1}{z-1}\right)$ in the range $1 < |z| < 2$.

$$\begin{aligned} & \sin\left(\frac{1}{z-1}\right) \\ &= \sin\left(\frac{1}{z(1-\frac{1}{z})}\right) \end{aligned}$$

Example

Determine the Laurent series of $\sin(\frac{1}{z-1})$ in the range $1 < |z| < 2$.

$$\sin(\frac{1}{z-1})$$

$$= \sin(\frac{1}{z(1-\frac{1}{z})})$$

$$= \sin(\frac{1}{z} + \frac{1}{z^2} + \frac{1}{z^3} \cdots)$$

Example

Determine the Laurent series of $\sin(\frac{1}{z-1})$ in the range $1 < |z| < 2$.

$$\sin(\frac{1}{z-1})$$

$$= \sin(\frac{1}{z(1-\frac{1}{z})})$$

$$= \sin(\frac{1}{z} + \frac{1}{z^2} + \frac{1}{z^3} \cdots)$$

$$= (\frac{1}{z} + \frac{1}{z^2} + \frac{1}{z^3} \cdots) - \frac{1}{3!}(\frac{1}{z} + \frac{1}{z^2} + \frac{1}{z^3} \cdots)^3 + \frac{1}{5!}(\frac{1}{z} + \frac{1}{z^2} + \frac{1}{z^3} \cdots)^5 + \cdots$$

Example

Determine the Laurent series of $\sin(\frac{1}{z-1})$ in the range $1 < |z| < 2$.

$$\begin{aligned}\sin\left(\frac{1}{z-1}\right) &= \sin\left(\frac{1}{z(1-\frac{1}{z})}\right) \\ &= \sin\left(\frac{1}{z} + \frac{1}{z^2} + \frac{1}{z^3} \cdots\right) \\ &= \left(\frac{1}{z} + \frac{1}{z^2} + \frac{1}{z^3} \cdots\right) - \frac{1}{3!}\left(\frac{1}{z} + \frac{1}{z^2} + \frac{1}{z^3} \cdots\right)^3 + \frac{1}{5!}\left(\frac{1}{z} + \frac{1}{z^2} + \frac{1}{z^3} \cdots\right)^5 + \cdots \\ &= \frac{1}{z} + \frac{1}{z^2} + \frac{5}{6}\frac{1}{z^3} + \frac{1}{2}\frac{1}{z^4} + \cdots\end{aligned}$$

Example

Determine the Laurent series of $f(z) = \frac{1}{z+2}$ around $z = 1$.

The given function has a pole at -2. So we will break up the Laurent series computation for the regions $|z - 1| < 3$ and the region $|z - 1| > 3$. In the region $|z - 1| < 3$, the Laurent series will coincide with the Taylor series since the function is holomorphic there.

For the region $|z - 1| < 3$, write $f(z) = \frac{1}{3+(z-1)} = \frac{1}{3} \frac{1}{1+\frac{z-1}{3}}$.

Since $|z - 1| < 3$, we can expand this into a geometric series and get $f(z) = \sum_0^\infty \frac{(-1)^n (z-1)^n}{3^{n+1}}$ in this region. For the region

$$\begin{aligned} |z - 1| > 3, \text{ write } f(z) &= \frac{1}{3+(z-1)} = \frac{1}{z-1} \frac{1}{1+\frac{3}{z-1}} = \frac{1}{z-1} \sum_0^\infty \frac{(-1)^n 3^n}{(z-1)^n} \\ &= \sum_0^\infty \frac{(-1)^n 3^n}{(z-1)^{n+1}} \end{aligned}$$