

MA 207 - Differential Equations-II

Amiya Kumar Pani


Department of Mathematics,
Indian Institute of Technology Bombay,
Powai, Mumbai 76
akp.ma207.2020@gmail.com

November 7, 2020

Start with

"The sciences do not try to explain, they hardly even try to interpret, they mainly make models. By a model is meant a mathematical construct which, with the addition of certain verbal interpretations, describes observed phenomena. The justification of such a mathematical construct is solely and precisely that it is expected to work—that is, correctly to describe phenomena from a reasonably wide area. Furthermore, it must satisfy certain esthetic criteria—that is, in relation to how much it describes, it must be rather simple."

by John von Neumann



Start with

"The sciences do not try to explain, they hardly even try to interpret, they mainly make models. By a model is meant a mathematical construct which, with the addition of certain verbal interpretations, describes observed phenomena. The justification of such a mathematical construct is solely and precisely that it is expected to work—that is, correctly to describe phenomena from a reasonably wide area. Furthermore, it must satisfy certain esthetic criteria—that is, in relation to how much it describes, it must be rather simple."

by John von Neumann



The ultimate goal of a meteorologist is to set up differential equations of the movements of the air and to obtain, as their integral, the general atmospheric circulation, and as particular integrals the cyclones, anticyclones, tornados, and thunderstorms.

Amirja Mourvic - QUOTESTATS.COM



Outline of the lecture

- Elliptic PDE: Laplace Equation, Poisson Equation
- Uniqueness of Solution
- Heat Conduction Equation - Separation of variables
- Non-homogeneous boundary data
- Non-homogeneous equation

Poisson equation in a bounded domain

- Let Ω be an open, bounded subset of \mathbb{R}^n with boundary $\partial\Omega$.

Poisson equation in a bounded domain

- Let Ω be an open, bounded subset of \mathbb{R}^n with boundary $\partial\Omega$.
- Consider
$$-\Delta u \equiv - \sum_{i=1}^n \frac{\partial^2 u}{\partial x_i^2} = f(x), \quad u = u(x_1, x_2, \dots, x_n), \quad x \in \Omega, \text{ with}$$
any of the following prescribed boundary conditions:

Poisson equation in a bounded domain

- Let Ω be an open, bounded subset of \mathbb{R}^n with boundary $\partial\Omega$.
- Consider

$$-\Delta u \equiv - \sum_{i=1}^n \frac{\partial^2 u}{\partial x_i^2} = f(x), \quad u = u(x_1, x_2, \dots, x_n), \quad x \in \Omega, \text{ with}$$

any of the following prescribed boundary conditions:

- **Dirichlet BC** : $u = g$ on $\partial\Omega$. ($g = 0 \implies$ homogeneous BC)

Poisson equation in a bounded domain

- Let Ω be an open, bounded subset of \mathbb{R}^n with boundary $\partial\Omega$.
- Consider

$$-\Delta u \equiv - \sum_{i=1}^n \frac{\partial^2 u}{\partial x_i^2} = f(x), \quad u = u(x_1, x_2, \dots, x_n), \quad x \in \Omega, \text{ with}$$

any of the following prescribed boundary conditions:

- **Dirichlet BC** : $u = g$ on $\partial\Omega$. ($g = 0 \implies$ homogeneous BC)
- **Neumann BC** : $\frac{\partial u}{\partial \nu} \equiv \nabla u \cdot \nu = g$ on $\partial\Omega$, where ν is the outward normal to Ω .

Poisson equation in a bounded domain

- Let Ω be an open, bounded subset of \mathbb{R}^n with boundary $\partial\Omega$.
- Consider

$$-\Delta u \equiv - \sum_{i=1}^n \frac{\partial^2 u}{\partial x_i^2} = f(x), \quad u = u(x_1, x_2, \dots, x_n), \quad x \in \Omega, \text{ with}$$

any of the following prescribed boundary conditions:

- **Dirichlet BC** : $u = g$ on $\partial\Omega$. ($g = 0 \implies$ homogeneous BC)
- **Neumann BC** : $\frac{\partial u}{\partial \nu} \equiv \nabla u \cdot \nu = g$ on $\partial\Omega$, where ν is the outward normal to Ω .
- **Robin BC** : $\frac{\partial u}{\partial \nu} + \alpha u = g$ on $\partial\Omega$, $\alpha > 0$ being a real constant.

Poisson equation in a bounded domain

- Let Ω be an open, bounded subset of \mathbb{R}^n with boundary $\partial\Omega$.
- Consider

$$-\Delta u \equiv - \sum_{i=1}^n \frac{\partial^2 u}{\partial x_i^2} = f(x), \quad u = u(x_1, x_2, \dots, x_n), \quad x \in \Omega, \text{ with}$$

any of the following prescribed boundary conditions:

- **Dirichlet BC** : $u = g$ on $\partial\Omega$. ($g = 0 \implies$ homogeneous BC)
- **Neumann BC** : $\frac{\partial u}{\partial \nu} \equiv \nabla u \cdot \nu = g$ on $\partial\Omega$, where ν is the outward normal to Ω .
- **Robin BC** : $\frac{\partial u}{\partial \nu} + \alpha u = g$ on $\partial\Omega$, $\alpha > 0$ being a real constant.
- **Mixed BC** : Let $\partial\Omega = \partial\bar{\Omega}_D \cup \partial\bar{\Omega}_N$, $\partial\Omega_D \cap \partial\Omega_N = \emptyset$.

Poisson equation in a bounded domain

- Let Ω be an open, bounded subset of \mathbb{R}^n with boundary $\partial\Omega$.
- Consider

$$-\Delta u \equiv - \sum_{i=1}^n \frac{\partial^2 u}{\partial x_i^2} = f(x), \quad u = u(x_1, x_2, \dots, x_n), \quad x \in \Omega, \text{ with}$$

any of the following prescribed boundary conditions:

- **Dirichlet BC** : $u = g$ on $\partial\Omega$. ($g = 0 \implies$ homogeneous BC)
- **Neumann BC** : $\frac{\partial u}{\partial \nu} \equiv \nabla u \cdot \nu = g$ on $\partial\Omega$, where ν is the outward normal to Ω .
- **Robin BC** : $\frac{\partial u}{\partial \nu} + \alpha u = g$ on $\partial\Omega$, $\alpha > 0$ being a real constant.
- **Mixed BC** : Let $\partial\Omega = \partial\bar{\Omega}_D \cup \partial\bar{\Omega}_N$, $\partial\Omega_D \cap \partial\Omega_N = \emptyset$.

$$u = g_1 \text{ on } \partial\Omega_D, \quad \frac{\partial u}{\partial \nu} = g_2 \text{ on } \partial\Omega_N.$$

Pierre-Simon, marquis de Laplace (23 March 1749 – 5 March 1827)

“ French truly polymath who made important to contributions to engineering, mathematics, statistics, physics, astronomy, and philosophy.

Called French Newton

His five-volumes on Celestial Mechanics (1799–1825) were classic .

Formulated Laplace’s equation, and pioneered the Laplace transform

The Laplacian differential operator, widely used in mathematics, is also named after him.

He restated and developed the nebular hypothesis of the origin of the Solar System and

He was one of the first scientists to postulate the existence of black holes and the notion of gravitational collapse.



Baron Simeon Denis Poisson (21 June 1781 – 25 April 1840)

*“ French mathematician, engineer, and physicist
who made many scientific advances
Known for his work on definite integrals,
electromagnetic theory, and probability.
In probability: Poisson Distribution is named after
him.
Worked on Celestial Mechanics (extended works of
Laplace and Lagrange)
300 papers published*



Theorem

Let Ω be bounded domain whose boundary $\partial\Omega$ is piece-wise smooth and orientable.

Theorem

Let Ω be bounded domain whose boundary $\partial\Omega$ is piece-wise smooth and orientable. Let \vec{F} be a vector function which is continuously differentiable.

Digression-Gauss Divergence Theorem

Theorem

Let Ω be bounded domain whose boundary $\partial\Omega$ is piece-wise smooth and orientable. Let \vec{F} be a vector function which is continuously differentiable. Then,

$$\underbrace{\int_{\Omega} \nabla \cdot \vec{F} \, dx}_{\sum_{i=1}^n \int_{\Omega} \frac{\partial F_i}{\partial x_i} \, dx} = \underbrace{\int_{\partial\Omega} \vec{F} \cdot \nu \, ds}_{\sum_{i=1}^n \int_{\partial\Omega} F_i \nu_i \, ds} .$$

As a consequence of the divergence theorem, we have

As a consequence of the divergence theorem, we have

$$\int_{\Omega} \Delta uv \, dx = \int_{\partial\Omega} v \frac{\partial u}{\partial \nu} \, ds - \int_{\Omega} \nabla u \cdot \nabla v \, dx$$

As a consequence of the divergence theorem, we have

$$\int_{\Omega} \Delta uv \, dx = \int_{\partial\Omega} v \frac{\partial u}{\partial \nu} \, ds - \int_{\Omega} \nabla u \cdot \nabla v \, dx$$

Proof:

$$\int_{\Omega} \Delta uv \, dx = \int_{\Omega} \nabla \cdot (\nabla u) v \, dx$$

As a consequence of the divergence theorem, we have

$$\int_{\Omega} \Delta uv \, dx = \int_{\partial\Omega} v \frac{\partial u}{\partial \nu} \, ds - \int_{\Omega} \nabla u \cdot \nabla v \, dx$$

Proof:

$$\begin{aligned} \int_{\Omega} \Delta uv \, dx &= \int_{\Omega} \nabla \cdot (\nabla u) v \, dx \\ &= \int_{\Omega} \left(\nabla \cdot (v \nabla u) - \nabla u \cdot \nabla v \right) dx \end{aligned}$$

As a consequence of the divergence theorem, we have

$$\int_{\Omega} \Delta uv \, dx = \int_{\partial\Omega} v \frac{\partial u}{\partial \nu} \, ds - \int_{\Omega} \nabla u \cdot \nabla v \, dx$$

Proof:

$$\begin{aligned} \int_{\Omega} \Delta uv \, dx &= \int_{\Omega} \nabla \cdot (\nabla u) v \, dx \\ &= \int_{\Omega} \left(\nabla \cdot (v \nabla u) - \nabla u \cdot \nabla v \right) dx \end{aligned}$$

Applying divergence theorem to the first term on the right hand side, we obtain

As a consequence of the divergence theorem, we have

$$\int_{\Omega} \Delta uv \, dx = \int_{\partial\Omega} v \frac{\partial u}{\partial \nu} \, ds - \int_{\Omega} \nabla u \cdot \nabla v \, dx$$

Proof:

$$\begin{aligned} \int_{\Omega} \Delta uv \, dx &= \int_{\Omega} \nabla \cdot (\nabla u) v \, dx \\ &= \int_{\Omega} \left(\nabla \cdot (v \nabla u) - \nabla u \cdot \nabla v \right) dx \end{aligned}$$

Applying divergence theorem to the first term on the right hand side, we obtain

$$\int_{\Omega} \Delta uv \, dx = \int_{\partial\Omega} v \frac{\partial u}{\partial \nu} \, ds - \int_{\Omega} \nabla u \cdot \nabla v \, dx.$$

As a consequence of the divergence theorem, we have

$$\int_{\Omega} \Delta uv \, dx = \int_{\partial\Omega} v \frac{\partial u}{\partial \nu} \, ds - \int_{\Omega} \nabla u \cdot \nabla v \, dx$$

Proof:

$$\begin{aligned} \int_{\Omega} \Delta uv \, dx &= \int_{\Omega} \nabla \cdot (\nabla u) v \, dx \\ &= \int_{\Omega} \left(\nabla \cdot (v \nabla u) - \nabla u \cdot \nabla v \right) dx \end{aligned}$$

Applying divergence theorem to the first term on the right hand side, we obtain

$$\int_{\Omega} \Delta uv \, dx = \int_{\partial\Omega} v \frac{\partial u}{\partial \nu} \, ds - \int_{\Omega} \nabla u \cdot \nabla v \, dx.$$

Elliptic problem (Uniqueness of solution)

Show that the problem

$$-\Delta u = f \text{ in } \Omega$$

Elliptic problem (Uniqueness of solution)

Show that the problem

$$-\Delta u = f \text{ in } \Omega$$

$$u = g \text{ on } \partial\Omega$$

Elliptic problem (Uniqueness of solution)

Show that the problem

$$-\Delta u = f \text{ in } \Omega$$

$$u = g \text{ on } \partial\Omega$$

has a unique solution.

If possible, let u_1 and u_2 be two distinct solutions of the Dirichlet problem, that is u_i , ($i = 1, 2$) satisfies

$$-\Delta u_i = f \text{ in } \Omega$$

Elliptic problem (Uniqueness of solution)

Show that the problem

$$-\Delta u = f \text{ in } \Omega$$

$$u = g \text{ on } \partial\Omega$$

has a unique solution.

If possible, let u_1 and u_2 be two distinct solutions of the Dirichlet problem, that is u_i , ($i = 1, 2$) satisfies

$$-\Delta u_i = f \text{ in } \Omega$$

$$u_i = g \text{ on } \partial\Omega$$

Elliptic problem (Uniqueness of solution)

Show that the problem

$$-\Delta u = f \text{ in } \Omega$$

$$u = g \text{ on } \partial\Omega$$

has a unique solution.

If possible, let u_1 and u_2 be two distinct solutions of the Dirichlet problem, that is u_i , ($i = 1, 2$) satisfies

$$-\Delta u_i = f \text{ in } \Omega$$

$$u_i = g \text{ on } \partial\Omega$$

Let $w = u_1 - u_2$.

Elliptic problem (Uniqueness of solution)

Show that the problem

$$-\Delta u = f \text{ in } \Omega$$

$$u = g \text{ on } \partial\Omega$$

has a unique solution.

If possible, let u_1 and u_2 be two distinct solutions of the Dirichlet problem, that is u_i , ($i = 1, 2$) satisfies

$$-\Delta u_i = f \text{ in } \Omega$$

$$u_i = g \text{ on } \partial\Omega$$

Let $w = u_1 - u_2$. Then, w satisfies

$$-\Delta w = 0 \text{ in } \Omega$$

Elliptic problem (Uniqueness of solution)

Show that the problem

$$-\Delta u = f \text{ in } \Omega$$

$$u = g \text{ on } \partial\Omega$$

has a unique solution.

If possible, let u_1 and u_2 be two distinct solutions of the Dirichlet problem, that is u_i , ($i = 1, 2$) satisfies

$$-\Delta u_i = f \text{ in } \Omega$$

$$u_i = g \text{ on } \partial\Omega$$

Let $w = u_1 - u_2$. Then, w satisfies

$$-\Delta w = 0 \text{ in } \Omega$$

$$w = 0 \text{ on } \partial\Omega.$$

Elliptic problem (Uniqueness of solution)

Show that the problem

$$-\Delta u = f \text{ in } \Omega$$

$$u = g \text{ on } \partial\Omega$$

has a unique solution.

If possible, let u_1 and u_2 be two distinct solutions of the Dirichlet problem, that is u_i , ($i = 1, 2$) satisfies

$$-\Delta u_i = f \text{ in } \Omega$$

$$u_i = g \text{ on } \partial\Omega$$

Let $w = u_1 - u_2$. Then, w satisfies

$$-\Delta w = 0 \text{ in } \Omega$$

$$w = 0 \text{ on } \partial\Omega.$$

Claim : $w \equiv 0$.

Elliptic problem (Uniqueness of solution)

Show that the problem

$$-\Delta u = f \text{ in } \Omega$$

$$u = g \text{ on } \partial\Omega$$

has a unique solution.

If possible, let u_1 and u_2 be two distinct solutions of the Dirichlet problem, that is u_i , ($i = 1, 2$) satisfies

$$-\Delta u_i = f \text{ in } \Omega$$

$$u_i = g \text{ on } \partial\Omega$$

Let $w = u_1 - u_2$. Then, w satisfies

$$-\Delta w = 0 \text{ in } \Omega$$

$$w = 0 \text{ on } \partial\Omega.$$

Claim : $w \equiv 0$.

Multiply the first equation above by w and integrate over Ω to obtain

Multiply the first equation above by w and integrate over Ω to obtain

$$0 = - \int_{\Omega} w \Delta w \, d\Omega$$

Multiply the first equation above by w and integrate over Ω to obtain

$$\begin{aligned} 0 &= - \int_{\Omega} w \Delta w \, d\Omega \\ &= - \int_{\partial\Omega} w \frac{\partial w}{\partial \nu} \, ds + \int_{\Omega} |\nabla w|^2 \, dx \end{aligned}$$

Multiply the first equation above by w and integrate over Ω to obtain

$$\begin{aligned} 0 &= - \int_{\Omega} w \Delta w \, d\Omega \\ &= - \int_{\partial\Omega} w \frac{\partial w}{\partial \nu} \, ds + \int_{\Omega} |\nabla w|^2 \, dx \end{aligned}$$

Now $w = 0$ on $\partial\Omega$.

Multiply the first equation above by w and integrate over Ω to obtain

$$\begin{aligned} 0 &= - \int_{\Omega} w \Delta w \, d\Omega \\ &= - \int_{\partial\Omega} w \frac{\partial w}{\partial \nu} \, ds + \int_{\Omega} |\nabla w|^2 \, dx \end{aligned}$$

Now $w = 0$ on $\partial\Omega$.

Hence $\int_{\Omega} |\nabla w|^2 \, dx = 0 \implies w$ is a constant in Ω .

Multiply the first equation above by w and integrate over Ω to obtain

$$\begin{aligned} 0 &= - \int_{\Omega} w \Delta w \, d\Omega \\ &= - \int_{\partial\Omega} w \frac{\partial w}{\partial \nu} \, ds + \int_{\Omega} |\nabla w|^2 \, dx \end{aligned}$$

Now $w = 0$ on $\partial\Omega$.

Hence $\int_{\Omega} |\nabla w|^2 \, dx = 0 \implies w$ is a constant in Ω .

$w = 0$ on $\partial\Omega$; w being a continuous function, we deduce $w \equiv 0$ in Ω .

Multiply the first equation above by w and integrate over Ω to obtain

$$\begin{aligned} 0 &= - \int_{\Omega} w \Delta w \, d\Omega \\ &= - \int_{\partial\Omega} w \frac{\partial w}{\partial \nu} \, ds + \int_{\Omega} |\nabla w|^2 \, dx \end{aligned}$$

Now $w = 0$ on $\partial\Omega$.

Hence $\int_{\Omega} |\nabla w|^2 \, dx = 0 \implies w$ is a constant in Ω .

$w = 0$ on $\partial\Omega$; w being a continuous function, we deduce $w \equiv 0$ in Ω .

$\implies u_1 \equiv u_2$ in Ω .

Multiply the first equation above by w and integrate over Ω to obtain

$$\begin{aligned} 0 &= - \int_{\Omega} w \Delta w \, d\Omega \\ &= - \int_{\partial\Omega} w \frac{\partial w}{\partial \nu} \, ds + \int_{\Omega} |\nabla w|^2 \, dx \end{aligned}$$

Now $w = 0$ on $\partial\Omega$.

Hence $\int_{\Omega} |\nabla w|^2 \, dx = 0 \implies w$ is a constant in Ω .

$w = 0$ on $\partial\Omega$; w being a continuous function, we deduce $w \equiv 0$ in Ω .

$\implies u_1 \equiv u_2$ in Ω .

\implies The Dirichlet problem has a unique solution.

Multiply the first equation above by w and integrate over Ω to obtain

$$\begin{aligned} 0 &= - \int_{\Omega} w \Delta w \, d\Omega \\ &= - \int_{\partial\Omega} w \frac{\partial w}{\partial \nu} \, ds + \int_{\Omega} |\nabla w|^2 \, dx \end{aligned}$$

Now $w = 0$ on $\partial\Omega$.

Hence $\int_{\Omega} |\nabla w|^2 \, dx = 0 \implies w$ is a constant in Ω .

$w = 0$ on $\partial\Omega$; w being a continuous function, we deduce $w \equiv 0$ in Ω .

$\implies u_1 \equiv u_2$ in Ω .

\implies The Dirichlet problem has a unique solution.

Exercise : 1. Verify for other BC's.

2. Tut. Sheet 4 - Qns. 6, 7(i), 8

Parabolic initial and boundary value problem (Uniqueness of solution)

Show that the problem

$$u_t - \Delta u = f \text{ in } \Omega, \quad t \in (0, T]$$

Parabolic initial and boundary value problem (Uniqueness of solution)

Show that the problem

$$u_t - \Delta u = f \text{ in } \Omega, \quad t \in (0, T]$$
$$BC : \quad \frac{\partial u}{\partial \nu} = g \text{ on } \partial\Omega, \quad t \in [0, T]$$

Parabolic initial and boundary value problem (Uniqueness of solution)

Show that the problem

$$u_t - \Delta u = f \text{ in } \Omega, \quad t \in (0, T]$$

$$BC : \quad \frac{\partial u}{\partial \nu} = g \text{ on } \partial\Omega, \quad t \in [0, T]$$

$$IC : \quad u(x, 0) = u_0(x), \quad x \in \Omega$$

Parabolic initial and boundary value problem (Uniqueness of solution)

Show that the problem

$$u_t - \Delta u = f \text{ in } \Omega, \quad t \in (0, T]$$

$$BC : \quad \frac{\partial u}{\partial \nu} = g \text{ on } \partial\Omega, \quad t \in [0, T]$$

$$IC : \quad u(x, 0) = u_0(x), \quad x \in \Omega$$

has a unique solution.

Parabolic initial and boundary value problem (Uniqueness of solution)

Show that the problem

$$u_t - \Delta u = f \text{ in } \Omega, \quad t \in (0, T]$$

$$BC : \quad \frac{\partial u}{\partial \nu} = g \text{ on } \partial\Omega, \quad t \in [0, T]$$

$$IC : \quad u(x, 0) = u_0(x), \quad x \in \Omega$$

has a unique solution.

Suppose that the solution is not unique, that is, there are two distinct solutions u_i , $i = 1, 2$ with $u_1 \neq u_2$.

Parabolic initial and boundary value problem (Uniqueness of solution)

Show that the problem

$$u_t - \Delta u = f \text{ in } \Omega, \quad t \in (0, T]$$

$$BC : \quad \frac{\partial u}{\partial \nu} = g \text{ on } \partial\Omega, \quad t \in [0, T]$$

$$IC : \quad u(x, 0) = u_0(x), \quad x \in \Omega$$

has a unique solution.

Suppose that the solution is not unique, that is, there are two distinct solutions u_i , $i = 1, 2$ with $u_1 \neq u_2$.

Then, $v = u_1 - u_2$ satisfies

Parabolic initial and boundary value problem (Uniqueness of solution)

Show that the problem

$$u_t - \Delta u = f \text{ in } \Omega, \quad t \in (0, T]$$

$$BC : \quad \frac{\partial u}{\partial \nu} = g \text{ on } \partial\Omega, \quad t \in [0, T]$$

$$IC : \quad u(x, 0) = u_0(x), \quad x \in \Omega$$

has a unique solution.

Suppose that the solution is not unique, that is, there are two distinct solutions u_i , $i = 1, 2$ with $u_1 \neq u_2$.

Then, $v = u_1 - u_2$ satisfies

$$v_t - \Delta v = 0 \text{ in } \Omega, \quad t \in (0, T]$$

Parabolic initial and boundary value problem (Uniqueness of solution)

Show that the problem

$$u_t - \Delta u = f \text{ in } \Omega, \quad t \in (0, T]$$

$$BC : \quad \frac{\partial u}{\partial \nu} = g \text{ on } \partial\Omega, \quad t \in [0, T]$$

$$IC : \quad u(x, 0) = u_0(x), \quad x \in \Omega$$

has a unique solution.

Suppose that the solution is not unique, that is, there are two distinct solutions u_i , $i = 1, 2$ with $u_1 \neq u_2$.

Then, $v = u_1 - u_2$ satisfies

$$v_t - \Delta v = 0 \text{ in } \Omega, \quad t \in (0, T]$$

$$BC : \quad \frac{\partial v}{\partial \nu} = 0 \text{ on } \partial\Omega, \quad t \in [0, T]$$

Parabolic initial and boundary value problem (Uniqueness of solution)

Show that the problem

$$u_t - \Delta u = f \text{ in } \Omega, \quad t \in (0, T]$$

$$BC : \quad \frac{\partial u}{\partial \nu} = g \text{ on } \partial\Omega, \quad t \in [0, T]$$

$$IC : \quad u(x, 0) = u_0(x), \quad x \in \Omega$$

has a unique solution.

Suppose that the solution is not unique, that is, there are two distinct solutions u_i , $i = 1, 2$ with $u_1 \neq u_2$.

Then, $v = u_1 - u_2$ satisfies

$$v_t - \Delta v = 0 \text{ in } \Omega, \quad t \in (0, T]$$

$$BC : \quad \frac{\partial v}{\partial \nu} = 0 \text{ on } \partial\Omega, \quad t \in [0, T]$$

$$IC : \quad v(x, 0) = 0, \quad x \in \Omega$$

Parabolic initial and boundary value problem (Uniqueness of solution)

Show that the problem

$$u_t - \Delta u = f \text{ in } \Omega, \quad t \in (0, T]$$

$$BC : \quad \frac{\partial u}{\partial \nu} = g \text{ on } \partial\Omega, \quad t \in [0, T]$$

$$IC : \quad u(x, 0) = u_0(x), \quad x \in \Omega$$

has a unique solution.

Suppose that the solution is not unique, that is, there are two distinct solutions u_i , $i = 1, 2$ with $u_1 \neq u_2$.

Then, $v = u_1 - u_2$ satisfies

$$v_t - \Delta v = 0 \text{ in } \Omega, \quad t \in (0, T]$$

$$BC : \quad \frac{\partial v}{\partial \nu} = 0 \text{ on } \partial\Omega, \quad t \in [0, T]$$

$$IC : \quad v(x, 0) = 0, \quad x \in \Omega$$

Claim : $v(x, t) \equiv 0$.

Parabolic initial and boundary value problem (Uniqueness of solution)

Show that the problem

$$u_t - \Delta u = f \text{ in } \Omega, \quad t \in (0, T]$$

$$BC : \quad \frac{\partial u}{\partial \nu} = g \text{ on } \partial\Omega, \quad t \in [0, T]$$

$$IC : \quad u(x, 0) = u_0(x), \quad x \in \Omega$$

has a unique solution.

Suppose that the solution is not unique, that is, there are two distinct solutions u_i , $i = 1, 2$ with $u_1 \neq u_2$.

Then, $v = u_1 - u_2$ satisfies

$$v_t - \Delta v = 0 \text{ in } \Omega, \quad t \in (0, T]$$

$$BC : \quad \frac{\partial v}{\partial \nu} = 0 \text{ on } \partial\Omega, \quad t \in [0, T]$$

$$IC : \quad v(x, 0) = 0, \quad x \in \Omega$$

Claim : $v(x, t) \equiv 0$.

Multiply the PDE by v and integrate over Ω to obtain

Multiply the PDE by v and integrate over Ω to obtain

$$0 = \int_{\Omega} v_t v \, dx - \int_{\Omega} (\Delta v) v \, dx$$

Multiply the PDE by v and integrate over Ω to obtain

$$\begin{aligned} 0 &= \int_{\Omega} v_t v \, dx - \int_{\Omega} (\Delta v) v \, dx \\ &= \frac{1}{2} \frac{d}{dt} \left(\int_{\Omega} v(x, t)^2 \, dx \right) - \cancel{\int_{\partial\Omega} v \frac{\partial v}{\partial \nu} \, ds} + \int_{\Omega} |\nabla v(x, t)|^2 \, dx \end{aligned}$$

Multiply the PDE by v and integrate over Ω to obtain

$$\begin{aligned} 0 &= \int_{\Omega} v_t v \, dx - \int_{\Omega} (\Delta v) v \, dx \\ &= \frac{1}{2} \frac{d}{dt} \left(\int_{\Omega} v(x, t)^2 \, dx \right) - \cancel{\int_{\partial\Omega} v \frac{\partial v}{\partial \nu} \, ds} + \int_{\Omega} |\nabla v(x, t)|^2 \, dx \end{aligned}$$

Integrate from 0 to t to obtain

$$\int_{\Omega} v(x, t)^2 \, dx + 2 \int_0^t \int_{\Omega} |\nabla v(x, t)|^2 \, dx = 0$$

Multiply the PDE by v and integrate over Ω to obtain

$$\begin{aligned} 0 &= \int_{\Omega} v_t v \, dx - \int_{\Omega} (\Delta v) v \, dx \\ &= \frac{1}{2} \frac{d}{dt} \left(\int_{\Omega} v(x, t)^2 \, dx \right) - \cancel{\int_{\partial\Omega} v \frac{\partial v}{\partial \nu} \, ds} + \int_{\Omega} |\nabla v(x, t)|^2 \, dx \end{aligned}$$

Integrate from 0 to t to obtain

$$\int_{\Omega} v(x, t)^2 \, dx + 2 \int_0^t \int_{\Omega} |\nabla v(x, t)|^2 \, dx = 0 \quad (v(x, 0) = 0)$$

Multiply the PDE by v and integrate over Ω to obtain

$$\begin{aligned} 0 &= \int_{\Omega} v_t v \, dx - \int_{\Omega} (\Delta v) v \, dx \\ &= \frac{1}{2} \frac{d}{dt} \left(\int_{\Omega} v(x, t)^2 \, dx \right) - \cancel{\int_{\partial\Omega} v \frac{\partial v}{\partial \nu} \, ds} + \int_{\Omega} |\nabla v(x, t)|^2 \, dx \end{aligned}$$

Integrate from 0 to t to obtain

$$\begin{aligned} \int_{\Omega} v(x, t)^2 \, dx + 2 \int_0^t \int_{\Omega} |\nabla v(x, t)|^2 \, dx &= 0 \quad (v(x, 0) = 0) \\ \implies \int_{\Omega} v(x, t)^2 \, dx = 0 &\implies v(x, t) \equiv 0. \end{aligned}$$

$\implies u_1 = u_2$; hence the solution is unique.

Multiply the PDE by v and integrate over Ω to obtain

$$\begin{aligned} 0 &= \int_{\Omega} v_t v \, dx - \int_{\Omega} (\Delta v) v \, dx \\ &= \frac{1}{2} \frac{d}{dt} \left(\int_{\Omega} v(x, t)^2 \, dx \right) - \cancel{\int_{\partial\Omega} v \frac{\partial v}{\partial \nu} \, ds} + \int_{\Omega} |\nabla v(x, t)|^2 \, dx \end{aligned}$$

Integrate from 0 to t to obtain

$$\begin{aligned} \int_{\Omega} v(x, t)^2 \, dx + 2 \int_0^t \int_{\Omega} |\nabla v(x, t)|^2 \, dx &= 0 \quad (v(x, 0) = 0) \\ \implies \int_{\Omega} v(x, t)^2 \, dx = 0 &\implies v(x, t) \equiv 0. \end{aligned}$$

$\implies u_1 = u_2$; hence the solution is unique.

Exercise : Tut. Sheet 4- Qns. 7 (ii), (iii), (iv), 10

Hyperbolic initial and boundary value problem (Uniqueness of solution)

Exercise : Show that the problem

$$u_{tt} - \Delta u = f \text{ in } \Omega, \quad t \in (0, T]$$

$$BC : \quad \frac{\partial u}{\partial \nu} = g \text{ on } \partial\Omega, \quad t \in [0, T]$$

$$IC : \quad u(x, 0) = u_0(x), \quad u_t(x, 0) = u_1(x), \quad x \in \Omega$$

has a unique solution.

Exercise : Tut. Sheet 4 - Qn. 7(iv) , 9

Heat Conduction Equation (One space variable)

- Consider a homogeneous rod of length l .

Heat Conduction Equation (One space variable)

- Consider a homogeneous rod of length l .
- The rod is sufficiently thin so that the heat is distributed equally over the cross-section at time t .

Heat Conduction Equation (One space variable)

- Consider a homogeneous rod of length l .
- The rod is sufficiently thin so that the heat is distributed equally over the cross-section at time t .
- The surface of the rod is insulated and hence there is no heat loss through the boundary.

The **temperature distribution** of the rod is given by the solution of the initial-boundary value problem :

$$u_t = ku_{xx}, \quad 0 < x < l, \quad t > 0 \quad (k > 0 \text{ is the thermal conductivity})$$

Heat Conduction Equation (One space variable)

- Consider a homogeneous rod of length l .
- The rod is sufficiently thin so that the heat is distributed equally over the cross-section at time t .
- The surface of the rod is insulated and hence there is no heat loss through the boundary.

The **temperature distribution** of the rod is given by the solution of the initial-boundary value problem :

$$u_t = ku_{xx}, \quad 0 < x < l, \quad t > 0 \quad (k > 0 \text{ is the thermal conductivity})$$

$$BC : u(0, t) = 0, \quad u(l, t) = 0$$

Heat Conduction Equation (One space variable)

- Consider a homogeneous rod of length l .
- The rod is sufficiently thin so that the heat is distributed equally over the cross-section at time t .
- The surface of the rod is insulated and hence there is no heat loss through the boundary.

The **temperature distribution** of the rod is given by the solution of the initial-boundary value problem :

$$u_t = ku_{xx}, \quad 0 < x < l, \quad t > 0 \quad (k > 0 \text{ is the thermal conductivity})$$

$$BC : u(0, t) = 0, \quad u(l, t) = 0$$

$$IC : u(x, 0) = \phi(x) \quad 0 \leq x \leq l.$$

Method of Separation of Variables

$$u_t - ku_{xx} = 0, \quad l = (0, l), \quad t > 0$$

Method of Separation of Variables

$$u_t - ku_{xx} = 0, \quad l = (0, l), \quad t > 0$$

$$BC : u(0, t) = 0, \quad u(l, t) = 0$$

Method of Separation of Variables

$$u_t - ku_{xx} = 0, \quad I = (0, l), \quad t > 0$$

$$BC : u(0, t) = 0, \quad u(l, t) = 0$$

$$IC : u(x, 0) = \phi(x) \quad 0 \leq x \leq l.$$

Method of Separation of Variables

$$u_t - ku_{xx} = 0, \quad l = (0, l), \quad t > 0$$

$$BC : u(0, t) = 0, \quad u(l, t) = 0$$

$$IC : u(x, 0) = \phi(x) \quad 0 \leq x \leq l.$$

BOUNDED DOMAIN, HOMOGENEOUS BCs.

Assume that the solution $u(x, t)$ could be expressed as

$$u(x, t) = X(x)T(t)$$

Method of Separation of Variables

$$u_t - ku_{xx} = 0, \quad l = (0, l), \quad t > 0$$

$$BC : u(0, t) = 0, \quad u(l, t) = 0$$

$$IC : u(x, 0) = \phi(x) \quad 0 \leq x \leq l.$$

BOUNDED DOMAIN, HOMOGENEOUS BCs.

Assume that the solution $u(x, t)$ could be expressed as

$$u(x, t) = X(x)T(t)$$

Substituting in the heat equation, we obtain

Method of Separation of Variables

$$u_t - ku_{xx} = 0, \quad l = (0, l), \quad t > 0$$

$$BC : u(0, t) = 0, \quad u(l, t) = 0$$

$$IC : u(x, 0) = \phi(x) \quad 0 \leq x \leq l.$$

BOUNDED DOMAIN, HOMOGENEOUS BCs.

Assume that the solution $u(x, t)$ could be expressed as

$$u(x, t) = X(x)T(t)$$

Substituting in the heat equation, we obtain

$$X(x)T'(t) = kX''(x)T(t)$$

Method of Separation of Variables

$$u_t - ku_{xx} = 0, \quad l = (0, l), \quad t > 0$$

$$BC : u(0, t) = 0, \quad u(l, t) = 0$$

$$IC : u(x, 0) = \phi(x) \quad 0 \leq x \leq l.$$

BOUNDED DOMAIN, HOMOGENEOUS BCs.

Assume that the solution $u(x, t)$ could be expressed as

$$u(x, t) = X(x)T(t)$$

Substituting in the heat equation, we obtain

$$\begin{aligned} X(x)T'(t) &= kX''(x)T(t) \\ \implies \frac{X''(x)}{X(x)} &= \frac{T'(t)}{kT(t)} = -\lambda, \text{ (say)} \end{aligned}$$

Method of Separation of Variables

$$u_t - ku_{xx} = 0, \quad l = (0, l), \quad t > 0$$

$$BC : u(0, t) = 0, \quad u(l, t) = 0$$

$$IC : u(x, 0) = \phi(x) \quad 0 \leq x \leq l.$$

BOUNDED DOMAIN, HOMOGENEOUS BCs.

Assume that the solution $u(x, t)$ could be expressed as

$$u(x, t) = X(x)T(t)$$

Substituting in the heat equation, we obtain

$$\begin{aligned} X(x)T'(t) &= kX''(x)T(t) \\ \implies \frac{X''(x)}{X(x)} &= \frac{T'(t)}{kT(t)} = -\lambda, \text{ (say)} \end{aligned}$$

(Note that the first expression is a function of x alone, the second expression is a function of t alone, since x and t are independent variables, hence both sides must be equal to a constant.)

Method of Separation of Variables

$$u_t - ku_{xx} = 0, \quad l = (0, l), \quad t > 0$$

$$BC : u(0, t) = 0, \quad u(l, t) = 0$$

$$IC : u(x, 0) = \phi(x) \quad 0 \leq x \leq l.$$

BOUNDED DOMAIN, HOMOGENEOUS BCs.

Assume that the solution $u(x, t)$ could be expressed as

$$u(x, t) = X(x)T(t)$$

Substituting in the heat equation, we obtain

$$\begin{aligned} X(x)T'(t) &= kX''(x)T(t) \\ \implies \frac{X''(x)}{X(x)} &= \frac{T'(t)}{kT(t)} = -\lambda, \text{ (say)} \end{aligned}$$

(Note that the first expression is a function of x alone, the second expression is a function of t alone, since x and t are independent variables, hence both sides must be equal to a constant.)

$$\Rightarrow \frac{T'(t)}{kT(t)} = -\lambda, \quad \frac{X''(x)}{X(x)} = -\lambda;$$

$$\implies \frac{T'(t)}{kT(t)} = -\lambda, \quad \frac{X''(x)}{X(x)} = -\lambda;$$

That is, we have two problems:

$$\implies \frac{T'(t)}{kT(t)} = -\lambda, \quad \frac{X''(x)}{X(x)} = -\lambda;$$

That is, we have two problems:

$$X''(x) + \lambda X(x) = 0$$

$$T'(t) + \lambda k T(t) = 0$$

$$\implies \frac{T'(t)}{kT(t)} = -\lambda, \quad \frac{X''(x)}{X(x)} = -\lambda;$$

That is, we have two problems:

$$X''(x) + \lambda X(x) = 0$$

$$u(0, t) = u(l, t) = 0$$

$$T'(t) + \lambda k T(t) = 0$$

$$u(x, 0) = \phi(x)$$

$$\implies \frac{T'(t)}{kT(t)} = -\lambda, \quad \frac{X''(x)}{X(x)} = -\lambda;$$

That is, we have two problems:

$$X''(x) + \lambda X(x) = 0$$

$$u(0, t) = u(l, t) = 0$$

$$T'(t) + \lambda k T(t) = 0$$

$$u(x, 0) = \phi(x)$$

That is, $X(x)$ is the solution of the **eigenvalue problem** defined by:

$$\implies \frac{T'(t)}{kT(t)} = -\lambda, \quad \frac{X''(x)}{X(x)} = -\lambda;$$

That is, we have two problems:

$$X''(x) + \lambda X(x) = 0$$

$$u(0, t) = u(l, t) = 0$$

$$T'(t) + \lambda k T(t) = 0$$

$$u(x, 0) = \phi(x)$$

That is, $X(x)$ is the solution of the **eigenvalue problem** defined by:

$$X''(x) + \lambda X(x) = 0$$

$$\implies \frac{T'(t)}{kT(t)} = -\lambda, \quad \frac{X''(x)}{X(x)} = -\lambda;$$

That is, we have two problems:

$$X''(x) + \lambda X(x) = 0$$

$$u(0, t) = u(l, t) = 0$$

$$T'(t) + \lambda k T(t) = 0$$

$$u(x, 0) = \phi(x)$$

That is, $X(x)$ is the solution of the **eigenvalue problem** defined by:

$$X''(x) + \lambda X(x) = 0$$

$$X(0)T(t) = 0 \quad \forall t \implies X(0) = 0$$

$$\implies \frac{T'(t)}{kT(t)} = -\lambda, \quad \frac{X''(x)}{X(x)} = -\lambda;$$

That is, we have two problems:

$$X''(x) + \lambda X(x) = 0$$

$$u(0, t) = u(l, t) = 0$$

$$T'(t) + \lambda k T(t) = 0$$

$$u(x, 0) = \phi(x)$$

That is, $X(x)$ is the solution of the **eigenvalue problem** defined by:

$$X''(x) + \lambda X(x) = 0$$

$$X(0)T(t) = 0 \quad \forall t \implies X(0) = 0$$

$$X(l)T(t) = 0 \quad \forall t \implies X(l) = 0$$

$$\implies \frac{T'(t)}{kT(t)} = -\lambda, \quad \frac{X''(x)}{X(x)} = -\lambda;$$

That is, we have two problems:

$$X''(x) + \lambda X(x) = 0$$

$$u(0, t) = u(l, t) = 0$$

$$T'(t) + \lambda k T(t) = 0$$

$$u(x, 0) = \phi(x)$$

That is, $X(x)$ is the solution of the **eigenvalue problem** defined by:

$$X''(x) + \lambda X(x) = 0$$

$$X(0)T(t) = 0 \quad \forall t \implies X(0) = 0$$

$$X(l)T(t) = 0 \quad \forall t \implies X(l) = 0$$

The cases $\lambda = 0$ and $\lambda < 0$ give **only trivial solutions** to the EVP.

$$\implies \frac{T'(t)}{kT(t)} = -\lambda, \quad \frac{X''(x)}{X(x)} = -\lambda;$$

That is, we have two problems:

$$X''(x) + \lambda X(x) = 0$$

$$u(0, t) = u(l, t) = 0$$

$$T'(t) + \lambda k T(t) = 0$$

$$u(x, 0) = \phi(x)$$

That is, $X(x)$ is the solution of the **eigenvalue problem** defined by:

$$X''(x) + \lambda X(x) = 0$$

$$X(0)T(t) = 0 \quad \forall t \implies X(0) = 0$$

$$X(l)T(t) = 0 \quad \forall t \implies X(l) = 0$$

The cases $\lambda = 0$ and $\lambda < 0$ give **only trivial solutions** to the EVP.

$$\lambda > 0$$

$$X(x) = A \cos \sqrt{\lambda} x + B \sin \sqrt{\lambda} x$$

$$\lambda > 0$$

$$X(x) = A \cos \sqrt{\lambda} x + B \sin \sqrt{\lambda} x$$
$$X(0) = 0 \implies A = 0$$

$$\lambda > 0$$

$$X(x) = A \cos \sqrt{\lambda} x + B \sin \sqrt{\lambda} x$$

$$X(0) = 0 \implies A = 0$$

$$X(l) = 0 \implies B \sin \sqrt{\lambda} l = 0 \quad (B \neq 0)$$

$$\lambda > 0$$

$$X(x) = A \cos \sqrt{\lambda} x + B \sin \sqrt{\lambda} x$$

$$X(0) = 0 \implies A = 0$$

$$X(l) = 0 \implies B \sin \sqrt{\lambda} l = 0 \quad (B \neq 0)$$

$$\implies \sin \sqrt{\lambda} l = 0$$

$$\lambda > 0$$

$$X(x) = A \cos \sqrt{\lambda} x + B \sin \sqrt{\lambda} x$$

$$X(0) = 0 \implies A = 0$$

$$X(l) = 0 \implies B \sin \sqrt{\lambda} l = 0 \quad (B \neq 0)$$

$$\implies \sin \sqrt{\lambda} l = 0 \implies \sqrt{\lambda} l = n\pi \implies \lambda = \left(\frac{n\pi}{l}\right)^2.$$

$$\lambda > 0$$

$$X(x) = A \cos \sqrt{\lambda}x + B \sin \sqrt{\lambda}x$$

$$X(0) = 0 \implies A = 0$$

$$X(l) = 0 \implies B \sin \sqrt{\lambda}l = 0 \quad (B \neq 0)$$

$$\implies \sin \sqrt{\lambda}l = 0 \implies \sqrt{\lambda}l = n\pi \implies \lambda = \left(\frac{n\pi}{l}\right)^2.$$

Hence the eigenvalues are $\lambda = \left(\frac{n\pi}{l}\right)^2$

$$\lambda > 0$$

$$X(x) = A \cos \sqrt{\lambda}x + B \sin \sqrt{\lambda}x$$

$$X(0) = 0 \implies A = 0$$

$$X(l) = 0 \implies B \sin \sqrt{\lambda}l = 0 \quad (B \neq 0)$$

$$\implies \sin \sqrt{\lambda}l = 0 \implies \sqrt{\lambda}l = n\pi \implies \lambda = \left(\frac{n\pi}{l}\right)^2.$$

Hence the eigenvalues are $\lambda = \left(\frac{n\pi}{l}\right)^2$ and eigenfunctions are

$$X_n(x) = B_n \sin\left(\frac{n\pi x}{l}\right).$$

$$\lambda > 0$$

$$X(x) = A \cos \sqrt{\lambda}x + B \sin \sqrt{\lambda}x$$

$$X(0) = 0 \implies A = 0$$

$$X(l) = 0 \implies B \sin \sqrt{\lambda}l = 0 \quad (B \neq 0)$$

$$\implies \sin \sqrt{\lambda}l = 0 \implies \sqrt{\lambda}l = n\pi \implies \lambda = \left(\frac{n\pi}{l}\right)^2.$$

Hence the eigenvalues are $\lambda = \left(\frac{n\pi}{l}\right)^2$ and eigenfunctions are

$$X_n(x) = B_n \sin\left(\frac{n\pi x}{l}\right).$$

Now consider,

$$T'(t) + \lambda k T(t) = 0$$

Now consider,

$$T'(t) + \lambda k T(t) = 0$$

$$u(x, 0) = \phi(x)$$

Now consider,

$$T'(t) + \lambda k T(t) = 0$$

$$u(x, 0) = \phi(x)$$

Integrating, we obtain

$$\ln T(t) = -k\lambda t + \ln A' \implies T(t) = A'e^{-k\lambda t}.$$

Now consider,

$$T'(t) + \lambda k T(t) = 0$$

$$u(x, 0) = \phi(x)$$

Integrating, we obtain

$$\ln T(t) = -k\lambda t + \ln A' \implies T(t) = A' e^{-k\lambda t}.$$

$$\text{For each } \lambda_n, T_n(t) = A'_n e^{-k\lambda_n t} = A'_n e^{-k(\frac{n\pi}{l})^2 t}.$$

Now consider,

$$T'(t) + \lambda k T(t) = 0$$

$$u(x, 0) = \phi(x)$$

Integrating, we obtain

$$\ln T(t) = -k\lambda t + \ln A' \implies T(t) = A' e^{-k\lambda t}.$$

$$\text{For each } \lambda_n, T_n(t) = A'_n e^{-k\lambda_n t} = A'_n e^{-k(\frac{n\pi}{l})^2 t}.$$

Let $u_n(x, t) = X_n(x) T_n(t)$. Now, $\sum_{n=1}^N u_n(x, t)$ satisfies the heat equation.

Now consider,

$$T'(t) + \lambda k T(t) = 0$$

$$u(x, 0) = \phi(x)$$

Integrating, we obtain

$$\ln T(t) = -k\lambda t + \ln A' \implies T(t) = A' e^{-k\lambda t}.$$

$$\text{For each } \lambda_n, T_n(t) = A'_n e^{-k\lambda_n t} = A'_n e^{-k(\frac{n\pi}{l})^2 t}.$$

Let $u_n(x, t) = X_n(x) T_n(t)$. Now, $\sum_{n=1}^N u_n(x, t)$ satisfies the heat equation.

Under suitable assumptions on the initial data (which will be specified later), $u(x, t) = \sum_{n=1}^{\infty} u_n(x, t)$ also satisfies the heat equation.

$$u(x, t) = \sum_{n=1}^{\infty} u_n(x, t)$$

ASSUMPTION (A1)

$$\begin{aligned} u(x, t) &= \sum_{n=1}^{\infty} u_n(x, t) && \text{ASSUMPTION (A1)} \\ &= \sum_{n=1}^{\infty} X_n(x) T_n(t) \end{aligned}$$

$$\begin{aligned}
 u(x, t) &= \sum_{n=1}^{\infty} u_n(x, t) && \text{ASSUMPTION (A1)} \\
 &= \sum_{n=1}^{\infty} X_n(x) T_n(t) \\
 &= \sum_{n=1}^{\infty} A_n \sin\left(\frac{n\pi x}{l}\right) e^{-k\left(\frac{n\pi}{l}\right)^2 t} && (A_n = A'_n B_n)
 \end{aligned}$$

$$\begin{aligned}
 u(x, t) &= \sum_{n=1}^{\infty} u_n(x, t) && \text{ASSUMPTION (A1)} \\
 &= \sum_{n=1}^{\infty} X_n(x) T_n(t) \\
 &= \sum_{n=1}^{\infty} A_n \sin\left(\frac{n\pi x}{l}\right) e^{-k\left(\frac{n\pi}{l}\right)^2 t} && (A_n = A'_n B_n)
 \end{aligned}$$

$$u(x, 0) = \phi(x) \implies \sum_{n=1}^{\infty} A_n \sin\left(\frac{n\pi x}{l}\right) = \phi(x) \quad \text{ASSUMPTION (A2) -}$$

Fourier sine series of $\phi(x)$ converges to $\phi(x)$.

$$\begin{aligned}
 u(x, t) &= \sum_{n=1}^{\infty} u_n(x, t) && \text{ASSUMPTION (A1)} \\
 &= \sum_{n=1}^{\infty} X_n(x) T_n(t) \\
 &= \sum_{n=1}^{\infty} A_n \sin\left(\frac{n\pi x}{l}\right) e^{-k\left(\frac{n\pi}{l}\right)^2 t} && (A_n = A'_n B_n)
 \end{aligned}$$

$$u(x, 0) = \phi(x) \implies \sum_{n=1}^{\infty} A_n \sin\left(\frac{n\pi x}{l}\right) = \phi(x) \quad \text{ASSUMPTION (A2) -}$$

Fourier sine series of $\phi(x)$ converges to $\phi(x)$.

Hence,
$$A_n = \frac{2}{l} \int_0^l \phi(x) \sin\left(\frac{n\pi x}{l}\right) dx.$$

$$\begin{aligned}
 u(x, t) &= \sum_{n=1}^{\infty} u_n(x, t) && \text{ASSUMPTION (A1)} \\
 &= \sum_{n=1}^{\infty} X_n(x) T_n(t) \\
 &= \sum_{n=1}^{\infty} A_n \sin\left(\frac{n\pi x}{l}\right) e^{-k\left(\frac{n\pi}{l}\right)^2 t} \quad (A_n = A'_n B_n)
 \end{aligned}$$

$$u(x, 0) = \phi(x) \implies \sum_{n=1}^{\infty} A_n \sin\left(\frac{n\pi x}{l}\right) = \phi(x) \quad \text{ASSUMPTION (A2) -}$$

Fourier sine series of $\phi(x)$ converges to $\phi(x)$.

Hence, $A_n = \frac{2}{l} \int_0^l \phi(x) \sin\left(\frac{n\pi x}{l}\right) dx.$ The formal solution of the

heat equation with Dirichlet BC & IC is given by

$$u(x, t) = \sum_{n=1}^{\infty} \left(\frac{2}{l} \int_0^l \phi(x) \sin\left(\frac{n\pi x}{l}\right) dx \right) \sin\left(\frac{n\pi x}{l}\right) e^{-k\left(\frac{n\pi}{l}\right)^2 t}.$$

Justification of (A1) & (A2)

If the function $\phi(x)$ satisfies the properties:

Justification of (A1) & (A2)

If the function $\phi(x)$ satisfies the properties:

- 1 $\phi(x)$ is continuous in $[0, l]$,

Justification of (A1) & (A2)

If the function $\phi(x)$ satisfies the properties:

- 1 $\phi(x)$ is continuous in $[0, l]$,
- 2 $\phi(0) = \phi(l) = 0$

Justification of (A1) & (A2)

If the function $\phi(x)$ satisfies the properties:

- 1 $\phi(x)$ is continuous in $[0, l]$,
- 2 $\phi(0) = \phi(l) = 0$
- 3 $\phi'(x)$ is piecewise continuous in $(0, l)$,

Justification of (A1) & (A2)

If the function $\phi(x)$ satisfies the properties:

- 1 $\phi(x)$ is continuous in $[0, l]$,
- 2 $\phi(0) = \phi(l) = 0$
- 3 $\phi'(x)$ is piecewise continuous in $(0, l)$,

then assumptions (A1) and (A2) are satisfied so that the formal solution is in fact a solution.

Justification of (A1) & (A2)

If the function $\phi(x)$ satisfies the properties:

- ① $\phi(x)$ is continuous in $[0, l]$,
- ② $\phi(0) = \phi(l) = 0$
- ③ $\phi'(x)$ is piecewise continuous in $(0, l)$,

then assumptions (A1) and (A2) are satisfied so that the formal solution is in fact a solution.

Exercise :

1. Is the solution unique?

Justification of (A1) & (A2)

If the function $\phi(x)$ satisfies the properties:

- ① $\phi(x)$ is continuous in $[0, l]$,
- ② $\phi(0) = \phi(l) = 0$
- ③ $\phi'(x)$ is piecewise continuous in $(0, l)$,

then assumptions (A1) and (A2) are satisfied so that the formal solution is in fact a solution.

Exercise :

1. Is the solution unique?
2. Tut. Sheet 5 : 1, 2, 3, 10 (i), (iii)

Justification of (A1) & (A2)

If the function $\phi(x)$ satisfies the properties:

- ① $\phi(x)$ is continuous in $[0, l]$,
- ② $\phi(0) = \phi(l) = 0$
- ③ $\phi'(x)$ is piecewise continuous in $(0, l)$,

then assumptions (A1) and (A2) are satisfied so that the formal solution is in fact a solution.

Exercise :

1. Is the solution unique?
2. Tut. Sheet 5 : 1, 2, 3, 10 (i), (iii)

Heat Equation - Non-homogeneous Boundary Data

$$u_t - ku_{xx} = 0, \quad I = (0, l), \quad t > 0$$

Heat Equation - Non-homogeneous Boundary Data

$$u_t - ku_{xx} = 0, \quad I = (0, l), \quad t > 0$$

$$BC : u(0, t) = g(t), \quad u(l, t) = h(t)$$

Heat Equation - Non-homogeneous Boundary Data

$$u_t - ku_{xx} = 0, \quad I = (0, l), \quad t > 0$$

$$BC : u(0, t) = g(t), \quad u(l, t) = h(t)$$

$$IC : u(x, 0) = \phi(x) \quad 0 \leq x \leq l.$$

Heat Equation - Non-homogeneous Boundary Data

$$u_t - ku_{xx} = 0, \quad I = (0, l), \quad t > 0$$

$$BC : u(0, t) = g(t), \quad u(l, t) = h(t)$$

$$IC : u(x, 0) = \phi(x) \quad 0 \leq x \leq l.$$

Introduce a new function :

Heat Equation - Non-homogeneous Boundary Data

$$u_t - ku_{xx} = 0, \quad I = (0, l), \quad t > 0$$

$$BC : u(0, t) = g(t), \quad u(l, t) = h(t)$$

$$IC : u(x, 0) = \phi(x) \quad 0 \leq x \leq l.$$

Introduce a new function :

$$U(x, t) = \frac{1}{l} \left[(l - x)g(t) + xh(t) \right].$$

Heat Equation - Non-homogeneous Boundary Data

$$u_t - ku_{xx} = 0, \quad I = (0, l), \quad t > 0$$

$$BC : u(0, t) = g(t), \quad u(l, t) = h(t)$$

$$IC : u(x, 0) = \phi(x) \quad 0 \leq x \leq l.$$

Introduce a new function :

$$U(x, t) = \frac{1}{l} \left[(l - x)g(t) + xh(t) \right]. \quad \text{Note that } U(x, t) = g(t) \text{ at } x = 0 \text{ and } U(x, t) = h(t) \text{ at } x = l.$$

Heat Equation - Non-homogeneous Boundary Data

$$u_t - ku_{xx} = 0, \quad I = (0, l), \quad t > 0$$

$$BC : u(0, t) = g(t), \quad u(l, t) = h(t)$$

$$IC : u(x, 0) = \phi(x) \quad 0 \leq x \leq l.$$

Introduce a new function :

$$U(x, t) = \frac{1}{l} \left[(l - x)g(t) + xh(t) \right]. \quad \text{Note that } U(x, t) = g(t) \text{ at}$$

$x = 0$ and $U(x, t) = h(t)$ at $x = l$.

Let $v(x, t) = u(x, t) - U(x, t)$.

Heat Equation - Non-homogeneous Boundary Data

$$u_t - ku_{xx} = 0, \quad I = (0, l), \quad t > 0$$

$$BC : u(0, t) = g(t), \quad u(l, t) = h(t)$$

$$IC : u(x, 0) = \phi(x) \quad 0 \leq x \leq l.$$

Introduce a new function :

$$U(x, t) = \frac{1}{l} \left[(l - x)g(t) + xh(t) \right]. \quad \text{Note that } U(x, t) = g(t) \text{ at}$$

$x = 0$ and $U(x, t) = h(t)$ at $x = l$.

Let $v(x, t) = u(x, t) - U(x, t)$.

Therefore,

$$v_t - kv_{xx} = (u_t - ku_{xx}) - (U_t - kU_{xx})$$

Heat Equation - Non-homogeneous Boundary Data

$$u_t - ku_{xx} = 0, \quad I = (0, l), \quad t > 0$$

$$BC : u(0, t) = g(t), \quad u(l, t) = h(t)$$

$$IC : u(x, 0) = \phi(x) \quad 0 \leq x \leq l.$$

Introduce a new function :

$$U(x, t) = \frac{1}{l} \left[(l-x)g(t) + xh(t) \right]. \quad \text{Note that } U(x, t) = g(t) \text{ at}$$

$x = 0$ and $U(x, t) = h(t)$ at $x = l$.

Let $v(x, t) = u(x, t) - U(x, t)$.

Therefore,

$$\begin{aligned} v_t - kv_{xx} &= (u_t - ku_{xx}) - (U_t - kU_{xx}) \\ &= 0 - \left[\frac{1}{l} (l-x)g'(t) + xh'(t) \right] \quad (U_{xx} = 0) \end{aligned}$$

$$\text{Hence, } v_t - kv_{xx} = -\left[\frac{1}{l}(l-x)g'(t) + xh'(t)\right]$$

Hence, $v_t - kv_{xx} = - \left[\frac{1}{l}(l-x)g'(t) + xh'(t) \right]$

$$v(x,0) = u(x,0) - U(x,0)$$

Hence, $v_t - kv_{xx} = -\left[\frac{1}{l}(l-x)g'(t) + xh'(t)\right]$

$$\begin{aligned} v(x, 0) &= u(x, 0) - U(x, 0) \\ &= \phi(x) - \left[\frac{1}{l}(l-x)g(0) + xh(0)\right]. \end{aligned}$$

$$\text{Hence, } v_t - kv_{xx} = - \left[\frac{1}{l}(l-x)g'(t) + xh'(t) \right]$$

$$\begin{aligned} v(x, 0) &= u(x, 0) - U(x, 0) \\ &= \phi(x) - \left[\frac{1}{l}(l-x)g(0) + xh(0) \right]. \end{aligned}$$

$$v(0, t) = u(0, t) - U(0, t)$$

Hence, $v_t - kv_{xx} = -\left[\frac{1}{l}(l-x)g'(t) + xh'(t)\right]$

$$\begin{aligned} v(x, 0) &= u(x, 0) - U(x, 0) \\ &= \phi(x) - \left[\frac{1}{l}(l-x)g(0) + xh(0)\right]. \end{aligned}$$

$$\begin{aligned} v(0, t) &= u(0, t) - U(0, t) \\ &= g(t) - g(t) = 0. \end{aligned}$$

$$\text{Hence, } v_t - kv_{xx} = - \left[\frac{1}{l}(l-x)g'(t) + xh'(t) \right]$$

$$\begin{aligned} v(x, 0) &= u(x, 0) - U(x, 0) \\ &= \phi(x) - \left[\frac{1}{l}(l-x)g(0) + xh(0) \right]. \end{aligned}$$

$$\begin{aligned} v(0, t) &= u(0, t) - U(0, t) \\ &= g(t) - g(t) = 0. \end{aligned}$$

$$v(l, t) = u(l, t) - U(l, t) = 0$$

Therefore, $v(x, t)$ is a solution of the **non-homogeneous heat equation** on $[0, l]$ with **homogeneous boundary data**,

$$v_t - kv_{xx} = - \left[\frac{1}{l}(l-x)g'(t) + xh'(t) \right]$$

$$\text{Hence, } v_t - kv_{xx} = - \left[\frac{1}{l}(l-x)g'(t) + xh'(t) \right]$$

$$\begin{aligned} v(x, 0) &= u(x, 0) - U(x, 0) \\ &= \phi(x) - \left[\frac{1}{l}(l-x)g(0) + xh(0) \right]. \end{aligned}$$

$$\begin{aligned} v(0, t) &= u(0, t) - U(0, t) \\ &= g(t) - g(t) = 0. \end{aligned}$$

$$v(l, t) = u(l, t) - U(l, t) = 0$$

Therefore, $v(x, t)$ is a solution of the **non-homogeneous heat equation** on $[0, l]$ with **homogeneous boundary data**,

$$\begin{aligned} v_t - kv_{xx} &= - \left[\frac{1}{l}(l-x)g'(t) + xh'(t) \right] \\ v(x, 0) &= \phi(x) - \left[\frac{1}{l}(l-x)g(0) + xh(0) \right] \end{aligned}$$

$$\text{Hence, } v_t - kv_{xx} = -\left[\frac{1}{l}(l-x)g'(t) + xh'(t)\right]$$

$$\begin{aligned} v(x, 0) &= u(x, 0) - U(x, 0) \\ &= \phi(x) - \left[\frac{1}{l}(l-x)g(0) + xh(0)\right]. \end{aligned}$$

$$\begin{aligned} v(0, t) &= u(0, t) - U(0, t) \\ &= g(t) - g(t) = 0. \end{aligned}$$

$$v(l, t) = u(l, t) - U(l, t) = 0$$

Therefore, $v(x, t)$ is a solution of the **non-homogeneous heat equation** on $[0, l]$ with **homogeneous boundary data**,

$$\begin{aligned} v_t - kv_{xx} &= -\left[\frac{1}{l}(l-x)g'(t) + xh'(t)\right] \\ v(x, 0) &= \phi(x) - \left[\frac{1}{l}(l-x)g(0) + xh(0)\right] \\ v(0, t) &= v(l, t) = 0. \end{aligned}$$

$$\text{Hence, } v_t - kv_{xx} = - \left[\frac{1}{l}(l-x)g'(t) + xh'(t) \right]$$

$$\begin{aligned} v(x, 0) &= u(x, 0) - U(x, 0) \\ &= \phi(x) - \left[\frac{1}{l}(l-x)g(0) + xh(0) \right]. \end{aligned}$$

$$\begin{aligned} v(0, t) &= u(0, t) - U(0, t) \\ &= g(t) - g(t) = 0. \end{aligned}$$

$$v(l, t) = u(l, t) - U(l, t) = 0$$

Therefore, $v(x, t)$ is a solution of the **non-homogeneous heat equation** on $[0, l]$ with **homogeneous boundary data**,

$$v_t - kv_{xx} = - \left[\frac{1}{l}(l-x)g'(t) + xh'(t) \right]$$

$$v(x, 0) = \phi(x) - \left[\frac{1}{l}(l-x)g(0) + xh(0) \right]$$

$$v(0, t) = v(l, t) = 0.$$

$$u(x, t) = v(x, t) + U(x, t)$$

Non-homogeneous Heat Equation- Diagonalization Method

$$u_t - ku_{xx} = h(x, t), \quad 0 < x < l, \quad t > 0$$

Non-homogeneous Heat Equation- Diagonalization Method

$$u_t - ku_{xx} = h(x, t), \quad 0 < x < l, \quad t > 0$$

$$BC : u(0, t) = 0, \quad u(l, t) = 0$$

Non-homogeneous Heat Equation- Diagonalization Method

$$u_t - ku_{xx} = h(x, t), \quad 0 < x < l, \quad t > 0$$

$$BC : u(0, t) = 0, \quad u(l, t) = 0$$

$$IC : u(x, 0) = \phi(x) \quad 0 \leq x \leq l.$$

$h(x, t) \neq 0$ - (non-homogeneous) Consider the following eigenvalue problem:

$$y'' + \lambda y = 0, \quad 0 < x < l,$$

Non-homogeneous Heat Equation- Diagonalization Method

$$u_t - ku_{xx} = h(x, t), \quad 0 < x < l, \quad t > 0$$

$$BC : u(0, t) = 0, \quad u(l, t) = 0$$

$$IC : u(x, 0) = \phi(x) \quad 0 \leq x \leq l.$$

$h(x, t) \neq 0$ - (non-homogeneous) Consider the following eigenvalue problem:

$$y'' + \lambda y = 0, \quad 0 < x < l,$$

$$y(0) = y(l) = 0$$

Non-homogeneous Heat Equation- Diagonalization Method

$$u_t - ku_{xx} = h(x, t), \quad 0 < x < l, \quad t > 0$$

$$BC : u(0, t) = 0, \quad u(l, t) = 0$$

$$IC : u(x, 0) = \phi(x) \quad 0 \leq x \leq l.$$

$h(x, t) \neq 0$ - (non-homogeneous) Consider the following eigenvalue problem:

$$y'' + \lambda y = 0, \quad 0 < x < l,$$

$$y(0) = y(l) = 0$$

We know that $\lambda_n = \left(\frac{n\pi}{l}\right)^2$ are the eigenvalues and

$y_n = \sqrt{\frac{2}{l}} \sin\left(\frac{n\pi}{l}x\right)$ are the corresponding eigenfunctions.

Non-homogeneous Heat Equation- Diagonalization Method

$$u_t - ku_{xx} = h(x, t), \quad 0 < x < l, \quad t > 0$$

$$BC : u(0, t) = 0, \quad u(l, t) = 0$$

$$IC : u(x, 0) = \phi(x) \quad 0 \leq x \leq l.$$

$h(x, t) \neq 0$ - (non-homogeneous) Consider the following eigenvalue problem:

$$y'' + \lambda y = 0, \quad 0 < x < l,$$

$$y(0) = y(l) = 0$$

We know that $\lambda_n = \left(\frac{n\pi}{l}\right)^2$ are the eigenvalues and

$y_n = \sqrt{\frac{2}{l}} \sin\left(\frac{n\pi}{l}x\right)$ are the corresponding eigenfunctions.

Expand $h(x, t) = \sum_{n=1}^{\infty} \gamma_n(t) y_n(x)$

Non-homogeneous Heat Equation- Diagonalization Method

$$u_t - ku_{xx} = h(x, t), \quad 0 < x < l, \quad t > 0$$

$$BC : u(0, t) = 0, \quad u(l, t) = 0$$

$$IC : u(x, 0) = \phi(x) \quad 0 \leq x \leq l.$$

$h(x, t) \neq 0$ - (non-homogeneous) Consider the following eigenvalue problem:

$$y'' + \lambda y = 0, \quad 0 < x < l,$$

$$y(0) = y(l) = 0$$

We know that $\lambda_n = \left(\frac{n\pi}{l}\right)^2$ are the eigenvalues and

$y_n = \sqrt{\frac{2}{l}} \sin\left(\frac{n\pi}{l}x\right)$ are the corresponding eigenfunctions.

Expand $h(x, t) = \sum_{n=1}^{\infty} \gamma_n(t) y_n(x)$ and $u(x, t) = \sum_{n=1}^{\infty} \alpha_n(t) y_n(x)$.

Non-homogeneous Heat Equation- Diagonalization Method

$$u_t - ku_{xx} = h(x, t), \quad 0 < x < l, \quad t > 0$$

$$BC : u(0, t) = 0, \quad u(l, t) = 0$$

$$IC : u(x, 0) = \phi(x) \quad 0 \leq x \leq l.$$

$h(x, t) \neq 0$ - (non-homogeneous) Consider the following eigenvalue problem:

$$y'' + \lambda y = 0, \quad 0 < x < l,$$

$$y(0) = y(l) = 0$$

We know that $\lambda_n = \left(\frac{n\pi}{l}\right)^2$ are the eigenvalues and

$y_n = \sqrt{\frac{2}{l}} \sin\left(\frac{n\pi}{l}x\right)$ are the corresponding eigenfunctions.

Expand $h(x, t) = \sum_{n=1}^{\infty} \gamma_n(t) y_n(x)$ and $u(x, t) = \sum_{n=1}^{\infty} \alpha_n(t) y_n(x)$.

Here $\gamma_n(t)$ can be computed as the Fourier coefficients and $\alpha_n(t)$ are unknown functions.

Here $\gamma_n(t)$ can be computed as the Fourier coefficients and $\alpha_n(t)$ are unknown functions.

Plug these expressions in the PDE to obtain

Here $\gamma_n(t)$ can be computed as the Fourier coefficients and $\alpha_n(t)$ are unknown functions.

Plug these expressions in the PDE to obtain

$$\sum_{n=1}^{\infty} \frac{d}{dt} \alpha_n(t) y_n(x) - k \sum_{n=1}^{\infty} \alpha_n(t) y_n''(x) = \sum_{n=1}^{\infty} \gamma_n(t) y_n(x).$$

Here $\gamma_n(t)$ can be computed as the Fourier coefficients and $\alpha_n(t)$ are unknown functions.

Plug these expressions in the PDE to obtain

$$\sum_{n=1}^{\infty} \frac{d}{dt} \alpha_n(t) y_n(x) - k \sum_{n=1}^{\infty} \alpha_n(t) y_n''(x) = \sum_{n=1}^{\infty} \gamma_n(t) y_n(x).$$

Now $y_n''(x) = -\lambda_n y_n(x).$

Here $\gamma_n(t)$ can be computed as the Fourier coefficients and $\alpha_n(t)$ are unknown functions.

Plug these expressions in the PDE to obtain

$$\sum_{n=1}^{\infty} \frac{d}{dt} \alpha_n(t) y_n(x) - k \sum_{n=1}^{\infty} \alpha_n(t) y_n''(x) = \sum_{n=1}^{\infty} \gamma_n(t) y_n(x).$$

Now $y_n''(x) = -\lambda_n y_n(x)$.

On substitution, we obtain

Here $\gamma_n(t)$ can be computed as the Fourier coefficients and $\alpha_n(t)$ are unknown functions.

Plug these expressions in the PDE to obtain

$$\sum_{n=1}^{\infty} \frac{d}{dt} \alpha_n(t) y_n(x) - k \sum_{n=1}^{\infty} \alpha_n(t) y_n''(x) = \sum_{n=1}^{\infty} \gamma_n(t) y_n(x).$$

Now $y_n''(x) = -\lambda_n y_n(x)$.

On substitution, we obtain

$$\sum_{n=1}^{\infty} \left(\frac{d}{dt} \alpha_n(t) + k \lambda_n \alpha_n(t) - \gamma_n(t) \right) y_n(t) = 0$$

Here $\gamma_n(t)$ can be computed as the Fourier coefficients and $\alpha_n(t)$ are unknown functions.

Plug these expressions in the PDE to obtain

$$\sum_{n=1}^{\infty} \frac{d}{dt} \alpha_n(t) y_n(x) - k \sum_{n=1}^{\infty} \alpha_n(t) y_n''(x) = \sum_{n=1}^{\infty} \gamma_n(t) y_n(x).$$

Now $y_n''(x) = -\lambda_n y_n(x)$.

On substitution, we obtain

$$\sum_{n=1}^{\infty} \left(\frac{d}{dt} \alpha_n(t) + k \lambda_n \alpha_n(t) - \gamma_n(t) \right) y_n(t) = 0$$

Taking innerproduct with $y_m(t)$ and using orthonormality of eigen functions, we obtain

Here $\gamma_n(t)$ can be computed as the Fourier coefficients and $\alpha_n(t)$ are unknown functions.

Plug these expressions in the PDE to obtain

$$\sum_{n=1}^{\infty} \frac{d}{dt} \alpha_n(t) y_n(x) - k \sum_{n=1}^{\infty} \alpha_n(t) y_n''(x) = \sum_{n=1}^{\infty} \gamma_n(t) y_n(x).$$

Now $y_n''(x) = -\lambda_n y_n(x)$.

On substitution, we obtain

$$\sum_{n=1}^{\infty} \left(\frac{d}{dt} \alpha_n(t) + k \lambda_n \alpha_n(t) - \gamma_n(t) \right) y_n(t) = 0$$

Taking innerproduct with $y_m(t)$ and using orthonormality of eigen functions, we obtain

$$\frac{d}{dt} \alpha_n(t) + k \lambda_n \alpha_n(t) = \gamma_n(t), \quad n = 1, 2, 3, \dots,$$

Here $\gamma_n(t)$ can be computed as the Fourier coefficients and $\alpha_n(t)$ are unknown functions.

Plug these expressions in the PDE to obtain

$$\sum_{n=1}^{\infty} \frac{d}{dt} \alpha_n(t) y_n(x) - k \sum_{n=1}^{\infty} \alpha_n(t) y_n''(x) = \sum_{n=1}^{\infty} \gamma_n(t) y_n(x).$$

Now $y_n''(x) = -\lambda_n y_n(x)$.

On substitution, we obtain

$$\sum_{n=1}^{\infty} \left(\frac{d}{dt} \alpha_n(t) + k \lambda_n \alpha_n(t) - \gamma_n(t) \right) y_n(t) = 0$$

Taking innerproduct with $y_m(t)$ and using orthonormality of eigen functions, we obtain

$$\frac{d}{dt} \alpha_n(t) + k \lambda_n \alpha_n(t) = \gamma_n(t), \quad n = 1, 2, 3, \dots,$$

$$\implies \alpha_n(t) = \alpha_n(0) e^{-\lambda_n t} + \int_0^t e^{-\lambda_n(t-s)} \gamma_n(s) ds$$

Here $\gamma_n(t)$ can be computed as the Fourier coefficients and $\alpha_n(t)$ are unknown functions.

Plug these expressions in the PDE to obtain

$$\sum_{n=1}^{\infty} \frac{d}{dt} \alpha_n(t) y_n(x) - k \sum_{n=1}^{\infty} \alpha_n(t) y_n''(x) = \sum_{n=1}^{\infty} \gamma_n(t) y_n(x).$$

Now $y_n''(x) = -\lambda_n y_n(x)$.

On substitution, we obtain

$$\sum_{n=1}^{\infty} \left(\frac{d}{dt} \alpha_n(t) + k \lambda_n \alpha_n(t) - \gamma_n(t) \right) y_n(t) = 0$$

Taking innerproduct with $y_m(t)$ and using orthonormality of eigen functions, we obtain

$$\frac{d}{dt} \alpha_n(t) + k \lambda_n \alpha_n(t) = \gamma_n(t), \quad n = 1, 2, 3, \dots,$$

$$\implies \alpha_n(t) = \alpha_n(0) e^{-\lambda_n t} + \int_0^t e^{-\lambda_n(t-s)} \gamma_n(s) ds$$

To find $\alpha_n(0)$ using the IC :

To find $\alpha_n(0)$ using the IC :

$$u(x, 0) = \phi(x) \implies \phi(x) = \sum_{n=1}^{\infty} \alpha_n(0) y_n(x)$$

where $\alpha_n(0) = \int_0^l \phi(x) y_n(x) dx.$

To find $\alpha_n(0)$ using the IC :

$$u(x, 0) = \phi(x) \implies \phi(x) = \sum_{n=1}^{\infty} \alpha_n(0) y_n(x)$$

where $\alpha_n(0) = \int_0^l \phi(x) y_n(x) dx$.

Hence $u(x, t) = \sum_{n=1}^{\infty} \alpha_n(t) y_n(x)$ with

To find $\alpha_n(0)$ using the IC :

$$u(x, 0) = \phi(x) \implies \phi(x) = \sum_{n=1}^{\infty} \alpha_n(0) y_n(x)$$

where $\alpha_n(0) = \int_0^l \phi(x) y_n(x) dx.$

Hence $u(x, t) = \sum_{n=1}^{\infty} \alpha_n(t) y_n(x)$ with

$$\implies \alpha_n(t) = \alpha_n(0) e^{-\lambda_n t} + \int_0^t e^{-\lambda_n(t-s)} \gamma_n(s) ds. \text{ (CHECK !)}$$

Exercise : Tut. Sheet 5 - Qns. 7, 8, 9 (i)