

Signal Processing - 1 by One

Sibi Raj B. Pillai
Dept of Electrical Engineering
IIT Bombay



- Impulse Replacement Operation
- Generalization
- Examples
- Digital Convolution
- Analog Domain



Outline

- Impulse Replacement Operation
- Generalization
- Examples
- Digital Convolution
- Analog Domain



Outline

- Impulse Replacement Operation
- Generalization
- **Examples**
- Digital Convolution
- Analog Domain



Outline

- Impulse Replacement Operation
- Generalization
- Examples
- Digital Convolution
- Analog Domain





Outline

- Impulse Replacement Operation
- Generalization
- Examples
- Digital Convolution
- Analog Domain



Dirac Formalism for Replacement





$$\delta(t - \tau) * x(t) = x(t - \tau)$$

$$\delta(t) * x(t - \tau) = x(t - \tau)$$

$$f(t) * \delta(t - \tau) = ?$$



Dirac Formalism for Replacement



$$\delta(t - \tau) * x(t) = x(t - \tau)$$

$$\delta(t) * x(t - \tau) = x(t - \tau)$$

$$f(t) * \delta(t - \tau) = ?$$



Desiderata on $*$

- ▶ $*$ should be commutative, i.e. $x(t) * y(t) = y(t) * x(t)$.
- ▶ $*$ with an impulse at τ should yield the function $x(t - \tau)$.
- ▶ Associativity: $(x(t) * y(t)) * z(t) = x(t) * (y(t) * z(t))$.
- ▶ Distribution over addition:

$$x(t) * (y(t) + z(t)) = x(t) * y(t) + x(t) * z(t).$$

- ▶ Consistency when $\delta(t - \tau)$ is used in place of any signal(s).



Desiderata on $*$

- ▶ $*$ should be commutative, i.e. $x(t) * y(t) = y(t) * x(t)$.
- ▶ $*$ with an impulse at τ should yield the function $x(t - \tau)$.

▶ Associativity: $(x(t) * y(t)) * z(t) = x(t) * (y(t) * z(t))$.

▶ Distribution over addition:

$$x(t) * (y(t) + z(t)) = x(t) * y(t) + x(t) * z(t).$$

▶ Consistency when $\delta(t - \tau)$ is used in place of any signal(s).



Desiderata on $*$

- ▶ $*$ should be commutative, i.e. $x(t) * y(t) = y(t) * x(t)$.
- ▶ $*$ with an impulse at τ should yield the function $x(t - \tau)$.
- ▶ Associativity: $(x(t) * y(t)) * z(t) = x(t) * (y(t) * z(t))$.
- ▶ Distribution over addition:

$$x(t) * (y(t) + z(t)) = x(t) * y(t) + x(t) * z(t).$$

- ▶ Consistency when $\delta(t - \tau)$ is used in place of any signal(s).



Desiderata on $*$

- ▶ $*$ should be commutative, i.e. $x(t) * y(t) = y(t) * x(t)$.
- ▶ $*$ with an impulse at τ should yield the function $x(t - \tau)$.
- ▶ Associativity: $(x(t) * y(t)) * z(t) = x(t) * (y(t) * z(t))$.
- ▶ Distribution over addition:

$$x(t) * (y(t) + z(t)) = x(t) * y(t) + x(t) * z(t).$$

- ▶ Consistency when $\delta(t - \tau)$ is used in place of any signal(s).



Desiderata on $*$

- ▶ $*$ should be commutative, i.e. $x(t) * y(t) = y(t) * x(t)$.
- ▶ $*$ with an impulse at τ should yield the function $x(t - \tau)$.
- ▶ Associativity: $(x(t) * y(t)) * z(t) = x(t) * (y(t) * z(t))$.
- ▶ Distribution over addition:

$$x(t) * (y(t) + z(t)) = x(t) * y(t) + x(t) * z(t).$$

- ▶ Consistency when $\delta(t - \tau)$ is used in place of any signal(s).



Desiderata on $*$

- ▶ $*$ should be commutative, i.e. $x(t) * y(t) = y(t) * x(t)$.
- ▶ $*$ with an impulse at τ should yield the function $x(t - \tau)$.
- ▶ Associativity: $(x(t) * y(t)) * z(t) = x(t) * (y(t) * z(t))$.
- ▶ Distribution over addition:

$$x(t) * (y(t) + z(t)) = x(t) * y(t) + x(t) * z(t).$$


- ▶ Consistency when $\delta(t - \tau)$ is used in place of any signal(s).

We hope that there may exist such an integral operation, which boils down to a summation for digital signals.




Examples

Assume the desired properties: then




Examples

Assume the desired properties: then




Examples

Assume the desired properties: then




Examples

Assume the desired properties: then



Another Superposition 'Law'



Discrete Version

$$x(t) = x_0\delta(t) + x_1\delta(t - T) + \dots + x_m\delta(t - mT)$$

$$y(t) = \delta(t) + 3\delta(t - T) + 2\delta(t - 2T)$$



Discrete Version

$$x(t) = x_0\delta(t) + x_1\delta(t - T) + \cdots + x_m\delta(t - mT)$$

$$y(t) = \delta(t) + 3\delta(t - T) + 2\delta(t - 2T)$$

$$x(t) * y(t) = x(t) + 3x(t - T) + 2x(t - 2T).$$



Discrete Version

$$x(t) = x_0\delta(t) + x_1\delta(t - T) + \cdots + x_m\delta(t - mT)$$

$$y(t) = \delta(t) + 3\delta(t - T) + 2\delta(t - 2T)$$

$$x(t) * y(t) = x(t) + 3x(t - T) + 2x(t - 2T).$$

Eg. let $\bar{x} = (x_0, \dots, x_m) = (2, 3, 5, 4, 5)$ and $\bar{y} = (y_0, \dots, y_2) = (1, 3, 2)$.



Discrete Version

$$x(t) = x_0\delta(t) + x_1\delta(t - T) + \cdots + x_m\delta(t - mT)$$

$$y(t) = \delta(t) + 3\delta(t - T) + 2\delta(t - 2T)$$

$$x(t) * y(t) = x(t) + 3x(t - T) + 2x(t - 2T).$$

Eg. let $\bar{x} = (x_0, \dots, x_m) = (2, 3, 5, 4, 5)$ and $\bar{y} = (y_0, \dots, y_2) = (1, 3, 2)$.

$$\begin{array}{ccccccccc} & & 2 & & 3 & & 5 & & 4 & & 5 \\ & & & & & & & & & & \\ & & & & & & 1 & & 3 & & 2 \\ \hline \end{array}$$



Discrete Version

$$x(t) = x_0\delta(t) + x_1\delta(t - T) + \dots + x_m\delta(t - mT)$$

$$y(t) = \delta(t) + 3\delta(t - T) + 2\delta(t - 2T)$$

$$x(t) * y(t) = x(t) + 3x(t - T) + 2x(t - 2T).$$

Eg. let $\bar{x} = (x_0, \dots, x_m) = (2, 3, 5, 4, 5)$ and $\bar{y} = (y_0, \dots, y_2) = (1, 3, 2)$.

| | | 2 | 3 | 5 | 4 | 5 |
|-------------|---|---|----|----|----|----|
| | | | | 1 | 3 | 2 |
| 2 $x(t-2T)$ | | 4 | 6 | 10 | 8 | 10 |
| 3 $x(t-T)$ | 6 | 9 | 15 | 12 | 15 | |
| 1 $x(t-0T)$ | 2 | 3 | 5 | 4 | 5 | |



Discrete Version

$$x(t) = x_0\delta(t) + x_1\delta(t-T) + \dots + x_m\delta(t-mT)$$

$$y(t) = \delta(t) + 3\delta(t-T) + 2\delta(t-2T)$$

$$x(t) * y(t) = x(t) + 3x(t-T) + 2x(t-2T).$$

Eg. let $\bar{x} = (x_0, \dots, x_m) = (2, 3, 5, 4, 5)$ and $\bar{y} = (y_0, \dots, y_2) = (1, 3, 2)$.

| | | | | | | | |
|-------------|---|---|----|----|----|----|----|
| | | | 2 | 3 | 5 | 4 | 5 |
| | | | | | 1 | 3 | 2 |
| 2 $x(t-2T)$ | | | 4 | 6 | 10 | 8 | 10 |
| 3 $x(t-T)$ | | 6 | 9 | 15 | 12 | 15 | |
| 1 $x(t-0T)$ | 2 | 3 | 5 | 4 | 5 | | |
| | 2 | 9 | 18 | 25 | 27 | 23 | 10 |



Discrete Version

$$x(t) = x_0\delta(t) + x_1\delta(t-T) + \dots + x_m\delta(t-mT)$$

$$y(t) = \delta(t) + 3\delta(t-T) + 2\delta(t-2T)$$

$$x(t) * y(t) = x(t) + 3x(t-T) + 2x(t-2T).$$

Eg. let $\bar{x} = (x_0, \dots, x_m) = (2, 3, 5, 4, 5)$ and $\bar{y} = (y_0, \dots, y_2) = (1, 3, 2)$.

| | | | | | | | |
|-------------|---|---|----|----|----|----|----|
| | | | 2 | 3 | 5 | 4 | 5 |
| | | | | | 1 | 3 | 2 |
| 2 $x(t-2T)$ | | | 4 | 6 | 10 | 8 | 10 |
| 3 $x(t-T)$ | | 6 | 9 | 15 | 12 | 15 | |
| 1 $x(t-0T)$ | 2 | 3 | 5 | 4 | 5 | | |
| | 2 | 9 | 18 | 25 | 27 | 23 | 10 |

For $\bar{z} = \bar{x} * \bar{y}$, makes sense to define $z_n := \sum_k y_k x_{n-k}$



Primary School Days

x_1 x_2 x_3 x_4 x_5 x_6 x_7 x_8 x_9 x_{10} \otimes

1 1

x_1 x_2 x_3 x_4 x_5 x_6 x_7 x_8 x_9 x_{10} \oplus

x_1 x_2 x_3 x_4 x_5 x_6 x_7 x_8 x_9 x_{10}

y_1 y_2 y_3 y_4 y_5 y_6 y_7 y_8 y_9 y_{10} y_{11}

Multiplication without carry addition is called **convolution**.

$$\bar{x} * \bar{u} = (\bar{x} \text{ multiply } \bar{u}) \text{ modulo MAXNUM.}$$



Primary School Days

x_1 x_2 x_3 x_4 x_5 x_6 x_7 x_8 x_9 x_{10} \otimes

1 1

x_1 x_2 x_3 x_4 x_5 x_6 x_7 x_8 x_9 x_{10} \oplus

x_1 x_2 x_3 x_4 x_5 x_6 x_7 x_8 x_9 x_{10}

y_1 y_2 y_3 y_4 y_5 y_6 y_7 y_8 y_9 y_{10} y_{11}

Multiplication without carry addition is called **convolution**.

$$\bar{x} * \bar{u} = (\bar{x} \text{ multiply } \bar{u}) \text{ modulo MAXNUM.}$$



Digital Convolution

Notation: Vector $\bar{x} := x_0, \dots, x_{N-1}$.

Distribution of $*$ over $+$

$$\bar{x} * (\bar{u}_1 + \bar{u}_2) = (\bar{x} * \bar{u}_1) + (\bar{x} * \bar{u}_2).$$

Examples:

$$\bar{x} * [1 \ 1] = \bar{x} * ([1 \ 0] + [0 \ 1]) = (\bar{x} * [1 \ 0]) + (\bar{x} * [0 \ 1])$$

$$\bar{x} * [\alpha \ \beta] = \bar{x} * ([\alpha \ 0] + [0 \ \beta]) = (\bar{x} * [\alpha \ 0]) + (\bar{x} * [0 \ \beta])$$

Makes sense to have $\bar{x} * [1 \ 0] = \bar{x}$ — “digital impulse”.



Digital Convolution

Notation: Vector $\bar{x} := x_0, \dots, x_{N-1}$.

Distribution of $*$ over $+$

$$\bar{x} * (\bar{u}_1 + \bar{u}_2) = (\bar{x} * \bar{u}_1) + (\bar{x} * \bar{u}_2).$$

Examples:

$$\bar{x} * [1 \ 1] = \bar{x} * ([1 \ 0] + [0 \ 1]) = (\bar{x} * [1 \ 0]) + (\bar{x} * [0 \ 1])$$

$$\bar{x} * [\alpha \ \beta] = \bar{x} * ([\alpha \ 0] + [0 \ \beta]) = (\bar{x} * [\alpha \ 0]) + (\bar{x} * [0 \ \beta])$$

Makes sense to have $\bar{x} * [1 \ 0] = \bar{x}$ — “digital impulse”.



Digital Convolution

Notation: Vector $\bar{x} := x_0, \dots, x_{N-1}$.

Distribution of $*$ over $+$

$$\bar{x} * (\bar{u}_1 + \bar{u}_2) = (\bar{x} * \bar{u}_1) + (\bar{x} * \bar{u}_2).$$

Examples:

$$\bar{x} * [1 \ 1] = \bar{x} * ([1 \ 0] + [0 \ 1]) = (\bar{x} * [1 \ 0]) + (\bar{x} * [0 \ 1])$$

$$\bar{x} * [\alpha \ \beta] = \bar{x} * ([\alpha \ 0] + [0 \ \beta]) = (\bar{x} * [\alpha \ 0]) + (\bar{x} * [0 \ \beta])$$

Makes sense to have $\bar{x} * [1 \ 0] = \bar{x}$ — “digital impulse”.



Digital Convolution

Notation: Vector $\bar{x} := x_0, \dots, x_{N-1}$.

Distribution of $*$ over $+$

$$\bar{x} * (\bar{u}_1 + \bar{u}_2) = (\bar{x} * \bar{u}_1) + (\bar{x} * \bar{u}_2).$$

Examples:

$$\bar{x} * [1 \ 1] = \bar{x} * ([1 \ 0] + [0 \ 1]) = (\bar{x} * [1 \ 0]) + (\bar{x} * [0 \ 1])$$

$$\bar{x} * [\alpha \ \beta] = \bar{x} * ([\alpha \ 0] + [0 \ \beta]) = (\bar{x} * [\alpha \ 0]) + (\bar{x} * [0 \ \beta])$$

Makes sense to have $\bar{x} * [1 \ 0] = \bar{x}$ “— *digital impulse*”.



Integration

$$I = \int_2^7 |x - \sin(x)| dx$$




Figure: Riemann Sum for $f(x) = x - \sin(x)$



Integration

$$I = \int_2^7 |x - \sin(x)| dx$$




Figure: Riemann Sum for $f(x) = x - \sin(x)$



Integration

$$I = \int_2^7 |x - \sin(x)| dx$$




Figure: Riemann Sum for $f(x) = x - \sin(x)$



Riemann Approximation




Figure: Piecewise approximation of $f(x) = x - \sin(x)$



Riemann Approximation




Figure: Piecewise approximation of $f(x) = x - \sin(x)$




Figure: $f(x)$ as a Composition



Riemann Approximation




Figure: Piecewise approximation of $f(x) = x - \sin(x)$





Figure: $f(x)$ as a Composition



Riemann Approximation




Figure: Piecewise approximation of $f(x) = x - \sin(x)$




Figure: $f(x)$ as a Composition



Riemann Approximation




Figure: Piecewise approximation of $f(x) = x - \sin(x)$





Figure: $f(x)$ as a Composition




Generalizing Our Operator



Function $x(t)$


*




Function $y(t)$




Generalizing Our Operator



Generalizing Our Operator



Generalizing Our Operator




Function $x(t)$

Function $y(t)$

$$\bar{x}_T = \sum_{m \in I} x(mT) \delta(t - mT) \quad , \quad \bar{y}_T = \sum_{n \in J} y(nT) \delta(t - nT).$$



Generalizing Our Operator



Function $x(t)$


Function $y(t)$

$$\bar{x}_T = \sum_{m \in I} x(mT) \delta(t - mT) \quad , \quad \bar{y}_T = \sum_{n \in J} y(nT) \delta(t - nT).$$

$$x(t) * y(t) \approx r_T(t) * \bar{x}_T * \bar{y}_T * r_T(t)$$



Generalizing Our Operator




$$\bar{x}_T = \sum_{m \in I} x(mT) \delta(t - mT) \quad , \quad \bar{y}_T = \sum_{n \in J} y(nT) \delta(t - nT).$$

$$\begin{aligned} x(t) * y(t) &\approx r_T(t) * \bar{x}_T * \bar{y}_T * r_T(t) \\ &= \bar{x}_T * \bar{y}_T * r_T(t) * r_T(t). \end{aligned}$$



Generalizing Our Operator




$$\bar{x}_T = \sum_{m \in I} x(mT) \delta(t - mT) \quad , \quad \bar{y}_T = \sum_{n \in J} y(nT) \delta(t - nT).$$

$$\begin{aligned} x(t) * y(t) &\approx r_T(t) * \bar{x}_T * \bar{y}_T * r_T(t) \\ &= \boxed{\bar{x}_T * \bar{y}_T} * r_T(t) * r_T(t). \end{aligned}$$

Discrete Convolution



Generalizing Our Operator



Function $x(t)$

Function $y(t)$

$$\bar{x}_T = \sum_{m \in I} x(mT) \delta(t - mT) \quad , \quad \bar{y}_T = \sum_{n \in J} y(nT) \delta(t - nT).$$

$$\begin{aligned}
 x(t) * y(t) &\approx r_T(t) * \bar{x}_T * \bar{y}_T * r_T(t) \\
 &= \boxed{\bar{x}_T * \bar{y}_T} * \boxed{r_T(t) * r_T(t)}.
 \end{aligned}$$

Discrete Convolution

Analog Convolutn



Leaky Bucket



Courtesy: Open Clipart

20-08-2020, Class-3

