

MA 207 - Differential Equations-II


Amiya Kumar Pani

Department of Mathematics,
Indian Institute of Technology Bombay,
Powai, Mumbai 76
akp.ma207.2020@gmail.com

November 18, 2020


Start with Quotation and One Cartoon

*"If God has made Nature, Man made Mathematics
to study it "*
by Anonymous




Start with Quotation and One Cartoon

"If God has made Nature, Man made Mathematics to study it "
by Anonymous



INDIFFERENTIAL EQUATIONS



Outline of the lecture

- Singular Sturm Liouville Problems
- Example
- Exercises

Sturm-Liouville Boundary Value Problems

Consider the second order linear DE of the form:

$$(r(x)y')' + (q(x) + \lambda p(x))y = 0 \quad a < x < b, \quad (1)$$

$$k_1y(a) + k_2y'(a) = 0 \quad (2)$$

$$l_1y(b) + l_2y'(b) = 0 \quad (3)$$

Sturm-Liouville Boundary Value Problems

Consider the second order linear DE of the form:

$$(r(x)y')' + (q(x) + \lambda p(x))y = 0 \quad a < x < b, \quad (1)$$

$$k_1 y(a) + k_2 y'(a) = 0 \quad (2)$$

$$l_1 y(b) + l_2 y'(b) = 0 \quad (3)$$

Here k_1, k_2 not both are zero, l_1, l_2 not both are zeros, λ is a parameter; k_1, k_2, l_1, l_2 are given constants.

Sturm-Liouville Boundary Value Problems

Consider the second order linear DE of the form:

$$(r(x)y')' + (q(x) + \lambda p(x))y = 0 \quad a < x < b, \quad (1)$$

$$k_1 y(a) + k_2 y'(a) = 0 \quad (2)$$

$$l_1 y(b) + l_2 y'(b) = 0 \quad (3)$$

Here k_1, k_2 not both are zero, l_1, l_2 not both are zeros, λ is a parameter; k_1, k_2, l_1, l_2 are given constants. (1)-(3) is referred to as **STURM-LIOUVILLE BVP**.

Sturm-Liouville Boundary Value Problems

Consider the second order linear DE of the form:

$$(r(x)y')' + (q(x) + \lambda p(x))y = 0 \quad a < x < b, \quad (1)$$

$$k_1 y(a) + k_2 y'(a) = 0 \quad (2)$$

$$l_1 y(b) + l_2 y'(b) = 0 \quad (3)$$

Here k_1, k_2 not both are zero, l_1, l_2 not both are zeros, λ is a parameter; k_1, k_2, l_1, l_2 are given constants. (1)-(3) is referred to as **STURM-LIOUVILLE BVP**. Often (1) is represented as

$$\mathcal{L}(y) + \lambda p(x)y = 0, \quad \mathcal{L}(y) = (r(x)y')' + q(x)y.$$

$y \equiv 0$ is always a solution of (1)-(3).

Sturm-Liouville Boundary Value Problems

Consider the second order linear DE of the form:

$$(r(x)y')' + (q(x) + \lambda p(x))y = 0 \quad a < x < b, \quad (1)$$

$$k_1 y(a) + k_2 y'(a) = 0 \quad (2)$$

$$l_1 y(b) + l_2 y'(b) = 0 \quad (3)$$

Here k_1, k_2 not both are zero, l_1, l_2 not both are zeros, λ is a parameter; k_1, k_2, l_1, l_2 are given constants. (1)-(3) is referred to as **STURM-LIOUVILLE BVP**. Often (1) is represented as

$$\mathcal{L}(y) + \lambda p(x)y = 0, \quad \mathcal{L}(y) = (r(x)y')' + q(x)y.$$

$y \equiv 0$ is always a solution of (1)-(3).

We are interested to determine the values of the parameter λ for which the BVP has **non-trivial solutions**.

Sturm-Liouville BVP

Regular Sturm Liouville Problems:

Sturm-Liouville BVP

Regular Sturm Liouville Problems:

A Sturm-Liouville problem

$$(r(x)y')' + (q(x) + \lambda p(x))y = 0 \quad a \leq x \leq b,$$

$$k_1 y(a) + k_2 y'(a) = 0$$

$$l_1 y(b) + l_2 y'(b) = 0$$

Sturm-Liouville BVP

Regular Sturm Liouville Problems:

A Sturm-Liouville problem

$$(r(x)y')' + (q(x) + \lambda p(x))y = 0 \quad a \leq x \leq b,$$

$$k_1 y(a) + k_2 y'(a) = 0$$

$$l_1 y(b) + l_2 y'(b) = 0$$

is said to be **regular** if

Regular Sturm Liouville Problems:

A Sturm-Liouville problem

$$(r(x)y')' + (q(x) + \lambda p(x))y = 0 \quad a \leq x \leq b,$$

$$k_1 y(a) + k_2 y'(a) = 0$$

$$l_1 y(b) + l_2 y'(b) = 0$$

is said to be **regular** if

- 1 $r(x) > 0$, $p(x) > 0$ for $x \in [a, b]$;
- 2 $p(x)$, $q(x)$, $r(x)$, $r'(x)$ are continuous on $[a, b]$.
- 3 (a, b) is a finite interval.

Regular Sturm Liouville Problems:

A Sturm-Liouville problem

$$(r(x)y')' + (q(x) + \lambda p(x))y = 0 \quad a \leq x \leq b,$$

$$k_1 y(a) + k_2 y'(a) = 0$$

$$l_1 y(b) + l_2 y'(b) = 0$$

is said to be **regular** if

- 1 $r(x) > 0$, $p(x) > 0$ for $x \in [a, b]$;
- 2 $p(x)$, $q(x)$, $r(x)$, $r'(x)$ are continuous on $[a, b]$.
- 3 (a, b) is a finite interval.

A linear BVP is called **SINGULAR**, if it is not REGULAR.

Regular Sturm Liouville Problems:

A Sturm-Liouville problem

$$(r(x)y')' + (q(x) + \lambda p(x))y = 0 \quad a \leq x \leq b,$$

$$k_1 y(a) + k_2 y'(a) = 0$$

$$l_1 y(b) + l_2 y'(b) = 0$$

is said to be **regular** if

- 1 $r(x) > 0$, $p(x) > 0$ for $x \in [a, b]$;
- 2 $p(x)$, $q(x)$, $r(x)$, $r'(x)$ are continuous on $[a, b]$.
- 3 (a, b) is a finite interval.

A linear BVP is called **SINGULAR**, if it is not REGULAR.

(Example : either $a = -\infty$, or $b = \infty$ or both; $r(x) = 0$ for at least one $x \in [a, b]$.)

Regular Sturm Liouville Problems:

A Sturm-Liouville problem

$$(r(x)y')' + (q(x) + \lambda p(x))y = 0 \quad a \leq x \leq b,$$

$$k_1 y(a) + k_2 y'(a) = 0$$

$$l_1 y(b) + l_2 y'(b) = 0$$

is said to be **regular** if

- 1 $r(x) > 0$, $p(x) > 0$ for $x \in [a, b]$;
- 2 $p(x)$, $q(x)$, $r(x)$, $r'(x)$ are continuous on $[a, b]$.
- 3 (a, b) is a finite interval.

A linear BVP is called **SINGULAR**, if it is not REGULAR.

(Example : either $a = -\infty$, or $b = \infty$ or both; $r(x) = 0$ for at least one $x \in [a, b]$.)

Questions related to Singular SL-BVP

Question

1. What type of boundary conditions can be allowed in a singular Sturm-Liouville Problem?

Questions related to Singular SL-BVP

Question

1. *What type of boundary conditions can be allowed in a singular Sturm-Liouville Problem?*

Question

2. *To what extent the properties of eigenvalues and eigenfunctions of regular Sturm-Liouville problem can be extended to eigenvalues and eigenfunctions of singular Sturm-Liouville problem. More precisely, are the eigenvalues real, are eigenfunctions orthogonal, can a given function be expanded as series of eigenfunctions?*

Questions related to Singular SL-BVP

Question

1. *What type of boundary conditions can be allowed in a singular Sturm-Liouville Problem?*

Question

2. *To what extent the properties of eigenvalues and eigenfunctions of regular Sturm-Liouville problem can be extended to eigenvalues and eigenfunctions of singular Sturm-Liouville problem. More precisely, are the eigenvalues real, are eigenfunctions orthogonal, can a given function be expanded as series of eigenfunctions?*

Both of these questions can be answered by studying the self-adjoint form.

Self-Adjoint Form

Let V be the space of twice continuously differentiable functions on $[a, b]$ with inner-product $(v, w) = \int_a^b v(x)w(x) dx$.

Self-Adjoint Form

Let V be the space of twice continuously differentiable functions on $[a, b]$ with inner-product $(v, w) = \int_a^b v(x)w(x) dx$.

Hence, we obtain the **LAGRANGE'S IDENTITY** :

Self-Adjoint Form

Let V be the space of twice continuously differentiable functions on $[a, b]$ with inner-product $(v, w) = \int_a^b v(x)w(x) dx$.

Hence, we obtain the **LAGRANGE'S IDENTITY** :

$$(\mathcal{L}u, v) - (u, \mathcal{L}v) = r(x)(u'v - uv') \Big|_{x=a}^{x=b}.$$

Definition

\mathcal{L} is called *self-adjoint*, if $(\mathcal{L}u, v) = (u, \mathcal{L}v)$.

In order to make \mathcal{L} self-adjoint, we could define V as the space of twice continuously differentiable functions satisfying the boundary conditions. That is,

Self-Adjoint Form

Let V be the space of twice continuously differentiable functions on $[a, b]$ with inner-product $(v, w) = \int_a^b v(x)w(x) dx$.

Hence, we obtain the **LAGRANGE'S IDENTITY** :

$$(\mathcal{L}u, v) - (u, \mathcal{L}v) = r(x)(u'v - uv') \Big|_{x=a}^{x=b}.$$

Definition

\mathcal{L} is called *self-adjoint*, if $(\mathcal{L}u, v) = (u, \mathcal{L}v)$.

In order to make \mathcal{L} self-adjoint, we could define V as the space of twice continuously differentiable functions satisfying the boundary conditions. That is, $u, v \in V$ satisfies

Self-Adjoint Form

Let V be the space of twice continuously differentiable functions on $[a, b]$ with inner-product $(v, w) = \int_a^b v(x)w(x) dx$.

Hence, we obtain the **LAGRANGE'S IDENTITY** :

$$(\mathcal{L}u, v) - (u, \mathcal{L}v) = r(x)(u'v - uv') \Big|_{x=a}^{x=b}.$$

Definition

\mathcal{L} is called *self-adjoint*, if $(\mathcal{L}u, v) = (u, \mathcal{L}v)$.

In order to make \mathcal{L} self-adjoint, we could define V as the space of twice continuously differentiable functions satisfying the boundary conditions. That is, $u, v \in V$ satisfies $k_1u(a) + k_2u'(a) = 0$, $l_1u(b) + l_2u'(b) = 0$,

Self-Adjoint Form

Let V be the space of twice continuously differentiable functions on $[a, b]$ with inner-product $(v, w) = \int_a^b v(x)w(x) dx$.

Hence, we obtain the **LAGRANGE'S IDENTITY** :

$$(\mathcal{L}u, v) - (u, \mathcal{L}v) = r(x)(u'v - uv') \Big|_{x=a}^{x=b}.$$

Definition

\mathcal{L} is called *self-adjoint*, if $(\mathcal{L}u, v) = (u, \mathcal{L}v)$.

In order to make \mathcal{L} self-adjoint, we could define V as the space of twice continuously differentiable functions satisfying the boundary conditions. That is, $u, v \in V$ satisfies

$$\begin{aligned} k_1 u(a) + k_2 u'(a) &= 0, & l_1 u(b) + l_2 u'(b) &= 0, \\ k_1 v(a) + k_2 v'(a) &= 0, & l_1 v(b) + l_2 v'(b) &= 0 \end{aligned}$$

Self-Adjoint Form——

Note that

$$\begin{aligned}(\mathcal{L}u, v) - (u, \mathcal{L}v) &= r(x)(u'v - uv') \Big|_{x=a}^{x=b} \\&= r(b)(u'(b)v(b) - v'(b)u(b)) \\&\quad - r(a)(u'(a)v(a) - v'(a)u(a))\end{aligned}$$

Self-Adjoint Form—

Note that

$$\begin{aligned}(\mathcal{L}u, v) - (u, \mathcal{L}v) &= r(x)(u'v - uv') \Big|_{x=a}^{x=b} \\&= r(b)(u'(b)v(b) - v'(b)u(b)) \\&\quad - r(a)(u'(a)v(a) - v'(a)u(a))\end{aligned}$$

If $l_2 \neq 0$ and $k_2 \neq 0$, then from the boundary condition:

$u'(b) = -(l_1/l_2)u(b)$, $v'(b) = -(l_1/l_2)v(b)$ and

$u'(a) = -(k_1/k_2)u(a)$, $v'(a) = -(k_1/k_2)v(a)$. On substitution

$$\begin{aligned}(\mathcal{L}u, v) - (u, \mathcal{L}v) &= r(b) \left(-\cancel{\frac{l_1}{l_2} u(b)v(b)} + \cancel{\frac{l_1}{l_2} v(b)u(b)} \right) \\&\quad - r(a) \left(-\cancel{\frac{k_1}{k_2} u(a)v(a)} + \cancel{\frac{k_1}{k_2} v(a)u(a)} \right) = 0\end{aligned}$$

Self-Adjoint Form—

Note that

$$\begin{aligned}(\mathcal{L}u, v) - (u, \mathcal{L}v) &= r(x)(u'v - uv') \Big|_{x=a}^{x=b} \\&= r(b)(u'(b)v(b) - v'(b)u(b)) \\&\quad - r(a)(u'(a)v(a) - v'(a)u(a))\end{aligned}$$

If $l_2 \neq 0$ and $k_2 \neq 0$, then from the boundary condition:

$u'(b) = -(l_1/l_2)u(b)$, $v'(b) = -(l_1/l_2)v(b)$ and

$u'(a) = -(k_1/k_2)u(a)$, $v'(a) = -(k_1/k_2)v(a)$. On substitution

$$\begin{aligned}(\mathcal{L}u, v) - (u, \mathcal{L}v) &= r(b) \left(-\cancel{\frac{l_1}{l_2} u(b)v(b)} + \cancel{\frac{l_1}{l_2} v(b)u(b)} \right) \\&\quad - r(a) \left(-\cancel{\frac{k_1}{k_2} u(a)v(a)} + \cancel{\frac{k_1}{k_2} v(a)u(a)} \right) = 0\end{aligned}$$

Hence, $(\mathcal{L}u, v) = (u, \mathcal{L}v)$.

How to write a DE in self-adjoint form?

QUESTION : Given a second linear DE,

$$a(x)y'' + b(x)y' + c(x)y = 0,$$

How to write a DE in self-adjoint form?

QUESTION : Given a second linear DE,

$$a(x)y'' + b(x)y' + c(x)y = 0, \quad (4)$$

can we write it in the self- adjoint form?

How to write a DE in self-adjoint form?

QUESTION : Given a second linear DE,

$$a(x)y'' + b(x)y' + c(x)y = 0, \quad (4)$$

can we write it in the self-adjoint form?

$$\underbrace{a(x)y'' + b(x)y'}_{(r(x)y')'} + c(x)y = 0.$$

How to write a DE in self-adjoint form?

QUESTION : Given a second linear DE,

$$a(x)y'' + b(x)y' + c(x)y = 0, \quad (4)$$

can we write it in the self-adjoint form?

$$\underbrace{a(x)y'' + b(x)y'}_{(r(x)y')'} + c(x)y = 0.$$

That is, we seek an integrating factor $\mu(x)$ such that (4) can be represented as :

How to write a DE in self-adjoint form?

QUESTION : Given a second linear DE,

$$a(x)y'' + b(x)y' + c(x)y = 0, \quad (4)$$

can we write it in the self- adjoint form?

$$\underbrace{a(x)y'' + b(x)y'}_{(r(x)y')'} + c(x)y = 0.$$

That is, we seek an integrating factor $\mu(x)$ such that (4) can be represented as :

$$(\mu(x)a(x)y')' + \mu(x)c(x)y = 0.$$

How to write a DE in self-adjoint form?

QUESTION : Given a second linear DE,

$$a(x)y'' + b(x)y' + c(x)y = 0, \quad (4)$$

can we write it in the self- adjoint form?

$$\underbrace{a(x)y'' + b(x)y'}_{(r(x)y')'} + c(x)y = 0.$$

That is, we seek an integrating factor $\mu(x)$ such that (4) can be represented as :

$$(\mu(x)a(x)y')' + \mu(x)c(x)y = 0. \quad (5)$$

From (4) (multiplied by $\mu(x)$) and (5), equate the coefficients of y' to obtain

$$\mu(x)a'(x) + \mu'(x)a(x) = \mu(x)b(x)$$

How to write a DE in self-adjoint form?

QUESTION : Given a second linear DE,

$$a(x)y'' + b(x)y' + c(x)y = 0, \quad (4)$$

can we write it in the self- adjoint form?

$$\underbrace{a(x)y'' + b(x)y'}_{(r(x)y')'} + c(x)y = 0.$$

That is, we seek an integrating factor $\mu(x)$ such that (4) can be represented as :

$$(\mu(x)a(x)y')' + \mu(x)c(x)y = 0. \quad (5)$$

From (4) (multiplied by $\mu(x)$) and (5), equate the coefficients of y' to obtain

$$\begin{aligned} \mu(x)a'(x) + \mu'(x)a(x) &= \mu(x)b(x) \\ \Rightarrow \mu'(x)a(x) &= (b(x) - a'(x))\mu(x) \end{aligned}$$

How to write a DE in self-adjoint form?

QUESTION : Given a second linear DE,

$$a(x)y'' + b(x)y' + c(x)y = 0, \quad (4)$$

can we write it in the self- adjoint form?

$$\underbrace{a(x)y'' + b(x)y'}_{(r(x)y')'} + c(x)y = 0.$$

That is, we seek an integrating factor $\mu(x)$ such that (4) can be represented as :

$$(\mu(x)a(x)y')' + \mu(x)c(x)y = 0. \quad (5)$$

From (4) (multiplied by $\mu(x)$) and (5), equate the coefficients of y' to obtain

$$\begin{aligned} \mu(x)a'(x) + \mu'(x)a(x) &= \mu(x)b(x) \\ \Rightarrow \mu'(x)a(x) &= (b(x) - a'(x))\mu(x) \end{aligned}$$

$$\mu'(x) = \frac{(b(x) - a'(x))}{a(x)} \mu(x) \quad (\text{assuming } a(x) \neq 0)$$

$$\mu'(x) = \frac{(b(x) - a'(x))}{a(x)} \mu(x) \quad (\text{assuming } a(x) \neq 0)$$

On solving for $\mu(x)$, we obtain

$$\mu'(x) = \frac{(b(x) - a'(x))}{a(x)} \mu(x) \quad (\text{assuming } a(x) \neq 0)$$

On solving for $\mu(x)$, we obtain

$$\mu(x) = \frac{1}{a(x)} e^{\int \frac{b(x)}{a(x)} dx} \quad (a(x) \neq 0).$$

$$\mu'(x) = \frac{(b(x) - a'(x))}{a(x)} \mu(x) \quad (\text{assuming } a(x) \neq 0)$$

On solving for $\mu(x)$, we obtain

$$\mu(x) = \frac{1}{a(x)} e^{\int \frac{b(x)}{a(x)} dx} \quad (a(x) \neq 0).$$

Example 1 : Legendre equation

$(1 - x^2)y'' - 2xy' + p(p+1)y = 0$, $x \in (-1, 1)$ can be put in the self-adjoint form as

$$\mu'(x) = \frac{(b(x) - a'(x))}{a(x)} \mu(x) \quad (\text{assuming } a(x) \neq 0)$$

On solving for $\mu(x)$, we obtain

$$\mu(x) = \frac{1}{a(x)} e^{\int \frac{b(x)}{a(x)} dx} \quad (a(x) \neq 0).$$

Example 1 : Legendre equation

$(1 - x^2)y'' - 2xy' + p(p+1)y = 0$, $x \in (-1, 1)$ can be put in the self-adjoint form as

$$\mathcal{L}y + \lambda y := ((1 - x^2)y')' + p(p+1)y = 0$$

On the set of all two time continuously differentiable functions v on $(-1, 1)$ with v, v' are bounded as $x \rightarrow \pm 1$, \mathcal{L} is Self-adjoint as

$$(\mathcal{L}u, v) - (u, \mathcal{L}v) = (1 - x^2)(u'v - uv') \Big|_{x=-1}^{x=1} = 0$$

$$(r(x) = 1 - x^2).$$

Example 2: Chebyshev's equation

$(1 - x^2)y'' - xy' + \lambda y = 0$, $x \in (-1, 1)$ can be put in the self-adjoint form

Example 2: Chebyshev's equation

$(1 - x^2)y'' - xy' + \lambda y = 0$, $x \in (-1, 1)$ can be put in the self-adjoint form

$$(\mu(x)a(x)y')' + \mu(x)c(x)y = 0, \quad (6)$$

by choosing $\mu(x) = \frac{1}{a(x)} e^{\int \frac{b(x)}{a(x)} dx}$.

Example 2: Chebyshev's equation

$(1 - x^2)y'' - xy' + \lambda y = 0$, $x \in (-1, 1)$ can be put in the self-adjoint form

$$(\mu(x)a(x)y')' + \mu(x)c(x)y = 0, \quad (6)$$

by choosing $\mu(x) = \frac{1}{a(x)} e^{\int \frac{b(x)}{a(x)} dx}$.

On the space of twice continuously differential functions on $(-1, 1)$ with function and its first derivative as bounded as $x \rightarrow \pm 1$

Example 2: Chebyshev's equation

$(1 - x^2)y'' - xy' + \lambda y = 0$, $x \in (-1, 1)$ can be put in the self-adjoint form

$$(\mu(x)a(x)y')' + \mu(x)c(x)y = 0, \quad (6)$$

by choosing $\mu(x) = \frac{1}{a(x)} e^{\int \frac{b(x)}{a(x)} dx}$.

On the space of twice continuously differential functions on $(-1, 1)$ with function and its first derivative as bounded as $x \rightarrow \pm 1$

$$\mu(x) = \frac{1}{1 - x^2} e^{-\int \frac{x}{1-x^2} dx}$$

Example 2: Chebyshev's equation

$(1 - x^2)y'' - xy' + \lambda y = 0$, $x \in (-1, 1)$ can be put in the self-adjoint form

$$(\mu(x)a(x)y')' + \mu(x)c(x)y = 0, \quad (6)$$

by choosing $\mu(x) = \frac{1}{a(x)} e^{\int \frac{b(x)}{a(x)} dx}$.

On the space of twice continuously differential functions on $(-1, 1)$ with function and its first derivative as bounded as $x \rightarrow \pm 1$

$$\mu(x) = \frac{1}{1 - x^2} e^{-\int \frac{x}{1-x^2} dx} \quad (\text{put } 1 - x^2 = t, \quad -2x dx = dt)$$

Example 2: Chebyshev's equation

$(1 - x^2)y'' - xy' + \lambda y = 0$, $x \in (-1, 1)$ can be put in the self-adjoint form

$$(\mu(x)a(x)y')' + \mu(x)c(x)y = 0, \quad (6)$$

by choosing $\mu(x) = \frac{1}{a(x)} e^{\int \frac{b(x)}{a(x)} dx}$.

On the space of twice continuously differential functions on $(-1, 1)$ with function and its first derivative as bounded as $x \rightarrow \pm 1$

$$\begin{aligned} \mu(x) &= \frac{1}{1 - x^2} e^{-\int \frac{x}{1-x^2} dx} \quad (\text{put } 1 - x^2 = t, \quad -2x dx = dt) \\ &= \frac{1}{1 - x^2} \times \sqrt{(1 - x^2)} = \frac{1}{\sqrt{1 - x^2}}. \end{aligned}$$

Example 2: Chebyshev's equation

$(1 - x^2)y'' - xy' + \lambda y = 0$, $x \in (-1, 1)$ can be put in the self-adjoint form

$$(\mu(x)a(x)y')' + \mu(x)c(x)y = 0, \quad (6)$$

by choosing $\mu(x) = \frac{1}{a(x)} e^{\int \frac{b(x)}{a(x)} dx}$.

On the space of twice continuously differential functions on $(-1, 1)$ with function and its first derivative as bounded as $x \rightarrow \pm 1$

$$\begin{aligned} \mu(x) &= \frac{1}{1 - x^2} e^{-\int \frac{x}{1-x^2} dx} \quad (\text{put } 1 - x^2 = t, \quad -2x dx = dt) \\ &= \frac{1}{1 - x^2} \times \sqrt{(1 - x^2)} = \frac{1}{\sqrt{1 - x^2}}. \end{aligned}$$

Hence, the self-adjoint form for Chebyshev's equation is

$$(\sqrt{1 - x^2} y')' + \frac{\lambda}{\sqrt{(1 - x^2)}} y = 0.$$

Example 2: Chebyshev's equation

$(1 - x^2)y'' - xy' + \lambda y = 0$, $x \in (-1, 1)$ can be put in the self-adjoint form

$$(\mu(x)a(x)y')' + \mu(x)c(x)y = 0, \quad (6)$$

by choosing $\mu(x) = \frac{1}{a(x)} e^{\int \frac{b(x)}{a(x)} dx}$.

On the space of twice continuously differential functions on $(-1, 1)$ with function and its first derivative as bounded as $x \rightarrow \pm 1$

$$\begin{aligned} \mu(x) &= \frac{1}{1 - x^2} e^{-\int \frac{x}{1-x^2} dx} \quad (\text{put } 1 - x^2 = t, \quad -2x dx = dt) \\ &= \frac{1}{1 - x^2} \times \sqrt{(1 - x^2)} = \frac{1}{\sqrt{1 - x^2}}. \end{aligned}$$

Hence, the self-adjoint form for Chebyshev's equation is

$$(\sqrt{1 - x^2} y')' + \frac{\lambda}{\sqrt{(1 - x^2)}} y = 0.$$

(Check: \mathcal{L} is self-adjoint?).

Example 3 : Hermite Equation

Let us try to write the Hermite equation

$y'' - xy' + \lambda y = 0 \quad -\infty < x < \infty$ in the self-adjoint form.

Example 3 : Hermite Equation

Let us try to write the Hermite equation

$y'' - xy' + \lambda y = 0 \quad -\infty < x < \infty$ in the self-adjoint form.

$$\mu(x) = e^{\int (-x) dx} = e^{-x^2/2}.$$

Example 3 : Hermite Equation

Let us try to write the Hermite equation

$y'' - xy' + \lambda y = 0 \quad -\infty < x < \infty$ in the self-adjoint form.

$$\mu(x) = e^{\int (-x) dx} = e^{-x^2/2}.$$

That is, the equation can be written as

Example 3 : Hermite Equation

Let us try to write the Hermite equation

$y'' - xy' + \lambda y = 0 \quad -\infty < x < \infty$ in the self-adjoint form.

$$\mu(x) = e^{\int (-x) dx} = e^{-x^2/2}.$$

That is, the equation can be written as

$$(e^{-x^2/2} y')' + \lambda e^{-x^2/2} y = 0, \quad -\infty < x < \infty.$$

Example 3 : Hermite Equation

Let us try to write the Hermite equation

$y'' - xy' + \lambda y = 0 \quad -\infty < x < \infty$ in the self-adjoint form.

$$\mu(x) = e^{\int (-x) dx} = e^{-x^2/2}.$$

That is, the equation can be written as

$$(e^{-x^2/2} y')' + \lambda e^{-x^2/2} y = 0, \quad -\infty < x < \infty.$$

To check self-adjointness of \mathcal{L} , where $\mathcal{L}u = (e^{-x^2/2} u')'$, we need to impose appropriate boundary conditions.

Example 3 : Hermite Equation

Let us try to write the Hermite equation

$y'' - xy' + \lambda y = 0 \quad -\infty < x < \infty$ in the self-adjoint form.

$$\mu(x) = e^{\int (-x) dx} = e^{-x^2/2}.$$

That is, the equation can be written as

$$(e^{-x^2/2} y')' + \lambda e^{-x^2/2} y = 0, \quad -\infty < x < \infty.$$

To check self-adjointness of \mathcal{L} , where $\mathcal{L}u = (e^{-x^2/2} u')'$, we need to impose appropriate boundary conditions.

$$(\mathcal{L}u, v) = \int_{-\infty}^{\infty} (e^{-x^2/2} u')' v dx$$

Example 3 : Hermite Equation

Let us try to write the Hermite equation

$y'' - xy' + \lambda y = 0 \quad -\infty < x < \infty$ in the self-adjoint form.

$$\mu(x) = e^{\int (-x) dx} = e^{-x^2/2}.$$

That is, the equation can be written as

$$(e^{-x^2/2} y')' + \lambda e^{-x^2/2} y = 0, \quad -\infty < x < \infty.$$

To check self-adjointness of \mathcal{L} , where $\mathcal{L}u = (e^{-x^2/2} u')'$, we need to impose appropriate boundary conditions.

$$\begin{aligned} (\mathcal{L}u, v) &= \int_{-\infty}^{\infty} (e^{-x^2/2} u')' v \, dx \\ &= \lim_{M \rightarrow \infty} \int_{-M}^M (e^{-x^2/2} u')' v \, dx \end{aligned}$$

Example 3 : Hermite Equation

Let us try to write the Hermite equation

$y'' - xy' + \lambda y = 0 \quad -\infty < x < \infty$ in the self-adjoint form.

$$\mu(x) = e^{\int (-x) dx} = e^{-x^2/2}.$$

That is, the equation can be written as

$$(e^{-x^2/2} y')' + \lambda e^{-x^2/2} y = 0, \quad -\infty < x < \infty.$$

To check self-adjointness of \mathcal{L} , where $\mathcal{L}u = (e^{-x^2/2} u')'$, we need to impose appropriate boundary conditions.

$$\begin{aligned} (\mathcal{L}u, v) &= \int_{-\infty}^{\infty} (e^{-x^2/2} u')' v \, dx \\ &= \lim_{M \rightarrow \infty} \int_{-M}^M (e^{-x^2/2} u')' v \, dx \\ &= \lim_{M \rightarrow \infty} \left\{ (e^{-x^2/2} u') v \Big|_{-M}^M - \int_{-M}^M (e^{-x^2/2} u') v \, dx \right\} \end{aligned}$$

Example 3 : Hermite Equation

Let us try to write the Hermite equation

$y'' - xy' + \lambda y = 0 \quad -\infty < x < \infty$ in the self-adjoint form.

$$\mu(x) = e^{\int (-x) dx} = e^{-x^2/2}.$$

That is, the equation can be written as

$$(e^{-x^2/2} y')' + \lambda e^{-x^2/2} y = 0, \quad -\infty < x < \infty.$$

To check self-adjointness of \mathcal{L} , where $\mathcal{L}u = (e^{-x^2/2} u')'$, we need to impose appropriate boundary conditions.

$$\begin{aligned} (\mathcal{L}u, v) &= \int_{-\infty}^{\infty} (e^{-x^2/2} u')' v \, dx \\ &= \lim_{M \rightarrow \infty} \int_{-M}^M (e^{-x^2/2} u')' v \, dx \\ &= \lim_{M \rightarrow \infty} \left\{ (e^{-x^2/2} u') v \Big|_{-M}^M - \int_{-M}^M (e^{-x^2/2} u') v \, dx \right\} \end{aligned}$$

$$(\mathcal{L}u, v) - (u, \mathcal{L}v) = \lim_{M \rightarrow \infty} \left(e^{-x^2/2} (u'v - uv') \right) \Big|_{-M}^M$$

$$\begin{aligned}
 (\mathcal{L}u, v) - (u, \mathcal{L}v) &= \lim_{M \rightarrow \infty} \left(e^{-x^2/2} (u'v - uv') \right) \Big|_{-M}^M \\
 &= \lim_{M \rightarrow \infty} e^{-x^2/2} (u'v - uv') \Big|_{-M}^M.
 \end{aligned}$$

$$\begin{aligned}
 (\mathcal{L}u, v) - (u, \mathcal{L}v) &= \lim_{M \rightarrow \infty} \left(e^{-x^2/2} (u'v - uv') \right) \Big|_{-M}^M \\
 &= \lim_{M \rightarrow \infty} e^{-x^2/2} (u'v - uv') \Big|_{-M}^M.
 \end{aligned}$$

To make \mathcal{L} self-adjoint, impose BC's on u and v as

$$\lim_{x \rightarrow \pm\infty} e^{-x^2/2} u(x) = 0, \quad \lim_{x \rightarrow \pm\infty} e^{-x^2/2} v(x) = 0.$$

$$\begin{aligned}
 (\mathcal{L}u, v) - (u, \mathcal{L}v) &= \lim_{M \rightarrow \infty} \left(e^{-x^2/2} (u'v - uv') \right) \bigg|_{-M}^M \\
 &= \lim_{M \rightarrow \infty} e^{-x^2/2} (u'v - uv') \bigg|_{-M}^M.
 \end{aligned}$$

To make \mathcal{L} self-adjoint, impose BC's on u and v as

$$\lim_{x \rightarrow \pm\infty} e^{-x^2/2} u(x) = 0, \quad \lim_{x \rightarrow \pm\infty} e^{-x^2/2} v(x) = 0.$$

Example 4 : Express the Laguerre equation

$xy'' + (1-x)y' + \lambda y = 0$, $0 < x < \infty$ in the self-adjoint form. (try it!)

Eigenvalues and eigenfunctions of Singular SL Problem

- Since

\mathcal{L} is self adjoint in $\mathcal{L}y + \lambda p(x)y = 0$, $\mathcal{L}(y) = (r(x)y')' + q(x)y$,

Eigenvalues and eigenfunctions of Singular SL Problem

- Since

\mathcal{L} is self adjoint in $\mathcal{L}y + \lambda p(x)y = 0$, $\mathcal{L}(y) = (r(x)y')' + q(x)y$,
all eigenvalues (if exist) are real with real eigenfunctions.

Eigenvalues and eigenfunctions of Singular SL Problem

- Since

\mathcal{L} is self adjoint in $\mathcal{L}y + \lambda p(x)y = 0$, $\mathcal{L}(y) = (r(x)y')' + q(x)y$,
all eigenvalues (if exist) are real with real eigenfunctions.

Therefore, all the four examples given in previous slides have eigenvalues with corresponding eigenfunctions are real and proofs are exactly on the similar lines with the proofs of regular Sturm-Liouville problem. (pl. check this!)

Eigenvalues and eigenfunctions of Singular SL Problem

- Since

\mathcal{L} is self adjoint in $\mathcal{L}y + \lambda p(x)y = 0$, $\mathcal{L}(y) = (r(x)y')' + q(x)y$,
all eigenvalues (if exist) are real with real eigenfunctions.

Therefore, all the four examples given in previous slides have eigenvalues with corresponding eigenfunctions are real and proofs are exactly on the similar lines with the proofs of regular Sturm-Liouville problem. (pl. check this!)

- Most striking difference between regular and singular SL problems: In a singular SL problem, the eigenvalues may not be discrete, that is, the problem may have nontrivial solution corresponding to every value of λ in some interval. In that case, we call that the problem has continuous spectrum.

Eigenvalues and eigenfunctions of Singular SL Problem

- Since

\mathcal{L} is self adjoint in $\mathcal{L}y + \lambda p(x)y = 0$, $\mathcal{L}(y) = (r(x)y')' + q(x)y$,
all eigenvalues (if exist) are real with real eigenfunctions.

Therefore, all the four examples given in previous slides have eigenvalues with corresponding eigenfunctions are real and proofs are exactly on the similar lines with the proofs of regular Sturm-Liouville problem. (pl. check this!)

- Most striking difference between regular and singular SL problems: In a singular SL problem, the eigenvalues may not be discrete, that is, the problem may have nontrivial solution corresponding to every value of λ in some interval. In that case, we call that the problem has continuous spectrum.

However, all four examples discussed in earlier slides have countable number of eigenvalues and eigenfunctions.

Orthogonality of Eigenfunctions

For all four examples (Legendre , Chebyshev's, Hermite and Laguerre equations), if $\{\phi_n\}_{n=0}^{\infty}$ be the set of eigenfunctions corresponding to eigenvalues $\{\lambda_n\}_{n=0}^{\infty}$, then eigenfunctions corresponding to distinct eigenvalues are orthogonal with respect to the weight function $p(x)$ (appearing in their respective self-adjoint form).

Orthogonality of Eigenfunctions

For all four examples (Legendre , Chebyshev's, Hermite and Laguerre equations), if $\{\phi_n\}_{n=0}^{\infty}$ be the set of eigenfunctions corresponding to eigenvalues $\{\lambda_n\}_{n=0}^{\infty}$, then eigenfunctions corresponding to distinct eigenvalues are orthogonal with respect to the weight function $p(x)$ (appearing in their respective self-adjoint form).

Example 1 : Legendre equation

$$(1 - x^2)y'' - 2xy' + n(n+1)y = 0, \quad x \in (-1, 1)$$

Orthogonality of Eigenfunctions

For all four examples (Legendre , Chebyshev's, Hermite and Laguerre equations), if $\{\phi_n\}_{n=0}^{\infty}$ be the set of eigenfunctions corresponding to eigenvalues $\{\lambda_n\}_{n=0}^{\infty}$, then eigenfunctions corresponding to distinct eigenvalues are orthogonal with respect to the weight function $p(x)$ (appearing in their respective self-adjoint form).

Example 1 : Legendre equation

$(1 - x^2)y'' - 2xy' + n(n+1)y = 0$, $x \in (-1, 1)$ in its self adjoint form: $((1 - x^2)y')' - 2xy' + \lambda y = 0$, $x \in (-1, 1)$.

Eigenvalues: $\lambda_n = n(n+1)$ with corresponding eigenfunction $P_n(x)$ for $n = 0, 1, \dots$.

Here, $p(x) = 1$ and orthogonality property:

$$\int_{-1}^1 P_n(x) P_m(x) dx = 0, \quad (m \neq n).$$

Note in addition, if we assume $y(0) = 0$, then eigenvalues $\lambda_n = n(n+1)$, when n is odd nonnegative integers.

Example 2: Chebyshev's equation

In its standard form: $(1 - x^2)y'' - xy' + \lambda y = 0$, $x \in (-1, 1)$,
with y, y' are bounded as $x \rightarrow \pm 1$.

Example 2: Chebyshev's equation

In its standard form: $(1 - x^2)y'' - xy' + \lambda y = 0$, $x \in (-1, 1)$,
with y, y' are bounded as $x \rightarrow \pm 1$. In its self-adjoint form :

$$(\sqrt{1 - x^2}y')' + \frac{\lambda}{\sqrt{(1 - x^2)}}y = 0.$$

Example 2: Chebyshev's equation

In its standard form: $(1 - x^2)y'' - xy' + \lambda y = 0$, $x \in (-1, 1)$,
with y, y' are bounded as $x \rightarrow \pm 1$. In its self-adjoint form :

$$(\sqrt{1 - x^2}y')' + \frac{\lambda}{\sqrt{(1 - x^2)}}y = 0.$$

Eigenvalues: $\{\lambda_n = n^2\}_{n=0}^{\infty}$ and the corresponding eigenfunctions
are the Chebyshev polynomials $\{T_n(x)\}_{n=0}^{\infty}$ with

$$T_0(x) = 1, T_1(x) = x, T_2(x) = 1 - 2x^2, \dots$$

Here, $p(x) = 1/\sqrt{1 - x^2}$ and orthogonality property:

$$\int_{-1}^1 (1 - x^2)^{-1/2} T_m(x) T_n(x) dx = 0, \quad (m \neq n).$$

Example 3: Hermite equation

In its standard form:

$$y'' - xy' + \lambda y = 0 \quad -\infty < x < \infty$$

with y and y' polynomially bounded at ∞ .

Example 3: Hermite equation

In its standard form:

$$y'' - xy' + \lambda y = 0 \quad -\infty < x < \infty$$

with y and y' polynomially bounded at ∞ .

In the self-adjoint form:

Example 3: Hermite equation

In its standard form:

$$y'' - xy' + \lambda y = 0 \quad -\infty < x < \infty$$

with y and y' polynomially bounded at ∞ .

In the self-adjoint form:

$$(e^{-x^2/2}y')' + \lambda e^{-x^2/2}y = 0, \quad -\infty < x < \infty.$$

Example 3: Hermite equation

In its standard form:

$$y'' - xy' + \lambda y = 0 \quad -\infty < x < \infty$$

with y and y' polynomially bounded at ∞ .

In the self-adjoint form:

$$(e^{-x^2/2} y')' + \lambda e^{-x^2/2} y = 0, \quad -\infty < x < \infty.$$

Eigenvalues: $\{\lambda_n = n^2\}_{n=0}^{\infty}$ and the corresponding eigenfunctions are the Hermite polynomials $\{H_n(x)\}_{n=0}^{\infty}$ with $H_0(x) = 1, H_1(x) = 2x, H_2(x) = 4x^2 - 2, \dots$

Here, $p(x) = e^{-x^2/2}$ and orthogonality property:

$$\int_{-\infty}^{\infty} e^{-x^2/2} H_m(x) H_n(x) dx = 0, \quad (m \neq n).$$

Example 4: Laguerre equation

In its standard form:

$$xy'' + (1-x)y' + \lambda y = 0 \quad 0 < x < \infty$$

with y and its derivative bounded at ∞ .

Example 4: Laguerre equation

In its standard form:

$$xy'' + (1-x)y' + \lambda y = 0 \quad 0 < x < \infty$$

with y and its derivative bounded at ∞ .

In the self-adjoint form:

Example 4: Laguerre equation

In its standard form:

$$xy'' + (1-x)y' + \lambda y = 0 \quad 0 < x < \infty$$

with y and its derivative bounded at ∞ .

In the self-adjoint form:

$$(xe^{-x}y')' + \lambda e^{-x}y = 0, \quad 0 < x < \infty.$$

Example 4: Laguerre equation

In its standard form:

$$xy'' + (1 - x)y' + \lambda y = 0 \quad 0 < x < \infty$$

with y and its derivative bounded at ∞ .

In the self-adjoint form:

$$(xe^{-x}y')' + \lambda e^{-x}y = 0, \quad 0 < x < \infty.$$

Eigenvalues: $\{\lambda_n = n\}_{n=0}^{\infty}$ and the corresponding eigenfunctions are the Laguerre polynomials $\{L_n(x)\}_{n=0}^{\infty}$ with

$$L_0(x) = 1, L_1(x) = 1 - x, L_2(x) = \frac{1}{2}(x^2 - 4x + 2), \dots$$

Here, $p(x) = e^{-x}$ and orthogonality property:

$$\int_0^{\infty} e^{-x} L_m(x) L_n(x) dx = 0, \quad (m \neq n).$$

Conservation Laws: Self Reading Hand out 2

Many PDE models come naturally from:

Conservation Principle

The rate at which a quantity of interest changes in a given domain = the rate at which the quantity flows across the boundary of the domain + the rate at which the quantity is created or destroyed within the domain.

For Air pollution model: Consider a tube of cross sectional area A

- $u = u(x, t)$ denotes the concentration of the pollutants.
Then, the amount of the concentration in a small section of width dx is given by $u(x, t)Adx$.

- $q = q(x, t)$ denotes the flux

Definition

Flux is defined as the amount of concentration crossing the section x at t and its unit is given by amount per unit area per unit time.

The amount of the concentration that is crossing at x and at t is $Aq(x, t)$. By convention, the flux is positive if the flow is from left to right and negative if it is from right to left.

- $f = f(x, t)$ denotes the rate at the quantity is created or destroyed within the section x and at time t . Then the amount is created or destroyed in a width dx per unit time is $f(x, t)Adx$.

Conservation Laws—

Formulate the conservation law by considering a fixed, but arbitrary section $a \leq x \leq b$ of the tube. By the Conservation principle:

Conservation Principle

$$\frac{d}{dt} \int_a^b u(x, t) A \, dx = Aq(a, t) - Aq(b, t) + \int_a^b f(x, t) A \, dx$$

Assuming u and q have continuous first derivatives, we rewrite as

$$\int_a^b \left(u_t(x, t) + q_x(x, t) - f(x, t) \right) dx = 0.$$

Since the section $[a, b]$ is arbitrary and the integrand is continuous, therefore,

$$u_t(x, t) + q_x(x, t) = f(x, t).$$

Here, there are two unknowns and one equation, therefore, a relation between the flux and the concentration, which should be given by a physical law. In air pollution model $q = \alpha u$.

Hence, we obtain

$$u_t + \alpha u_x = f(x, t).$$

Here, α is the wind velocity.

This is the mathematical model of air pollution (in the absence of diffusion). How does one solve it?