

Sequential Learning Algorithms

Samplers

D Manjunath & Jayakrishnan Nair

EE, IIT Bombay

January 8, 2022

Sampler 1: Predicting a Bernoulli sequence

- $X_t \in \{0, 1\}$ is a binary sequence for $t = 1, 2, \dots$
- Let $\{X_t\}$ be i.i.d. Bernoulli(p) sequence.
- Let \hat{X}_t be the prediction from the algorithm.
- Consider a randomized prediction, i.e., \hat{X}_t is a Bernoulli(α) random variable.
- **Claim 1:** If p were known, the ‘best’ algorithm, the one that minimizes the expected number of mistakes upto to time T , would be, for

$$\hat{X}_t = \begin{cases} 1 & \text{if } p \geq 0.5 \\ 0 & \text{otherwise} \end{cases}$$

$$\hat{X}_t = \begin{cases} 1 & \text{if } p \geq 0.5 \\ 0 & \text{otherwise} \end{cases}$$

Make fixed prediction of 1 if $p \geq 0.5$ and 0 otherwise.

Knowing the sequence X_1, \dots, X_{t-1} has no bearing on prediction for X_t .

What if p is not known? And we want to predict for

Sampler 1: Predicting a Bernoulli sequence

- $X_t \in \{0, 1\}$ is a binary sequence for $t = 1, 2, \dots$
- Let $\{X_t\}$ be i.i.d. Bernoulli(p) sequence.
- Let \hat{X}_t be the prediction from the algorithm.
- Consider a randomized prediction, i.e., \hat{X}_t is a Bernoulli(α) random variable.
- **Claim 1:** If p were known, the ‘best’ algorithm, the one that minimizes the expected number of mistakes upto to time T , would be,

$$\hat{X}_t = \begin{cases} 1 & \text{if } p \geq 0.5 \\ 0 & \text{otherwise} \end{cases}$$

Can we find prediction of 1 if $p \geq 0.5$ and 0 otherwise.

Knowing the sequence X_1, \dots, X_{t-1} has no bearing on prediction for X_t .
If we know p we know α . And we want to predict for

Sampler 1: Predicting a Bernoulli sequence

- $X_t \in \{0, 1\}$ is a binary sequence for $t = 1, 2, \dots$
- Let $\{X_t\}$ be i.i.d. Bernoulli(p) sequence.
- Let \hat{X}_t be the prediction from the algorithm.
- Consider a randomized prediction, i.e., \hat{X}_t is a Bernoulli(α) random variable.
- **Claim 1:** If p were known, the ‘best’ algorithm, the one that minimizes the expected number of mistakes upto to time T , would be,

$$\hat{X}_t = \begin{cases} 1 & \text{if } p \geq 0.5 \\ 0 & \text{otherwise} \end{cases}$$

Can we do better if $p \in [0, 0.5]$ and 0 otherwise.

Can we do better if $p \in [0.5, 1]$ and 1 otherwise.

Can we do better if $p \in [0, 1]$ and p otherwise.

Sampler 1: Predicting a Bernoulli sequence

- $X_t \in \{0, 1\}$ is a binary sequence for $t = 1, 2, \dots$
- Let $\{X_t\}$ be i.i.d. Bernoulli(p) sequence.
- Let \hat{X}_t be the prediction from the algorithm.
- Consider a randomized prediction, i.e., \hat{X}_t is a Bernoulli(α) random variable.
- **Claim 1:** If p were known, the ‘best’ algorithm, the one that minimizes the expected number of mistakes upto to time T , would be,

Sampler 1: Predicting a Bernoulli sequence

- $X_t \in \{0, 1\}$ is a binary sequence for $t = 1, 2, \dots$
- Let $\{X_t\}$ be i.i.d. Bernoulli(p) sequence.
- Let \hat{X}_t be the prediction from the algorithm.
- Consider a randomized prediction, i.e., \hat{X}_t is a Bernoulli(α) random variable.
- **Claim 1:** If p were known, the ‘best’ algorithm, the one that minimizes the expected number of mistakes upto to time T , would be, for $t = 1, \dots, T$,

$$\hat{X}_t = \mathcal{I}_{p \geq 0.5}$$

i.e., a fixed prediction of 1 if $p \geq 0.5$ and 0 otherwise.

Sampler 1: Predicting a Bernoulli sequence

- $X_t \in \{0, 1\}$ is a binary sequence for $t = 1, 2, \dots$
- Let $\{X_t\}$ be i.i.d. Bernoulli(p) sequence.
- Let \hat{X}_t be the prediction from the algorithm.
- Consider a randomized prediction, i.e., \hat{X}_t is a Bernoulli(α) random variable.
- **Claim 1:** If p were known, the ‘best’ algorithm, the one that minimizes the expected number of mistakes upto to time T , would be, for $t = 1, \dots, T$,

$$\hat{X}_t = \mathcal{I}_{p \geq 0.5}$$

i.e., a fixed prediction of 1 if $p \geq 0.5$ and 0 otherwise.

- Knowing the sequence X_1, \dots, X_{t-1} has no bearing on prediction for X_t .
- The best prediction algorithm would be a fixed one.

Sampler 1: Predicting a Bernoulli sequence

- $X_t \in \{0, 1\}$ is a binary sequence for $t = 1, 2, \dots$
- Let $\{X_t\}$ be i.i.d. Bernoulli(p) sequence.
- Let \hat{X}_t be the prediction from the algorithm.
- Consider a randomized prediction, i.e., \hat{X}_t is a Bernoulli(α) random variable.
- **Claim 1:** If p were known, the ‘best’ algorithm, the one that minimizes the expected number of mistakes upto to time T , would be, for $t = 1, \dots, T$,

$$\hat{X}_t = \mathcal{I}_{p \geq 0.5}$$

i.e., a fixed prediction of 1 if $p \geq 0.5$ and 0 otherwise.

- Knowing the sequence X_1, \dots, X_{t-1} has no bearing on prediction for X_t .
- What if p is not known? And we want to predict for

Sampler 1: Predicting a Bernoulli sequence

- $X_t \in \{0, 1\}$ is a binary sequence for $t = 1, 2, \dots$
- Let $\{X_t\}$ be i.i.d. Bernoulli(p) sequence.
- Let \hat{X}_t be the prediction from the algorithm.
- Consider a randomized prediction, i.e., \hat{X}_t is a Bernoulli(α) random variable.
- **Claim 1:** If p were known, the ‘best’ algorithm, the one that minimizes the expected number of mistakes upto to time T , would be, for $t = 1, \dots, T$,

$$\hat{X}_t = \mathcal{I}_{p \geq 0.5}$$

i.e., a fixed prediction of 1 if $p \geq 0.5$ and 0 otherwise.

- Knowing the sequence X_1, \dots, X_{t-1} has no bearing on prediction for X_t .
- What if p is not known? And we want to predict for

Sampler 1: Predicting a Bernoulli sequence

- $X_t \in \{0, 1\}$ is a binary sequence for $t = 1, 2, \dots$
- Let $\{X_t\}$ be i.i.d. Bernoulli(p) sequence.
- Let \hat{X}_t be the prediction from the algorithm.
- Consider a randomized prediction, i.e., \hat{X}_t is a Bernoulli(α) random variable.
- **Claim 1:** If p were known, the ‘best’ algorithm, the one that minimizes the expected number of mistakes upto to time T , would be, for $t = 1, \dots, T$,

$$\hat{X}_t = \mathcal{I}_{p \geq 0.5}$$

i.e., a fixed prediction of 1 if $p \geq 0.5$ and 0 otherwise.

- Knowing the sequence X_1, \dots, X_{t-1} has no bearing on prediction for X_t .
- What if p is not known? And we want to predict for

Predicting a Bernoulli sequence

- **Claim 2:** More generally, let \hat{X}_t be the prediction using the data X_1, \dots, X_{t-1} , from any algorithm. Then,

$$\Pr(\hat{X}_t \neq X_t \mid X_1, \dots, X_{t-1}) \geq \min\{p, (1-p)\}$$

i.e., a fixed prediction conditioned on the data has the minimum expected error.

Proof

Predicting a Berouilli sequence

- The last claim leads us to suggest the following *historical majority* algorithm (HMA) for prediction of a Benroulli sequence when p is unknown.

$$\hat{X}_t = \text{majority from } \{X_1, X_2, \dots, X_{t-1}\}$$

- The expected loss for this algorithm upto time T is

$$\sum_{t=1}^T \Pr(\hat{X}_t \neq X_t) - \min\{p, (1-p)\}$$

- The best error that any algorithm could have had is with the knowledge of p . It is the difference that is the loss.
- The number of mistakes that the HM algorithm would make up to time T is the number of times that X_t was not equal to the majority of $\{X_1, X_2, \dots, X_{t-1}\}$ for $t = 1, \dots, T$.

Predicting a Berouilli sequence

- The last claim leads us to suggest the following *historical majority* algorithm (HMA) for prediction of a Benroulli sequence when p is unknown.

$$\hat{X}_t = \text{majority from } \{X_1, X_2, \dots, X_{t-1}\}$$

- The expected loss for this algorithm upto time T is

$$\sum_{t=1}^T \Pr(\hat{X}_t \neq X_t) - \min\{p, (1-p)\}$$

- The best error that any algorithm could have had is with the knowledge of p . It is the difference that is the loss.
- The number of mistakes that the HM algorithm would make up to time T is the number of times that X_t was not equal to the majority of $\{X_1, X_2, \dots, X_{t-1}\}$ for $t = 1, \dots, T$.

Predicting a Berouilli sequence

- The last claim leads us to suggest the following *historical majority* algorithm (HMA) for prediction of a Benroulli sequence when p is unknown.

$$\hat{X}_t = \text{majority from } \{X_1, X_2, \dots, X_{t-1}\}$$

- The expected loss for this algorithm upto time T is

$$\sum_{t=1}^T \Pr(\hat{X}_t \neq X_t) - \min\{p, (1-p)\}$$

- The best error that any algorithm could have had is with the knowledge of p . It is the difference that is the loss.
- The number of mistakes that the HM algorithm would make up to time T is the number of times that X_t was not equal to the majority of $\{X_1, X_2, \dots, X_{t-1}\}$ for $t = 1, \dots, T$.

Predicting a Berouilli sequence

- The last claim leads us to suggest the following *historical majority* algorithm (HMA) for prediction of a Benroulli sequence when p is unknown.

$$\hat{X}_t = \text{majority from } \{X_1, X_2, \dots, X_{t-1}\}$$

- The expected loss for this algorithm upto time T is

$$\sum_{t=1}^T \Pr(\hat{X}_t \neq X_t) - \min\{p, (1-p)\}$$

- The best error that any algorithm could have had is with the knowledge of p . It is the difference that is the loss.
- The number of mistakes that the HM algorithm would make up to time T is the number of times that X_t was not equal to the majority of $\{X_1, X_2, \dots, X_{t-1}\}$ for $t = 1, \dots, T$.

Loss for the HMA

Worksheet

Loss for the HMA

- Without loss of generality, assume $p \geq 0.5$.
- Let L_t be the probability that the prediction for step t is different from the “full knowledge case”, i.e.,

$$L_t := \Pr(\hat{X}_t \neq 1)$$

- Clearly, L_t is the loss at step t .

$$L_t = \Pr\left(\sum_{k=1}^{t-1} X_k \leq \frac{t-1}{2}\right)$$

$$= \Pr\left(\sum_{k=1}^{t-1} (2X_k - 1) \leq -\frac{t-1}{2} + (t-1)p\right)$$

$$= \Pr\left(-1 \leq \frac{1}{t-1} \sum_{k=1}^{t-1} (2X_k - 1) \leq 2p - 1\right) \quad \text{Applying Chebyshev Inequality}$$

$$\leq \frac{1}{(2p-1)^2} + \frac{1}{(1-2p)^2}$$

Loss for the HMA

- Without loss of generality, assume $p \geq 0.5$.
- Let L_t be the probability that the prediction for step t is different from the “full knowledge case”, i.e.,

$$L_t := \Pr(\hat{X}_t \neq 1)$$

- Clearly, L_t is the loss at step t .

$$L_t = \Pr\left(\sum_{k=1}^{t-1} X_k \leq \frac{t-1}{2}\right)$$

$$= \Pr\left(\sum_{k=1}^{t-1} (2X_k - 1) \leq (t-1) - 2\sum_{k=1}^{t-1} X_k\right) \\ = \Pr\left(\sum_{k=1}^{t-1} (2X_k - 1) \leq (t-1) - 2\sum_{k=1}^{t-1} (1 - (2X_k - 1))\right) \\ = \Pr\left(\sum_{k=1}^{t-1} (2X_k - 1) \leq (t-1) - 2(t-1) + 2\sum_{k=1}^{t-1} (2X_k - 1)\right) \\ = \Pr\left(\sum_{k=1}^{t-1} (2X_k - 1) \leq - (t-1) + 2\sum_{k=1}^{t-1} (2X_k - 1)\right) \\ = \Pr\left(\sum_{k=1}^{t-1} (2X_k - 1) \geq (t-1)\right)$$

Loss for the HMA

- Without loss of generality, assume $p \geq 0.5$.
- Let L_t be the probability that the prediction for step t is different from the “full knowledge case”, i.e.,

$$L_t := \Pr(\hat{X}_t \neq 1)$$

- Clearly, L_t is the loss at step t .

$$\begin{aligned} L_t &= \Pr\left(\sum_{k=1}^{t-1} X_k \leq \frac{t-1}{2}\right) \\ &= \Pr\left(\sum_{k=1}^{t-1} (X_k - p) \leq \frac{t-1}{2} - (t-1)p\right) \\ &\leq \exp \frac{-2 \left(\frac{t-1}{2} - (t-1)p\right)^2}{t-1} \quad \text{Applying Hoeffding Inequality} \\ &= \exp -2(t-1)(0.5 - p)^2 \end{aligned}$$

Loss for the HMA

- Without loss of generality, assume $p \geq 0.5$.
- Let L_t be the probability that the prediction for step t is different from the “full knowledge case”, i.e.,

$$L_t := \Pr(\hat{X}_t \neq 1)$$

- Clearly, L_t is the loss at step t .

$$\begin{aligned} L_t &= \Pr\left(\sum_{k=1}^{t-1} X_k \leq \frac{t-1}{2}\right) \\ &= \Pr\left(\sum_{k=1}^{t-1} (X_k - p) \leq \frac{t-1}{2} - (t-1)p\right) \\ &\leq \exp \frac{-2 \left(\frac{t-1}{2} - (t-1)p\right)^2}{t-1} \quad \text{Applying Hoeffding Inequality} \\ &= \exp -2(t-1)(0.5 - p)^2 \end{aligned}$$

Loss for the HMA

- Without loss of generality, assume $p \geq 0.5$.
- Let L_t be the probability that the prediction for step t is different from the “full knowledge case”, i.e.,

$$L_t := \Pr(\hat{X}_t \neq 1)$$

- Clearly, L_t is the loss at step t .

$$\begin{aligned} L_t &= \Pr\left(\sum_{k=1}^{t-1} X_k \leq \frac{t-1}{2}\right) \\ &= \Pr\left(\sum_{k=1}^{t-1} (X_k - p) \leq \frac{t-1}{2} - (t-1)p\right) \\ &\leq \exp \frac{-2 \left(\frac{t-1}{2} - (t-1)p\right)^2}{t-1} \quad \text{Applying Hoeffding Inequality} \\ &= \exp -2(t-1)(0.5 - p)^2 \end{aligned}$$

Loss for the HMA

- Without loss of generality, assume $p \geq 0.5$.
- Let L_t be the probability that the prediction for step t is different from the “full knowledge case”, i.e.,

$$L_t := \Pr(\hat{X}_t \neq 1)$$

- Clearly, L_t is the loss at step t .

$$\begin{aligned} L_t &= \Pr\left(\sum_{k=1}^{t-1} X_k \leq \frac{t-1}{2}\right) \\ &= \Pr\left(\sum_{k=1}^{t-1} (X_k - p) \leq \frac{t-1}{2} - (t-1)p\right) \\ &\leq \exp \frac{-2 \left(\frac{t-1}{2} - (t-1)p\right)^2}{t-1} \quad \text{Applying Hoeffding Inequality} \\ &= \exp -2(t-1)(0.5 - p)^2 \end{aligned}$$

Loss for the HMA

- Without loss of generality, assume $p \geq 0.5$.
- Let L_t be the probability that the prediction for step t is different from the “full knowledge case”, i.e.,

$$L_t := \Pr(\hat{X}_t \neq 1)$$

- Clearly, L_t is the loss at step t .

$$\begin{aligned} L_t &= \Pr\left(\sum_{k=1}^{t-1} X_k \leq \frac{t-1}{2}\right) \\ &= \Pr\left(\sum_{k=1}^{t-1} (X_k - p) \leq \frac{t-1}{2} - (t-1)p\right) \\ &\leq \exp \frac{-2\left(\frac{t-1}{2} - (t-1)p\right)^2}{t-1} \quad \text{Applying Hoeffding Inequality} \\ &= \exp -2(t-1)(0.5 - p)^2 \end{aligned}$$

Loss for the HMA

- Without loss of generality, assume $p \geq 0.5$.
- Let L_t be the probability that the prediction for step t is different from the “full knowledge case”, i.e.,

$$L_t := \Pr(\hat{X}_t \neq 1)$$

- Clearly, L_t is the loss at step t .

$$\begin{aligned} L_t &= \Pr\left(\sum_{k=1}^{t-1} X_k \leq \frac{t-1}{2}\right) \\ &= \Pr\left(\sum_{k=1}^{t-1} (X_k - p) \leq \frac{t-1}{2} - (t-1)p\right) \\ &\leq \exp \frac{-2\left(\frac{t-1}{2} - (t-1)p\right)^2}{t-1} \quad \text{Applying Hoeffding Inequality} \\ &= \exp -2(t-1)(0.5 - p)^2 \end{aligned}$$

Loss for the HMA

- The cumulative regret in T steps is

$$\begin{aligned}\sum_{t=1}^T L_t &\leq \sum_{t=1}^{T-1} \exp -2(t-1)(0.5-p)^2 \\ &\leq \frac{1}{1 - \exp -2(0.5-p)^2}\end{aligned}$$

So, we observe that the regret is upper bounded by a constant!

Loss for the HMA

- The cumulative regret in T steps is

$$\begin{aligned}\sum_{t=1}^T L_t &\leq \sum_{t=1}^{T-1} \exp -2(t-1)(0.5-p)^2 \\ &\leq \frac{1}{1 - \exp -2(0.5-p)^2}\end{aligned}$$

- We thus see that the regret is upper bounded by a constant.

Loss for the HMA

- The cumulative regret in T steps is

$$\begin{aligned}\sum_{t=1}^T L_t &\leq \sum_{t=1}^{T-1} \exp -2(t-1)(0.5-p)^2 \\ &\leq \frac{1}{1 - \exp -2(0.5-p)^2}\end{aligned}$$

■ We thus see that the regret is upper bounded by a constant.

Loss for the HMA

- The cumulative regret in T steps is

$$\begin{aligned}\sum_{t=1}^T L_t &\leq \sum_{t=1}^{T-1} \exp -2(t-1)(0.5-p)^2 \\ &\leq \frac{1}{1 - \exp -2(0.5-p)^2}\end{aligned}$$

- We thus see that the regret is upper bounded by a constant.

Sampler 2: Predicting a “fixed point”

- Consider the following shooting game: You are shooting into a unit square and you have to “predict” the fixed point, say \hat{X}_t for the t -th attempt.
- Let X_t be the actual location of the hit.
- For every \hat{X}_t there is a cost $\|\hat{X}_t - X_t\|^2$.
- First consider the offline problem, or the one shot problem: the sequence X_t for $t = 1, \dots, T$, is available to you and you have to choose the *best* fixed point

$$X_T^* := \arg \min_{x \in [0,1]^2} \sum_{t=1}^T \|X_t - x\|^2$$

- Question: The optimal fixed point for X would be

$$X_T^* = \frac{1}{T} \sum_{t=1}^T X_t$$

-

Sampler 2: Predicting a “fixed point”

- Consider the following shooting game: You are shooting into a unit square and you have to “predict” the fixed point, say \hat{X}_t for the t -th attempt.
- Let X_t be the actual location of the hit.
- For every \hat{X}_t there is a cost $\|\hat{X}_t - X_t\|^2$.
- First consider the offline problem, or the one shot problem: the sequence X_t for $t = 1, \dots, T$, is available to you and you have to choose the *best* fixed point

$$X_T^* := \arg \min_{x \in [0,1]^2} \sum_{t=1}^T \|X_t - x\|^2$$

Sampler 2: Predicting a “fixed point”

- Consider the following shooting game: You are shooting into a unit square and you have to “predict” the fixed point, say \hat{X}_t for the t -th attempt.
- Let X_t be the actual location of the hit.
- For every \hat{X}_t there is a cost $\|\hat{X}_t - X_t\|^2$.
- First consider the offline problem, or the one shot problem: the sequence X_t for $t = 1, \dots, T$, is available to you and you have to choose the *best* fixed point

$$X_T^* := \arg \min_{x \in [0,1]^2} \sum_{t=1}^T \|X_t - x\|^2$$

Sampler 2: Predicting a “fixed point”

- Consider the following shooting game: You are shooting into a unit square and you have to “predict” the fixed point, say \hat{X}_t for the t -th attempt.
- Let X_t be the actual location of the hit.
- For every \hat{X}_t there is a cost $\|\hat{X}_t - X_t\|^2$.
- First consider the offline problem, or the one shot problem: the sequence X_t for $t = 1, \dots, T$, is available to you and you have to choose the *best* fixed point

$$X_T^* := \arg \min_{x \in [0,1]^2} \sum_{t=1}^T \|X_t - x\|^2$$

■ Claim: The explicit formula for X_T^* would be

$$X_T^* = \frac{1}{T} \sum_{t=1}^T X_t$$

Sampler 2: Predicting a “fixed point”

- Consider the following shooting game: You are shooting into a unit square and you have to “predict” the fixed point, say \hat{X}_t for the t -th attempt.
- Let X_t be the actual location of the hit.
- For every \hat{X}_t there is a cost $\|\hat{X}_t - X_t\|^2$.
- First consider the offline problem, or the one shot problem: the sequence X_t for $t = 1, \dots, T$, is available to you and you have to choose the *best* fixed point

$$X_T^* := \arg \min_{x \in [0,1]^2} \sum_{t=1}^T \|X_t - x\|^2$$

■ Claim: The explicit formula for X_T^* would be

$$X_T^* = \frac{1}{T} \sum_{t=1}^T X_t$$

■ $\hat{X}_t = X_T^*$ minimizes the total cost.

Sampler 2: Predicting a “fixed point”

- Consider the following shooting game: You are shooting into a unit square and you have to “predict” the fixed point, say \hat{X}_t for the t -th attempt.
- Let X_t be the actual location of the hit.
- For every \hat{X}_t there is a cost $\|\hat{X}_t - X_t\|^2$.
- First consider the offline problem, or the one shot problem: the sequence X_t for $t = 1, \dots, T$, is available to you and you have to choose the *best* fixed point

$$X_T^* := \arg \min_{x \in [0,1]^2} \sum_{t=1}^T \|X_t - x\|^2$$

- **Claim:** The explicit formula for X_T^* would be

$$X_T^* = \frac{1}{T} \sum_{t=1}^T X_t$$

- $\hat{X}_t = X_T^*$ minimizes the total cost.

Sampler 2: Predicting a “fixed point”

- Consider the following shooting game: You are shooting into a unit square and you have to “predict” the fixed point, say \hat{X}_t for the t -th attempt.
- Let X_t be the actual location of the hit.
- For every \hat{X}_t there is a cost $\|\hat{X}_t - X_t\|^2$.
- First consider the offline problem, or the one shot problem: the sequence X_t for $t = 1, \dots, T$, is available to you and you have to choose the *best* fixed point

$$X_T^* := \arg \min_{x \in [0,1]^2} \sum_{t=1}^T \|X_t - x\|^2$$

- **Claim:** The explicit formula for X_T^* would be

$$X_T^* = \frac{1}{T} \sum_{t=1}^T X_t$$

- $\hat{X}_t = X_T^*$ minimizes the total cost.

Sampler 2: Predicting a “fixed point”

- Now convert this to an online question: Predict the point after every shot such that the cumulative error is minimized.
- Specifically,

$$\min \sum_{t=1} \|\hat{X}_t - X_t\|^2$$

with \hat{X}_t to be determined before the t -th shot and can be done using the information X_1, \dots, X_{t-1} .

- Consider the following algorithm

$$\hat{X}_t = X_{t-1}^*$$

where X_{t-1}^* is the point in the previous shot which was closest to the goal. Specifically, we can see that the cumulative loss in the online algorithm is bounded by the full information offline error.

Sampler 2: Predicting a “fixed point”

- Now convert this to an online question: Predict the point after every shot such that the cumulative error is minimized.
- Specifically,

$$\min \sum_{t=1} \|\hat{X}_t - X_t\|^2$$

with \hat{X}_t to be determined before the t -th shot and can be done using the information X_1, \dots, X_{t-1} .

- Consider the following algorithm

$$\hat{X}_t = X_{t-1}^*$$

Sampler 2: Predicting a “fixed point”

- Now convert this to an online question: Predict the point after every shot such that the cumulative error is minimized.
- Specifically,

$$\min \sum_{t=1} \|\hat{X}_t - X_t\|^2$$

with \hat{X}_t to be determined before the t -th shot and can be done using the information X_1, \dots, X_{t-1} .

- Consider the following algorithm

$$\hat{X}_t = X_{t-1}^* = \arg \min_{x \in [0,1]^2} \sum_{s=1}^{t-1} \|X_s - x\|^2 = \frac{1}{t-1} \sum_{s=1}^{t-1} X_s$$

Sampler 2: Predicting a “fixed point”

- Now convert this to an online question: Predict the point after every shot such that the cumulative error is minimized.
- Specifically,

$$\min \sum_{t=1} \|\hat{X}_t - X_t\|^2$$

with \hat{X}_t to be determined before the t -th shot and can be done using the information X_1, \dots, X_{t-1} .

- Consider the following algorithm

$$\hat{X}_t = X_{t-1}^* = \arg \min_{x \in [0,1]^2} \sum_{s=1}^{t-1} \|X_s - x\|^2 = \frac{1}{t-1} \sum_{s=1}^{t-1} X_s$$

- How does this compare with the “offline” algorithm above.
- How does this compare with the “online” algorithm above.
- How does this compare with the “fixed point” algorithm above.

Sampler 2: Predicting a “fixed point”

- Now convert this to an online question: Predict the point after every shot such that the cumulative error is minimized.
- Specifically,

$$\min \sum_{t=1} \|\hat{X}_t - X_t\|^2$$

with \hat{X}_t to be determined before the t -th shot and can be done using the information X_1, \dots, X_{t-1} .

- Consider the following algorithm

$$\hat{X}_t = X_{t-1}^* = \arg \min_{X \in [0,1]^2} \sum_{s=1}^{t-1} \|X_s - x\|^2 = \frac{1}{t-1} \sum_{s=1}^{t-1} X_s$$

- How does this compare with the “offline” algorithm above.

Sampler 2: Predicting a “fixed point”

- Now convert this to an online question: Predict the point after every shot such that the cumulative error is minimized.
- Specifically,

$$\min \sum_{t=1} \|\hat{X}_t - X_t\|^2$$

with \hat{X}_t to be determined before the t -th shot and can be done using the information X_1, \dots, X_{t-1} .

- Consider the following algorithm

$$\hat{X}_t = X_{t-1}^* = \arg \min_{x \in [0,1]^2} \sum_{s=1}^{t-1} \|X_s - x\|^2 = \frac{1}{t-1} \sum_{s=1}^{t-1} X_s$$

- How does this compare with the “offline” algorithm above.

Specifically, what is the cumulative *loss* in the online algorithm compared to the full information offline case, i.e.,

$$\sum_{t=1}^T \|\hat{X}_t - X_t\|^2 - \sum_{t=1}^T \|\hat{X}_T - X_t\|^2 = \sum_{t=1}^T \|X_{t-1}^* - X_t\|^2 - \sum_{t=1}^T \|X_T^* - X_t\|^2$$

Sampler 2: Predicting a “fixed point”

- Now convert this to an online question: Predict the point after every shot such that the cumulative error is minimized.
- Specifically,

$$\min \sum_{t=1} \|\hat{X}_t - X_t\|^2$$

with \hat{X}_t to be determined before the t -th shot and can be done using the information X_1, \dots, X_{t-1} .

- Consider the following algorithm

$$\hat{X}_t = X_{t-1}^* = \arg \min_{x \in [0,1]^2} \sum_{s=1}^{t-1} \|X_s - x\|^2 = \frac{1}{t-1} \sum_{s=1}^{t-1} X_s$$

- How does this compare with the “offline” algorithm above.

Specifically, what is the cumulative *loss* in the online algorithm compared to the full information offline case, i.e.,

$$\sum_{t=1}^T \|\hat{X}_t - X_t\|^2 - \sum_{t=1}^T \|\hat{X}_T - X_t\|^2 = \sum_{t=1}^T \|X_{t-1}^* - X_t\|^2 - \sum_{t=1}^T \|X_T^* - X_t\|^2$$

Sampler 2: Predicting a “fixed point”

- Now convert this to an online question: Predict the point after every shot such that the cumulative error is minimized.
- Specifically,

$$\min \sum_{t=1} \|\hat{X}_t - X_t\|^2$$

with \hat{X}_t to be determined before the t -th shot and can be done using the information X_1, \dots, X_{t-1} .

- Consider the following algorithm

$$\hat{X}_t = X_{t-1}^* = \arg \min_{x \in [0,1]^2} \sum_{s=1}^{t-1} \|X_s - x\|^2 = \frac{1}{t-1} \sum_{s=1}^{t-1} X_s$$

- How does this compare with the “offline” algorithm above. Specifically, what is the cumulative *loss* in the online algorithm compared to the full information offline case, i.e.,

$$\sum_{t=1}^T \|\hat{X}_t - X_t\|^2 - \sum_{t=1}^T \|\hat{X}_T - X_t\|^2 = \sum_{t=1}^T \|X_{t-1}^* - X_t\|^2 - \sum_{t=1}^T \|X_T^* - X_t\|^2$$

Analyzing Follow the Leader (FTL)

Lemma 1

$$\sum_{t=1}^T \|X_t^* - X_t\|^2 \leq \sum_{t=1}^T \|X_T^* - X_t\|^2$$

Proof

- Proof is by induction;
- Trivially true for $T = 1$ with equality;
- Inductive hypothesis: For $T > 1$,

$$\sum_{t=1}^T \|X_t^* - X_t\|^2 \leq \sum_{t=1}^T \|X_T^* - X_t\|^2$$

- We will now prove

$$\sum_{t=1}^T \|X_t^* - X_t\|^2 \leq \sum_{t=1}^T \|X_T^* - X_t\|^2$$

Analyzing Follow the Leader (FTL)

Lemma 1

$$\sum_{t=1}^T \|X_t^* - X_t\|^2 \leq \sum_{t=1}^T \|X_T^* - X_t\|^2$$

Proof

- Proof is by induction;
- Trivially true for $T = 1$ with equality.
- Inductive hypothesis: For $T > 1$,

$$\sum_{t=1}^T \|X_t^* - X_t\|^2 \leq \sum_{t=1}^T \|X_T^* - X_t\|^2$$

- We will now prove

$$\sum_{t=1}^T \|X_t^* - X_t\|^2 \leq \sum_{t=1}^T \|X_T^* - X_t\|^2$$

Analyzing Follow the Leader (FTL)

Lemma 1

$$\sum_{t=1}^T \|X_t^* - X_t\|^2 \leq \sum_{t=1}^T \|X_T^* - X_t\|^2$$

Proof

- Proof is by induction;
- Trivially true for $T = 1$ with equality.
- Induction hypothesis: For $T > 1$,

$$\sum_{t=1}^{T-1} \|X_t^* - X_t\|^2 \leq \sum_{t=1}^{T-1} \|X_{T-1}^* - X_t\|^2$$

- We will now prove

$$\sum_{t=1}^T \|X_t^* - X_t\|^2 \leq \sum_{t=1}^T \|X_T^* - X_t\|^2$$

Analyzing Follow the Leader (FTL)

Lemma 1

$$\sum_{t=1}^T \|X_t^* - X_t\|^2 \leq \sum_{t=1}^T \|X_T^* - X_t\|^2$$

Proof

- Proof is by induction;
- Trivially true for $T = 1$ with equality.
- Induction hypothesis: For $T > 1$,

$$\sum_{t=1}^{T-1} \|X_t^* - X_t\|^2 \leq \sum_{t=1}^{T-1} \|X_{T-1}^* - X_t\|^2$$

- We will now prove

$$\sum_{t=1}^T \|X_t^* - X_t\|^2 \leq \sum_{t=1}^T \|X_T^* - X_t\|^2$$

Analyzing Follow the Leader (FTL)

Lemma 1

$$\sum_{t=1}^T \|X_t^* - X_t\|^2 \leq \sum_{t=1}^T \|X_T^* - X_t\|^2$$

Proof

- Proof is by induction;
- Trivially true for $T = 1$ with equality.
- Induction hypothesis: For $T > 1$,

$$\sum_{t=1}^{T-1} \|X_t^* - X_t\|^2 \leq \sum_{t=1}^{T-1} \|X_{T-1}^* - X_t\|^2$$

- We will now prove

$$\sum_{t=1}^T \|X_t^* - X_t\|^2 \leq \sum_{t=1}^T \|X_T^* - X_t\|^2$$

Analyzing Follow the Leader (FTL)

Lemma 1

$$\sum_{t=1}^T \|X_t^* - X_t\|^2 \leq \sum_{t=1}^T \|X_T^* - X_t\|^2$$

Proof

- Proof is by induction;
- Trivially true for $T = 1$ with equality.
- Induction hypothesis: For $T > 1$,

$$\sum_{t=1}^{T-1} \|X_t^* - X_t\|^2 \leq \sum_{t=1}^{T-1} \|X_{T-1}^* - X_t\|^2$$

- We will now prove

$$\sum_{t=1}^T \|X_t^* - X_t\|^2 \leq \sum_{t=1}^T \|X_T^* - X_t\|^2$$

Analyzing Follow the Leader (FTL)

Lemma 1

$$\begin{aligned}\sum_{t=1}^T \|X_t^* - X_t\|^2 &\leq \sum_{t=1}^T \|X_T^* - X_t\|^2 \\ \Rightarrow \sum_{t=1}^{T-1} \|X_t^* - X_t\|^2 &\leq \sum_{t=1}^{T-1} \|X_T^* - X_t\|^2 \text{ knocking off term for } t = T\end{aligned}$$

We now prove the preceding inequality.

$$\begin{aligned}\sum_{t=1}^{T-1} \|X_t^* - X_t\|^2 &\leq \sum_{t=1}^{T-1} \|X_{T-1}^* - X_t\|^2 \quad \text{from induction hypothesis} \\ &\leq \sum_{t=1}^{T-1} \|X_T^* - X_t\|^2 \quad \text{because } X_{T-1}^* \text{ is not the minimizer} \quad \square\end{aligned}$$

Analyzing Follow the Leader (FTL)

Lemma 1

$$\begin{aligned}\sum_{t=1}^T \|X_t^* - X_t\|^2 &\leq \sum_{t=1}^T \|X_T^* - X_t\|^2 \\ \Rightarrow \sum_{t=1}^{T-1} \|X_t^* - X_t\|^2 &\leq \sum_{t=1}^{T-1} \|X_T^* - X_t\|^2 \text{ knocking of term for } t = T\end{aligned}$$

We now prove the preceding inequality.

$$\begin{aligned}\sum_{t=1}^{T-1} \|X_t^* - X_t\|^2 &\leq \sum_{t=1}^{T-1} \|X_{T-1}^* - X_t\|^2 \quad \text{from induction hypothesis} \\ &\leq \sum_{t=1}^{T-1} \|X_T^* - X_t\|^2 \quad \text{because } X_{T-1}^* \text{ is not the minimizer} \quad \square\end{aligned}$$

Analyzing Follow the Leader (FTL)

Lemma 1

$$\begin{aligned}\sum_{t=1}^T \|X_t^* - X_t\|^2 &\leq \sum_{t=1}^T \|X_T^* - X_t\|^2 \\ \Rightarrow \sum_{t=1}^{T-1} \|X_t^* - X_t\|^2 &\leq \sum_{t=1}^{T-1} \|X_T^* - X_t\|^2 \text{ knocking of term for } t = T\end{aligned}$$

We now prove the preceding inequality.

$$\begin{aligned}\sum_{t=1}^{T-1} \|X_t^* - X_t\|^2 &\leq \sum_{t=1}^{T-1} \|X_{T-1}^* - X_t\|^2 \quad \text{from induction hypothesis} \\ &\leq \sum_{t=1}^{T-1} \|X_T^* - X_t\|^2 \quad \text{because } X_{T-1}^* \text{ is not the minimizer} \quad \square\end{aligned}$$

Analyzing Follow the Leader (FTL)

Lemma 1

$$\begin{aligned}\sum_{t=1}^T \|X_t^* - X_t\|^2 &\leq \sum_{t=1}^T \|X_T^* - X_t\|^2 \\ \Rightarrow \sum_{t=1}^{T-1} \|X_t^* - X_t\|^2 &\leq \sum_{t=1}^{T-1} \|X_T^* - X_t\|^2 \text{ knocking of term for } t = T\end{aligned}$$

We now prove the preceding inequality.

$$\begin{aligned}\sum_{t=1}^{T-1} \|X_t^* - X_t\|^2 &\leq \sum_{t=1}^{T-1} \|X_{T-1}^* - X_t\|^2 \quad \text{from induction hypothesis} \\ &\leq \sum_{t=1}^{T-1} \|X_T^* - X_t\|^2 \quad \text{because } X_{T-1}^* \text{ is not the minimizer} \quad \square\end{aligned}$$

Analyzing Follow the Leader (FTL)

Lemma 1

$$\begin{aligned}\sum_{t=1}^T \|X_t^* - X_t\|^2 &\leq \sum_{t=1}^T \|X_T^* - X_t\|^2 \\ \Rightarrow \sum_{t=1}^{T-1} \|X_t^* - X_t\|^2 &\leq \sum_{t=1}^{T-1} \|X_T^* - X_t\|^2 \text{ knocking of term for } t = T\end{aligned}$$

We now prove the preceding inequality.

$$\begin{aligned}\sum_{t=1}^{T-1} \|X_t^* - X_t\|^2 &\leq \sum_{t=1}^{T-1} \|X_{T-1}^* - X_t\|^2 \quad \text{from induction hypothesis} \\ &\leq \sum_{t=1}^{T-1} \|X_T^* - X_t\|^2 \quad \text{because } X_{T-1}^* \text{ is not the minimizer} \quad \square\end{aligned}$$

Analyzing Follow the Leader (FTL)

Lemma 1

$$\begin{aligned}\sum_{t=1}^T \|X_t^* - X_t\|^2 &\leq \sum_{t=1}^T \|X_T^* - X_t\|^2 \\ \Rightarrow \sum_{t=1}^{T-1} \|X_t^* - X_t\|^2 &\leq \sum_{t=1}^{T-1} \|X_T^* - X_t\|^2 \text{ knocking of term for } t = T\end{aligned}$$

We now prove the preceding inequality.

$$\begin{aligned}\sum_{t=1}^{T-1} \|X_t^* - X_t\|^2 &\leq \sum_{t=1}^{T-1} \|X_{T-1}^* - X_t\|^2 \quad \text{from induction hypothesis} \\ &\leq \sum_{t=1}^{T-1} \|X_T^* - X_t\|^2 \quad \text{because } X_T^* \text{ is not the minimizer} \quad \square\end{aligned}$$

Analyzing Follow the Leader (FTL)

$$\begin{aligned} L_T &:= \sum_{t=1}^T \|\hat{X}_t - X_t\|^2 - \sum_{t=1}^T \|\hat{X}_T^* - X_t\|^2 \\ &= \sum_{t=1}^T \|X_{t-1}^* - X_t\|^2 - \sum_{t=1}^T \|\hat{X}_T^* - X_t\|^2 \\ &\leq \sum_{t=1}^T \|X_{t-1}^* - X_t\|^2 - \sum_{t=1}^T \|\hat{X}_t^* - X_t\|^2 \quad \text{by Lemma 1 above} \\ &= \sum_{t=1}^T \langle X_{t-1}^* - X_t^*, X_{t-1}^* + X_t^* - 2X_t \rangle \quad \text{using } \|x\|^2 - \|y\|^2 = \langle x+y, x-y \rangle \end{aligned}$$

Analyzing Follow the Leader (FTL)

$$\begin{aligned} L_T &:= \sum_{t=1}^T \|\hat{X}_t - X_t\|^2 - \sum_{t=1}^T \|\hat{X}_T^* - X_t\|^2 \\ &= \sum_{t=1}^T \|X_{t-1}^* - X_t\|^2 - \sum_{t=1}^T \|\hat{X}_T^* - X_t\|^2 \\ &\leq \sum_{t=1}^T \|X_{t-1}^* - X_t\|^2 - \sum_{t=1}^T \|\hat{X}_t^* - X_t\|^2 \quad \text{by Lemma 1 above} \\ &= \sum_{t=1}^T \langle X_{t-1}^* - X_t^*, X_{t-1}^* + X_t^* - 2X_t \rangle \quad \text{using } \|x\|^2 - \|y\|^2 = \langle x+y, x-y \rangle \end{aligned}$$

Analyzing Follow the Leader (FTL)

$$\begin{aligned} L_T &:= \sum_{t=1}^T \|\hat{X}_t - X_t\|^2 - \sum_{t=1}^T \|\hat{X}_T^* - X_t\|^2 \\ &= \sum_{t=1}^T \|X_{t-1}^* - X_t\|^2 - \sum_{t=1}^T \|\hat{X}_T^* - X_t\|^2 \\ &\leq \sum_{t=1}^T \|X_{t-1}^* - X_t\|^2 - \sum_{t=1}^T \|\hat{X}_t^* - X_t\|^2 \quad \text{by Lemma 1 above} \\ &= \sum_{t=1}^T \langle X_{t-1}^* - X_t^*, X_{t-1}^* + X_t^* - 2X_t \rangle \quad \text{using } \|x\|^2 - \|y\|^2 = \langle x+y, x-y \rangle \end{aligned}$$

Analyzing Follow the Leader (FTL)

$$\begin{aligned} L_T &:= \sum_{t=1}^T \|\hat{X}_t - X_t\|^2 - \sum_{t=1}^T \|\hat{X}_T^* - X_t\|^2 \\ &= \sum_{t=1}^T \|X_{t-1}^* - X_t\|^2 - \sum_{t=1}^T \|\hat{X}_T^* - X_t\|^2 \\ &\leq \sum_{t=1}^T \|X_{t-1}^* - X_t\|^2 - \sum_{t=1}^T \|\hat{X}_t^* - X_t\|^2 \quad \text{by Lemma 1 above} \\ &= \sum_{t=1}^T \langle X_{t-1}^* - X_t^*, X_{t-1}^* + X_t^* - 2X_t \rangle \quad \text{using } \|x\|^2 - \|y\|^2 = \langle x+y, x-y \rangle \end{aligned}$$

Analyzing Follow the Leader (FTL)

$$L_T \leq \sum_{t=1}^T \|X_{t-1}^* - X_t^*\| \cdot \|X_{t-1}^* + X_t^* - 2X_t\|$$

using Cauchy-Schwartz inequality $|\langle x, y \rangle| \leq \|x\| \cdot \|y\|$.

$$\leq \sum_{t=1}^T \|X_{t-1}^* - X_t^*\| \cdot (\|X_{t-1}^*\| + \|X_t^*\| + \|2X_t\|) \quad \text{using triangle inequality}$$

$$\leq \sum_{t=1}^T 4 \|X_{t-1}^* - X_t^*\| \quad \text{using the upper bound for } \|\cdot\| \text{ in 2nd term}$$

Use the formula for x_t^* , triangle inequality and upper bound for $\|\cdot\|$

$$\begin{aligned} \|X_{t-1}^* - X_t^*\| &= \left\| X_{t-1}^* - \frac{(t-1)X_{t-1}^* + X_t}{t} \right\| = \frac{1}{t} \|X_{t-1}^* - X_t\| \\ &= \frac{1}{t} (\|X_{t-1}^*\| + \|X_t\|) \leq \frac{2}{t} \end{aligned}$$

Analyzing Follow the Leader (FTL)

$$\begin{aligned} L_T &\leq \sum_{t=1}^T \|X_{t-1}^* - X_t^*\| \cdot \|X_{t-1}^* + X_t^* - 2X_t\| \\ &\quad \text{using Cauchy-Schwartz inequality } |\langle x, y \rangle| \leq \|x\| \cdot \|y\|. \\ &\leq \sum_{t=1}^T \|X_{t-1}^* - X_t^*\| \cdot (\|X_{t-1}^*\| + \|X_t^*\| + \|2X_t\|) \quad \text{using triangle inequality} \\ &\leq \sum_{t=1}^T 4 \|X_{t-1}^* - X_t^*\| \quad \text{using the upper bound for } \|\cdot\| \text{ in 2nd term} \end{aligned}$$

Use the formula for x_t^* , triangle inequality and upper bound for $\|\cdot\|$

$$\begin{aligned} \|X_{t-1}^* - X_t^*\| &= \left\| X_{t-1}^* - \frac{(t-1)X_{t-1}^* + X_t}{t} \right\| = \frac{1}{t} \|X_{t-1}^* - X_t\| \\ &= \frac{1}{t} (\|X_{t-1}^*\| + \|X_t\|) \leq \frac{2}{t} \end{aligned}$$

Analyzing Follow the Leader (FTL)

$$\begin{aligned} L_T &\leq \sum_{t=1}^T \|X_{t-1}^* - X_t^*\| \cdot \|X_{t-1}^* + X_t^* - 2X_t\| \\ &\quad \text{using Cauchy-Schwartz inequality } |\langle x, y \rangle| \leq \|x\| \cdot \|y\|. \\ &\leq \sum_{t=1}^T \|X_{t-1}^* - X_t^*\| \cdot (\|X_{t-1}^*\| + \|X_t^*\| + \|2X_t\|) \quad \text{using triangle inequality} \\ &\leq \sum_{t=1}^T 4 \|X_{t-1}^* - X_t^*\| \quad \text{using the upper bound for } \|\cdot\| \text{ in 2nd term} \end{aligned}$$

Use the formula for x_t^* , triangle inequality and upper bound for $\|\cdot\|$

$$\begin{aligned} \|X_{t-1}^* - X_t^*\| &= \left\| X_{t-1}^* - \frac{(t-1)X_{t-1}^* + X_t}{t} \right\| = \frac{1}{t} \|X_{t-1}^* - X_t\| \\ &= \frac{1}{t} (\|X_{t-1}^*\| + \|X_t\|) \leq \frac{2}{t} \end{aligned}$$

Analyzing Follow the Leader (FTL)

$$\begin{aligned} L_T &\leq \sum_{t=1}^T \|X_{t-1}^* - X_t^*\| \cdot \|X_{t-1}^* + X_t^* - 2X_t\| \\ &\quad \text{using Cauchy-Schwartz inequality } |\langle x, y \rangle| \leq \|x\| \cdot \|y\|. \\ &\leq \sum_{t=1}^T \|X_{t-1}^* - X_t^*\| \cdot (\|X_{t-1}^*\| + \|X_t^*\| + \|2X_t\|) \quad \text{using triangle inequality} \\ &\leq \sum_{t=1}^T 4 \|X_{t-1}^* - X_t^*\| \quad \text{using the upper bound for } \|\cdot\| \text{ in 2nd term} \end{aligned}$$

Use the formula for x_t^* , triangle inequality and upper bound for $\|\cdot\|$

$$\begin{aligned} \|X_{t-1}^* - X_t^*\| &= \left\| X_{t-1}^* - \frac{(t-1)X_{t-1}^* + X_t}{t} \right\| = \frac{1}{t} \|X_{t-1}^* - X_t\| \\ &= \frac{1}{t} (\|X_{t-1}^*\| + \|X_t\|) \leq \frac{2}{t} \end{aligned}$$

Analyzing Follow the Leader (FTL)

$$R_T \leq 4 \sum_{t=1}^T \frac{8}{T} \leq 8(1 + \ln(T))$$

Theorem For any sequence X_1, \dots, X_T , in the unit circle the FTL algorithm has a loss of at most $(1 + 8 \ln(T))$.

Analyzing Follow the Leader (FTL)

$$R_T \leq 4 \sum_{t=1}^T \frac{8}{T} \leq 8(1 + \ln(T))$$

Theorem For any sequence X_1, \dots, X_T , in the unit circle the FTL algorithm has a loss of at most $(1 + 8 \ln(T))$.