

CIS580 Problem Set 7

Sheil Sarda <sheils@seas.upenn.edu>

CIS580 Spring 2021

Contents

1	Scale Invariant Detection	1
1.1	Laplacian of Gaussian	1
1.2	Approximating a LoG by a DoG	2
1.2.1	Difference of Gaussians	2
1.2.2	DoG gets closer to LoG as $k \rightarrow 1$	3
1.2.3	Dropping the normalizng factor	3

1 Scale Invariant Detection

1.1 Laplacian of Gaussian




Figure 1

1.2 Approximating a LoG by a DoG

1.2.1 Difference of Gaussians




Figure 2

The plots obtained indicate that as k increases from $1 \rightarrow 2$, the Difference of Gaussians approximation is scaled down in magnitude.




Figure 3

1.2.2 DoG gets closer to LoG as $k \rightarrow 1$

We expect this to happen because in the provided expression, as $k \rightarrow 1$, the LoG will approach the DoG.

$$\text{LoG}_\sigma = \frac{1}{(k-1)\sigma^2} \text{DoG}_\sigma$$

1.2.3 Dropping the normalizing factor

We intentionally forget the normalizing factor and just use the difference of Gaussians because the product of a scalar and a Normal distribution will simply apply the constant as a scaling factor to both terms in the DoG expression.