

# CIS580 Problem Set 6


Sheil Sarda <sheils@seas.upenn.edu>

CIS580 Spring 2021


## Contents

1	Convolution of image with a Gaussian	1
2	Convolution of Gaussians	1
3	Convolution of Step Edge with Gaussian derivative	2
4	Box Function	3
5	1D FFT Quiz	4
6	2D Fourier Transform	5
7	Filter Design	6


# 1 Convolution of image with a Gaussian



(a) Original Image



(b)  $\sigma = 1$



(c)  $\sigma = 0.1$

Figure 1: Three simple graphs

The difference between the three images included above is that by increasing  $\sigma$  from  $\sigma = 0$  (original image) to 0.1 and 1, we increase the Gaussian blur applied to the image. This blurring reduces the image noise and detail.

## 2 Convolution of Gaussians

We know that the 1-D Gaussian  $g(t)$  is:

$$g(t) = \frac{1}{\sigma\sqrt{2\pi}} e^{-\frac{t^2}{2\sigma^2}}$$

Convolving two Gaussian kernels with  $\sigma_1$  and  $\sigma_2$  standard deviations:

$$g_1(t) = \frac{1}{\sigma_1\sqrt{2\pi}} e^{-\frac{t^2}{2\sigma_1^2}}$$

$$g_2(t) = \frac{1}{\sigma_2\sqrt{2\pi}} e^{-\frac{t^2}{2\sigma_2^2}}$$

$$(g_1 * g_2)(t) = \int_{-\infty}^{\infty} 2$$

### 3 Convolution of Step Edge with Gaussian derivative




Figure 2

## 4 Box Function




Figure 3

## 5 1D FFT Quiz




Figure 4

Function	FFT Plot
1	D
2	C
3	B
4	E
5	A

My reasoning for the above matching is that in frequency domain, the distance between the two peaks is proportional to the frequency of the curves.

## 6 2D Fourier Transform




Figure 5: Functions and corresponding FFTs

Function	FFT Plot
A	F
B	G
C	H
D	E

My reasoning for the above matching is that the separation of the white dots in the frequency domain is proportional to the frequency of the black and white signals. This reasoning is similar to the 1-dimensional case where the distance between the two peaks in the frequency domain was proportional to the frequency of the curves.

## **7 Filter Design**